

| • | | |
|---|--|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Digitized by the Internet Archive in 2017 with funding from CLIR

https://archive.org/details/speciesaccountsg00even

| | ÷ | | |
|---|---|--|--|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| • | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |




Val.6

Evenden, Fred G.

species index

Gavindae to Picifornes



Observer Fred Lvenden, Jr.

Tumber or field traps on much observed:

Lesser Loon 57-70-141-193-439-441-567-651-775-984-1055-1079-110-1113-1731-1751.

Gavia immer elasson

| | Late | | Location and observations |
|-------|-------|-------|---|
| April | 23, | 1:4% | |
| | | | uctoner, Mah. at ma enet, most that the first second most energy as |
| Apqil | 50, | 194. | Corvalling of tom chamaly, is a semant. It wish a war is |
| 1 5 | ept. | 4, 46 | oyst i mill. The process of the man-made liake at the north end |
| | | | of twon at ll a.m. this morning. |
| Mar 2 | 7, 19 | 47 | Cape Lookout 2 in full summer plumage. Many others noted off end of cape, |
| | | | and there were always several in sight flying low and singly northward |

ere were always several in sight flying low and singly northward (By Tom McCallister). Apr. 25, 1950, Humbolat Bay, Calif. many seen. 15-20 in entrance channel in a co. pact

unit, facingwest i to swells, about 2 p.m. I saw this one in mid-reservoir, Oct. 17,1951 with a fish, flashing wings in water, dashing ahead in short darts, with fish in bill, and even rolling over with the belly up in plain view. Watched for several minutes thru 7x35 binocs.

Oct. 29, 1952 Walker Take, Minteral Co. Nevada. 9 a.m. 4 seen from hiway. I watched one call at infrequent intervals while it was near another.

Observer Frea Ivenaen, Jr.

2661-2669-

Gavia arctica pacilica

Tate | Location and observitions

leb. 28, 1942 Respect: The diller of incivious and incivio

Jan. 25, 1947, Lint Slough, Waldport, one, probably of this species but could be red-throated in diving it does not rise out of water as do the cormorants. It just seems to slip downward as it goes forward. It spent much more time submerged than on top of the bay waters.

Apr. 18, 1954 Ft. Bragg-Westport, Mendocino Co. Calif. do ens flying north in lines low to the waves.

,

Observer Fred Evenden, Jr.


i.ea-throatec Joon 2011-2206-2254-2273-2311-2343-2402-3-2592-2665-

Gavia stellata

Late Location and observations

Legert say: 07 welt siming nor be, just of ma-o, in go, in . May 25, 1941 I matched it through the file of the second sold in the ipril 60, 1944 -- Corv in Org. which well it is a solution in the the opinal z v z z záme or. Twin in the

which the second of the same of the contract of the second this bill in the mean of the first t



Observer Fred Evenden, Jr.

Holboell': Grebe

lui. en of Tiena trip. 1 ...ic. c., ervet:
70-80-218-249-567-568-775-922-987-1079-1313-1761-1343-2403-2661-64

Colymbus grisegene holboelli

3-20-43 Washingal River, Wash, just above the dam.

Those of them & they were very acared. They seem to be there the

year around as live seen them before.

7-27-48 Klamath Marker, Rocky Pt li a couple seen

1-7-48

1 1, 1, 1, 1, 1, 1, 1, 200-10 by Farrer + Dave Marchell

Feb. 26, 1949 Yaquina Bay, Oregon (IBM) a total or about 8 came within my view. They were individuals scattered about. Found one which was injured and which we picked up. Some of these birds showed a considerable ant. of the readish plumare coming in on the neck

,

Observer Frea Evenden, Jr.

HORNED Errec Grebe lumbagor riena trica on thich obtervec:

Volices AURITUS 2591-2-2648-68-65-65-

141-223 351 = 2513-70-102-193-251-576-722-775-778-973-1004-1039-1028a(?)-1055-1079-1104-1107-1140-1334-1351-1360-1361-1362-1380-1381-1383-1384. 1386-1450-1811-1341-1904-1908-7209-2254-2744-2402-3-2421-2428-2441-2487-2499-2500-2511-2583-2584.

Late

Location and observations

Oct. 4, 1943

Vashon Island, Magnolia Beach, Tacoma Washington.

6 of them in winter plumage, hung around the area at Magnolia Beach. I sat on the prch of the cabin and watched them through the glasses for over an hour. I watched one of them dive down and come up with an unidentified fish about 5" long. This was 200 feet off shore. He would hold it crosswise in his bill and shake his head vigorously from side to side. He did this with his head in the air and also with the fish immersed in the water. Then he would lay the fish in the water and jab at it with his bill. As the fish was still alive, it would try to weakly swim away from the Grebe and so the grebe would have to dive after it again. Later on, 4 of them came in about 25 feet off shore and would dive under after sculpins in the shallow water. While I watched, I saw 4 of them catch 6 sculpins in less than 20 minutes. When they had numbed or killed the captured fish, they would work it around in their bill, so that the head would go down the throat first. One of the grebes had to cough it back up one time, because the fish had too much life as it went down the throat. When swallowing the fish, the neck was worked up and down and to the side, to facilitate the downward movement of the fish. Also one brebe chased another one that hac just come up with a sculpin. The black and white areas on the head were evenly divided with the black above.

12-mile Swamp, Corvallis // male in full plumage. Apr. 1 1947

Feb. 26, 1949

Yaquina Bay, Oregon (RBM). I am almost sure of one being a horned, an individual which allowed us to approache within about 6 fe t of him. Watched this bird swim underwater. Its mag and foot motion reminded me of a human swimming under water. Saw other small grebes scattered loosely about. Nov. 3, 1958 Pope Peach, T. Taboe, Chlif. I be sistently perched on sandy heach after flushing into water each time. Vodachrome.



Number of field trips on which observed:

Eared Grebe 89-192-258-307-328-339-422-423-424-457-10-,002 1003 - 1004 - 1028a(7) - 1028-1079 - 1107 - 1100 - 1130 - 1147 - 1285 - 1217 - 1239 - 1245 - 1279 - 1299 - 1310 - Date Location and observations: May 12 1944 Mtn. Home, Idaho, Reservoir - There were perhaps 30 or 40 (total) on the reservoir this a.m. None were present 5 days before. Test side Klama th Laws, at Focky Point: several sech - I heavily July 27, 1948. tinged chestnut on side of neck, and much larger. Hould possibly have edera a'Loodica arebe Silver Lake, Amador, Co. Calif. 7,200' elev. Either this or a horned grebe, Aug. 4, 1951 l in full breeding plumage observed for over a half hour with 7% bincos at distances from 50 to 125 yards. Light was fine, and observations were made with rests, from granite rocks above the water. Lighter colored plumes on side of head had position of that shown inpics for the eared grebe, but no black crest was observed. Further, there appeared to be a rusty color on the neck. Also, the sides were a rich and quite bright chestnut or rufous filling in a considerable portion of the sideview of the bird riding on the water. The bird

Oct. 21,1951

served to dive to escape these boats.
Mono Lake, Mono Co.Calif. 100's and 100's well-scattered over the lake area visible from the hiway. Most abundant bird.

would swim away from passing rowboats, motor boats and canoes, but was not ob-

Secret Se

•

•

• •

. .

·

Late

Location and observations

6-2-43 K. Camath Lake, O. ...

One adult with young. also several pairs doing love Lance.

May 12 1944 Mtn. Home Reservoir, Idaho. Saw 2 pair of them. One pair was evidently courting, as they were doing a lot of "neck bobbing". They gave very high, rasping squeek notes.


July 27, 1948 has traine, Klamath Lake they alout, alexans, it is a laid bear on back, toward rear, jon light any occupations 1/2 and 1, 16 to to.

Sept. 25, 1948: Grey Lodge Refuge: 2 different sets of 4 birds on one large pond. Each set had 2 adults and 2 young birds. More seen elsewhere during the p.m.

June 13, 1951 Lahontan Res, Churchill-Lyon Co. Nevada - several dozen seen, and one pair courting with neck twists, etc.

pune

Sept.28, 1957 Conway Ranch, Yolo By-pass, Yolo Co.Calif. 1 in deep water with mottled gray plumage, red eye and yellow bill.



•

will amette fiver sking to made. It would by Inc. betver, at one time, without coming up.

Corvallis on the Willamette River: Watched one for fifteen minutes Oct. 15, 1942 and timed its gives. The longest was for 21 seconds and the others ranged from 14 to 18 seconds each.

1 March 11, 1943 Willamette liver just south of albany orgon Watched one swimming in the slough just of the main river.

Hagerman Valley, Snake River, Idaho. These birds seem abde to sink Oct 17 1944 below the water without actually diving, just like a submarine.

12-mile Swamp: saw one dive just before we left. About 60 feet from the Apr. 8, 1947 road. It rear peared with only the bill and top of head showing, submerged, and swam some 150 feet just below the surface, and we could follow its wake. Then it rose with only the pate showing.

11-mile Over p - one care to lithin off or toward us as we show on the boldge senses the swim; I may solving and saimmins up a citch, yerhay to area a rest.

12-mile to me Of (originally enter a under Cost ! ! :: May 17, 1948 16.5 p.m. - hourd no oal to I down the schelch tall call. Out an an answer alm to immediately in a splittle-stage close 2001 in more rout. i uring a lo-mina of the horses by the collections. To come started out anxiotely the society of the out, but ended with a series of "peeyon's". At First trey was as look and a true of "barked" notes, but became income active a success, and notes in pater, or example the pater

being barely arcibe of reint of a letter to.

12-41- or: Sp.n. or: Sp.n. or greath arcibe of the reinter the content of the co May 15 1946 the lines it up a a court out to 10 - read of the Unit Tire-plants disher nest. au di ca passo ora, intten recat, tr. Televare 7 timbe stained egg, , very having alotted lite wing, in the contract of vity von set, but came. The egisters on the year rest, and and a soggy material similar to that on which the next in on the ct. that the node unlair rooms chine just a size or rotting couning. In page.

June 25, 1948 Cottage Grove Forerviir Pril h are my out "cost collar" too, , as a retracing them communicated the state of the s Some birthan ! Ha! is them in a civil at them. I steed one a indits tail and head and slip week pater in a "V" lithout madi at the rip ie.

July 6, 1948 1. -mile Cap: organin ature on process and secretaries of con-

Sacra. Refuge: 15 or so adults seen. 2 pairs with young less than Sept. 8, 1948. a week old. Young were good divers, too.

Grey Lodge Refuge: many seen, and they would "fold-up" to submerge. Watched Sept. 25, 1948 one steal a small Centrarchidae from another one. The fish was about $2\frac{1}{2}$ long, and $1\frac{1}{4}-1\frac{1}{2}$ deep. The grebe took it and swallowed it head first, flat side d own-up.

eb. 24, 1949 12-mile Swamp (DBM) watched one from the bridge swim with only his head out of water.

Stillwater Wildlife Mngmnt Area, Nevaoa. IBM. nest of 6 eggs floading in May 12, 1950 hardstem bulrush. Covered with soaking wet bulrush. Embryos in eggs well along. Openend egg thinking nest was abandoned.



Camp Seco Pona, 3 mi. N of Valley Springs, Call veras Co. moon. June 30, 1950. Saw about 30 young birds adattered about the pond, but not a single adult (full). The young had gray is bodies, darker on to, and pa with a blackish head, with two white strages on each side, and a little rea-brown color on the head as well as on the up or mardible.

Cap Seco Pond, Calaveras Co. Calif. 3 pairs - watched neck-hooking March 19, 1951 courtship antics of one pair at 3 p.m.

Oct. 3, 1951

Jan. 10, 1002

- Crop Screen of J. Mile office. Saluts and 3 loudly-striped young.

Dec. 26, 1954

Dec. 26, 1954

Southside Tark, Sacrayento Culif. 3hop. of the residence from an top apparently maiting. I persistently chosing of en, trabling from an top and behind of hear, banding on, both sometics submers no, and rear one with aplearance of mounting of or in deep water it surface.

James and the Attack. The many well of the throad there was small known to be a 2 gotton.

| | · | | |
|---|---|--|--|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| , | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

Diemotea nie ribas

oct. 16, 1955 To Farallons: 1 at 3:15 p.m on return. 5 others ameured 1 by 1 out of thin air between Sl and North group. Lany other sea birds shout them.

Fulmanus dacielis

October 16, 1955 To Farallons: 1st one sections over half way out and was worm don't form.

Others seen during day were not so don't, and one or two very list. Saw as many as a individuals in half been resting on rates. Then build, and short light bill with tube seared them. They like a correctors, but with more wing-bouts, and with parret-like head.

| | · | | |
|---|---|--|--|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| • | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

PACE-FOOTED SHEARWATER.

10/2/56 2 OFF PT. PINOS, MONTEREY CO. CACIE.

| | | • |
|--|--|---|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Puffinus creatopus

- Notaber 5, 1949, Neptune State Park, coast of Lane Co. Oregon. About 1 p.m. I saw about 6 of them paralleling swell crests about 1-mile offshore, in company with sooty shearwaters and Pacific kittiwakes.
- May 9-12, 1954 Off Pt. Pinos, Monterey Co. Calif. Dozens in company w/ 100's Sooty's.
- October 16:1055 To Farallons: Drite below, Cark back and Wingtops, largest form bot. Fout your islands, and seem only in a.m. to Lean + 1 just 1 of a free b.

| | | • |
|---|--|---|
| | | |
| • | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Fuf imus bulleri

October 16, 1955 To Farallons: Seen over half way out. This below, large, and with light back set off by black can and very dark teil. The most account of mina-length visible at consider blo distance with named eye. Zibby flight. 2 or 3% seen between 18 and 11 groups.

| · | | |
|---|--|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Puffinus griseus (Gmelin)

- Sept. 6, 1946, Seaside, Lewis & Clark Turn-around, 7 p.m. 10.0's and 1 00's of them flying north, just beyond the breakers. They were going for at least ly hours when I left. Huge "rafts" of them could be seen on the surface, too. Feeding, I suppose.
- Sept. 7, 1946, same place, 4 p;m., 1000's of them flying south just beyond the breaker line. The other evening they were heading north. A few rafts of floating birds could be seen as well.
- October 5, 1949 Nepture State Park, Lane Co. Oregon $\frac{1}{4}$ -mile offshore, paralleling swell crests. About 25 were seen in company with pink-fo tea shearwaters and Pacific kittiwakes.
- July 21, 1951 Off Pacific Grove Pt., Monterey Co. Calif. Noon. 1,000's passing northeast offshore. Their density varied, but two different counts showed 100 and 125 passing a given point per minute. The time we watched was about 20 minutes.
- July 22, 1951 Monterey Peninsula & Monterey Bay, Calif. 1,000's of them passing northeast offshore, and 1,000's circling in concentration of gulls, cormorants, and shearwaters off Marine station, many on water.
- May 9-12,1954 Off Pt. Pinos, Monterey Co. Calif. 1100-1500 offshore. On water in early morning, and busy soaring in afternoons.



Juff nus tenuirostris

October 16, 1935, To Farallons: A very few sean near the Soction. Sliver and shorter build and with durk underwings. May have bid of at 1 ast one of them.
About 12 seen between N and SE gr u s.

| | · | |
|---|---|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| , | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

MANX Puffinus opisthomelas

·lack-vented Enearwater

| | | | • |
|---|--|--|---|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| , | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

(comodrona furcita

3 457 - 3545-

1. o. c. o. a. . com

3545

Oceanodroma castro Harcourt's Petrel

Off Ocean City Md. May 23, 1970.

l crossed wake of boat, steady flight, back clear in going away view. Slight notch as in Robbins book, with complete white band as on Wilson's and not broken by dark, as on Leach's. Was seen 32-4 iles out.



Beal's Petrel

Oceanogroma leucorboa beali

Late

Location and observations

16, 2, 40


Pittock dira (inctual : Tour it hend on to India who we will be a fit had evidently been been in by or of the interpretary. It is a first one found in Fortland. It is still and it is in the windle to Glabert was sit, we seek all that it.



Beal Petrel Found in Portland.—While touring the grounds of the Pittock Bird Sanctuary in Portland, Oregon, on May 16, 1940, Harold S. Gilbert, Willard A. Eliot, and I found the dried up body of a Beal petrel (Oceanodroma leucorhoa beali) lying on the lower branches of a fir tree, not far from the Sanctuary's pond.

The assumption is that it had been blown inland by a severe winter storm, because Portland is certainly a long way from its normal habitat. The specimen was turned over to Mr. Stanley G. Jewett.

Jewett and Gabrielson (Birds of Oregon, 1940, p. 89) mention one specimen collected near Portland in 1923.—Fred G. Evenden, Jr., Oregon State College, Corvallis, Oregon.



•

•

(ceanoimora horochrea

| | · | |
|---|---|---|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| • | | |
| | | |
| | | · |
| | | |
| | | , |
| | | |

'Blac' etrel

Inomelonia relatia

.

Observer Fred Lvenden, Jr. humber of field tried on hich day to wa: late Location and observations Klamath Lake, Une - 8 30 p.m. Saw many white pelicane flying around the lake the overrook the train types of we see we make some for the set doing over 50 at the time Dec. 3, 1943 Wallula, Wash., along Columbia River. I saw two of them flying downstream. August 14, 1946 Fern Fiage Lam, Lane Co. Bill Graf saw White Felicans on this date ! Fern Fiage Lam, Lane Co there were between 25 and 10 of them sucttered June 25, 1947 over the south end of the recervoir. At a distance, one bind appeared to be on a nest (not guaranteed). Fern Fiage lam, Or I sat them thouting on the water, or perched in a ratio of July 24, 1947 island, and one group of 7 took to might at the same time. Form lidge land about (1) from, additioned well over the filence. August 8, 1947 lein lingo: which belon this bound to have the possible to be den-Cest. 20, 1:47 Then (cycles wheren) whilst 15^{11} km . Let 11^{11} 10^{11} 10^{11} 10^{11} 10^{11} the liter, proticing wiles. Otherwise very time. To be a topled observed about 50 binds carbain som evan about somet a gera de . Pern Bioge: " a tot I on the ut a community of a least to be a least ; but I one were 5 in the way of high a country as he have " on " June 11, 1948 orveloped on ill. I frame or only pinch as the chains a country courant time cause of anoth. Sept. 8, 1948: Sacramento Refuge: at least 200 seen on the area. Most in small, scattered groups, but one compact flock of 60-75 birds, ½ adults. The immatures were more alert than the adults, and had more pinkish bills, and fuzz-downy heads. No horns seen. Sept. 29, 1948 From 10 mi. SN of Merced to Merced, Merced Co: 150& seen. We saw about 40 of them perched on the ground with Amer. Egrets, but these took off and joined a larger flight that passed over. april 4, 1949 - Lacto. Lignal Depot - 830 a.m. a flight of 42 going N, vatter low - 2-300 feet up. also 3 seen et 9:30 over Yolo By-Proo, Yolo Co., leading N. Gray Loage Refuge, ST Butte Co. Lat least 50 seen. Up to 30 in the air at one May 5, 1949 time. All were in full adult plumage, and about 25 passessed the breeding "horn" on the upper bill. John Cowan, Mgr., was surprised when I suggested they might be nesting at the refuge. They were present and last summer, but no though they were immatures. He will keep his eye open this year. June 3, 1949 Sprague River, along U.S.395, Klamath Co. 550 p.m. 5 cheating in river and perchea on rocks in the stream. Also 5 seen in river near Chianquin Jct. July 25, 1949 Naga R, and Can Pable any marches 13 0-250 seem it in s. no to a e Dec. 6, 1949 Salinas R., hear mout h - 245 p.m. Observed 4 white pelicans on the river, and about 20 of ///e//just 100 yards away. Dis is te ; 1st time I've seen them at /the brown pelicans the same place at the same time. anaho Islana, Pyramia L., Nevaua IEI. at all stares of reproduction. 6 pous found, and a 7th having been so far along that he birds were greatent. One pod was seen on top of the island liter we left the island and here out in the

found, and a 7th having been so far along that he birds were present. One pod was seen on top of the island efter we left the island and were out in the boat. Giles estimated 2500 youngwere present. This does not include the pod at the top of the island. Fish coughed up by the young birds were carp, Lahontan chubs and cui-ui. Possibly 75% were carp. Trelease counted 25% carp average two inches long from one joung pealean. One grown bullinged was also found.

eggs Ling of the state of

•

Page 2.

July 21, 1951 Pt. Lobos State Park, Monterey Co. Calif. 2 white young with grayish head and black primaries. On Bird Rock.

July 22, 1951 Fonterey Peninsula, near Bird Rock? 5 flew north overhead.

May 1, 1952, Sutter Refuge, Calif. 25 seen, some with breeding "horns".

Feb. 3, 1053 Fonte Fat, S. Inis Shishe Father, Settle of the Father, and 60 in other at sandy edge on west side of bay.

May 30, 1953 Fone Beach, S. Corner Take Tahon, El Dorne Co. Calif. of sper.

June 27, 1954 Echo Take, Aldorado Co. Calif. 7 over Flend of Taho.

June 27, 1954 Tore Beach, Take Toloc, Golff. 3 seen.

April 23, 1958 25 of some yello Bypaso Safaborado Co. Calif. 2 circling high over Sierras and gliding west, to north of U.S. 50.

Yelder Manual Reference Company Remains and Sierras and gliding west, to north of U.S. 50.

.

•

Eastern 1854-2239-

Observer Fred Evenden, Jr.

Pelecanus occidentalis californicus

Late

Location and observations

Oct. 30, 1948

Seal Rocks, Cliff House, San Fran.: keveral 100 on the rocks. Many immatures. Most crouched down out of the strong wind.

Oct. 15, 1950.

S.Frisco Bay to Farallon Islands, Calif. First seen just north of lightship on trip out (at 122041125 and 3404512 N. 4 were flying westerly. A few others seen a few miles Moof Farallons. On the islands, several hundred were perched high, particularly of the cone-like cost island of the group. Few were flying aroun. On the return trip a few cone to the boat in gulls die, grabbing fish cleanings tosled overboald by fish from . A few landed on water with gulls and moved along with us. Parkness cane before we reached the Golden Gate. However, silhouetted against the city lights I saw brown pelicans heading toward S.F. bay from the ocean. A total of 55- or 40 were seen between 1830 and 1800 PCT. Also saw 5 in one group heading out to sea at this same time.

July 21, 1951 Monterey Peninsula very common, including many young on bird rock in Pt. Lobos park.

FebF 3, 1953 Morro Bay, Calif. 8 or 10 seen. Saw one make 4 unsaccessful dives from 3-60 feet heights.

May 11, 1954 Pt. Pinos, Monterey Co. Calif. 1 pair seen w/ Sandy Sprunt

Oct. 16, 1955 To Farllens: few near Golden G to in a.m. Tany on return in .m. ome diving for Tish. .ves Fellowed boot at thes.

Oct 4+5, 20 Marin Co. 101, 5/100 - 75 - o feeling or the deline of the second of the s

Aug. 28, 1962, Reat Island, Sonoma Co. Chlif. In the first solf there, and liter as fog rolled in, solder a nevite on the tot be same island.s

Observer Fred Evenden, Jr. Lamas, Wash. - 230 p.m. it were and of men as the me on a without the off like Cools, from the water. Dec. 3, 1943: Wallula Wash - along Columbia River. San Two at close range. I was surprised to see them this far up the river. Kings Hill to Glenns Ferry Idaho. around noontime. Saw one that was flying Oct 10 1944 downstream, but not over the water. It was brownish-bluck and had a yellowish throat pouch. Havasu L. - any - Calif: Seathered in small groupes ourstile take during the day, & hunched Mar. 19, 1947 in certain trees at nite. Many "sun-going" at moonline. Fern Liage Tam Area at least a half cozen were seen perched on dead June 25, 1947 snags over water. Fern liege Lam or several perched in what there out in the Law, and I July 254, 1947 watched one drop into a small lead below the dam to reco.

Bayshore Drive, Ber keley-Oakland preening and sun-bathing with wings Nov. 13, '48 held out and drooped on a rock within 5 feet of the hiway at 10 a.m. I stopped 5' from it and watched a long while; Iris gray-green. Others swimming Franklin Fanch slough E of Courlana, Sacto Co. one had to beat much of March 12, 1949 its wings into the water to take off. June 40, 1949 Rye Patch Res., Barsning Co. Wevada: 25 flew over orwnstream inl grap. October 5, 1949 Yachats Park, Lincoln Co. Oregon. two observed about 20' irom water on sand flats exposed by low tides. With gulls. They waddled around. Dec. 30, 1949, Hogan Res. - Calaveras Jo (2) 25 in area - 1 disported a 41" Perca flavescens as it flew over us. The purched of the snegs in the reservoir. June 24, 1950 17-mile Prive, Monterey Co. Calif. In the evening at 7 p.m. we arrived at Bira Rock along the Trive in the e to count about 4,000 of them on one side of one of the 3 rocks in the group, and all the lake here groups of 2 or 3 to 15 or 20 were suriving area both the north and the couth, although most of them were coming from the north. To estimated that 10,000 or more used the rocks that evening. The following marning we saw flights helding north all ng the east eage of monterey Bay, with directive pur oses. Last evening the vere flying west along the south and//side of the bay out to the ocean, and today flying cast and north along the south and east side of the bay to the day's feeding grounds. The bird rocks must draw cormorants iron a great distance - at least 15 miles. following the bay coa trine. S.F. Bay to Barallon Islands & return: Thuge numbers free ng over Potato-Oct. 15, 1950 patch Shoals (4-fathom bank) to the West of Bonita Paint and Channel. They would flush ahead of our boat. In few sec all the way out and back. At the island, 1,000's were on near-vertical slopes as well as on that areas. Al so 100's feeding in waters around island. After & p.m. low-ilying strings of 35-125 came in from the south and east. Each bird flew behind and a little to the side of the one shead. One time & strings from different directions crossed each o ther and still maintained formation. Strings broke up as they swept upward to land on the islands. Some of the Baird or Franct's

July 22, 1951 Bird Rock, 17-mile Drive, Monterey Penin. (1,000's of them passing from NE to 5 or 5.) to this rock, in strings of various sizes. Observed threat of sea lions to nests of mud built on these rocks.

May 14,1952 Advanged Z., 5 of Alamo, Lincoln Co. Newsdar _ 35 w/ nests in live cash (1) over water.

Dec. 5, 1950 Playa Lel Rey , LA Co. Calif. 6 perchea on hi power lines overBallona

Creek Canyon this a.m.


were also in these strings. Flights were still arriving when we left at 330 p

| | +* *. | | | | | |
|---|-------|---|--|---|---|--|
| | | | | • | | |
| | | | | | | |
| | - * | • | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| • | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | • | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |

÷

Olivaceous Cormorant
Phalacrocorax olivaceous
mexicanus

Feb. 22, 1951, Laguna Madre, Port Isabel, Texas - several seen.



Observer fred venden, er.

776-178-1004-1079-1107-1110-1113-1567-1631-1689-1761-1901-1954-1988-1956-

2060-2-01-2-17-226/-1311-2344.2317-9-9-2511-2669-

Fraiderocord pericillatus

Late | Location and object tions

Feb. 3, 1953 Morro Rock, Morro Bay, Calif. roosts on ledges and wind "blow-holes" high on west face of Morro Rock. Seen only on ochean side of rock and cut on ocean.

May 11, 1954 Pt. Lobos, Monterey Co. Calif. On nests on Pt. Lobos. Others carrying seaweed to nest sites.

.

.

Observer Fred Lvenden, Jr.

Phalacrocorex belasious res dincin,

rate

Location and observation .

arch 30, 1941

Agate beach mighthense? In a rocky iskura that the board on tirus with home cold with the rest of mineral section of the cold the body. The need has to have the mineral sections and the cold the section of the cold the col when picking or profiler them were. on small projection or to distribute the. The in for a Lancing of an introduct, say A of the Language and in the sening come to ano. it from their as a compare to be only one of the bernet symmetry on the energy of the light of officers for the day of

May 11, 140

Checkin to Loud 1 or a level neve to the both below to the Light.

On coast, S of Malpaso Cr., Monterey Co: 3 nests on Jeage 20 to grow head of quiet bay. Wests & feet a art. Mests built u with lant material, and adults on nest. Nuch excrete. Chiri swallow rests within a reet or them.

Pt. Lobos Park, Monterey Co. Culii. Young 2 to 2/3 grown in nests. July 21, 1951

Things.

4/19/67 and to any in sheer free, attracted a 6. 2 accided with the besting and single to any in sheer free, attracted a 6. 2 accided with 31, but film way a single to any in sheer free, attracted a 6. 2 accided with 31, but film way

.

•

Bluet12+77 - 1841-1854-

osciver into venct, o...

Tregunza's heron

1309-1405-156/a-1566a-1581a-15816-1612-1612-1615-1637-1646-1743-1751-1838-1839-1840-1841-1948-2000-2002-2005-2019-

Location and observ tions _. Uč

Colo. R. Indian Res. south of Parker Ariz: 2 with nests in tall cottonwoods 3-20-47 Used the same nest sites last year.

٠., , •

.

April 27, 1950 9:30am to 12:30pm With LeRoy Giles Heron Rockery and Vicinity at near Tule Lake

Pied-billed Grebe - heard one several times
White Pelican - Noticed about 200 circling over Mutgrass Area
Double-creasted Cormorant - Noticed a couple flying about over the
rookery. Possibly some were nesting in it.

Blue Heron - 24 nests were investigated. All were in hardaten bulrush and were constructed of this species and Suaeda sp. Young were not were hardaten as the night herons. Many were only several days old. Data on the 24 hests are as follows:

2 eggs, two nests
3 eggs and young, 1 nest
4 eggs and 2 young, 1 nest
4 eggs and 1 young, 1 nest
1 egg and 4 young, 1 nest
4 eggs, 9 nests

4 young, 1 nest 5 young, 2 nests 5 eggs, 6 nests.

American Egret - at least 3 were resent around the rockery, but we didn't run onto any of their nests.

Snowy Egret - 10 nests found. Data as follows:

l egg, 2 nests 2 eggs, 1 nest 3 eggs, 4 nests 4 eggs, 3 nests.

It is apparent that the egrets have were last of the herons to start nesting. In several of the blue herons nests we found small eggs which might have been out there by this species.

Black-crowned Night Heron - Found 50 nests. Some of the young had already left the nests and were flapping about. Others could already fly. Data on nests as follows:

l egg, 2 nests
2 young, 2 nests
2 eggs and 2 young, 1 nest
2 eggs, 1 nest
5 young, 7 nests
4 eggs, 5 nests
5 young, 5 nests
5 eggs and 2 young, 1 nest
5 eggs, 7 nests
4 young, 5 nests
6 eggs, 1 nest
6 young, 1 nest
1 egg and 1 young, 1 nest
4 young and 1 egg, 1 nest.

hite-faced Clossy Ibis - .e noticed reveral hanging around the rockery and wondered if They the ibis won't nest here too.

Mallard - pair
Mallard - pair
Mintenen Teal - saw 8 pairs, 2 pairs were courting together and then seperated.

Pintail - pair
Madwall - pair
Madwall - pair
Shoveller - pair
Turkey Vulture
Marsh Hawks
Milldeer
Milson Phalarope - saw 20 circling on Morthwest Pond as we drove out.

Tree Swallows - over northwest Pond
Darn bwallows - over Morth est pond. Thist seen this spring on area.

Marsh Trens

Savannah S arrows

Song Sparrows

More Data on the rookery

The rockery was sighted from the apirpline when we went up on april 25 to sensus the area. It is ap reminetely 1500 feet long and 500 feet wide, which is the extent of the hardsten bullash stand which it is entirely in. e walked a strip through the middle of it and tock data on the nexts visable. The estimate that 500 heats are present, about 250 of dich are night heron, 150 blue herons and 100 snowy egrats. The aclony at mark to be a new one considering the size of the nexts and the fact that all were escuring or showed evidence of being essential this year. The acres care and night becomes nexts were consider this year. The acres cultush, while the blue herons nexts also centain sea-white (Suaeda se). The estimates above are preliminary, subject to change.

Lie.

Shaker the state of the state o

SURVEY SWITTERS TO SOLD OX

April 29, 1950 10:00 moon to 4:00pm
Data below is on 866 mests found in the heron equally at the north end of the numeronarea through a search of the area of the order of the area through a search of the area of the order.

The colony is about 1500 feet lany and has a miximum that her most than 500 feet. It is situated in a stancof bologue action which is ever rounded by a zi m of Typha. Tome small pools and a chemical are present in the doir us. Host of the new two were demonstrated are presented in the doir us.

ond a white-freel phase, i he seeked and below the probably of early in the identified tion of the rests. The policy of a part begun to come into the velley for the summer and the bulk of the birds have not yet arrived, there is a possibility of more ibis nesting in the eclony. The colony appears to be a new one. To old nests were found and all except the blue herens nests soutsined achaids rable material so there is a possibility of the second and all except the blue herens nests soutsined achaids rable material so there is a possibility of there being used last year.

Black-orowned Wint Heron

Night heren neets were constructed entirely of beiroes sentes. Young were already off some of the nests, as flapper night herens were everywhere present as well as some young which could fig. this so-counts for the large number of night heren mests which were about ened. Night heren were halfway in sine between the blue heren and snowy egret.

Total Nests found 422 Total occupied nests 545 Total abandoned nests 76

The following combinations of each and young were found along with the number of grant nests with each.

| Smpty Neats 78 i eas suc i young legg 15 i eas and 5 wo mg 2 ease 13 f ease and 1 young | |
|---|--|
| dogge 96 1 goung 1 ogge and 1 young 5 ogge and 1 young 1 ogge and 1 young | |

In addition to the above I enowy egret ogg was found in each of 2 night heron nests.

and the state of t

en en la comparte de la comparte del la comparte del la comparte de la comparte del la comparte de la comparte del la comparte d

there is a property of the state of the stat

.

I is now the company of the second of the second property of the second second second is the second of the second second

STIDIWATER WINDLIF! MARKO MARKE ARGA, CHURCHILL GO., NOV.

SURVEY OF HERON COLORY (Continued)

Aoril 29, 1950

Black-crowned Bight Heron (Continued)

Clutch figured on basis of eggs and/or young combined:

| 1 egg or young 19 neats = 19 total 2 | eage end your sam |
|--------------------------------------|-------------------|
|--------------------------------------|-------------------|

Average per clutch on basis of these figures 3.614.

It should be pointed out that these figures are not entirely valid for basing avage and out that these figures are not entirely plotted laying some acre young hid following and acres young hid following a figure of the nest.

Clutch figured on basis of nests which centained all young or some young. This eliminates the MANASSINIII to the figuring in the nests which did not have him clutches completed.

| 2 | eggies young 4 poung or 2 egg | | The state of the s | |
|-------|-------------------------------|-----|--|-----------|
| 34567 | and 1 young 16 | 284 | | and young |

Average per clutch on bosis of these figures 5.591

FILORE THE STATE

heron nests, but averaged smaller in diameter. Eggs were somewhat smaller than those of the night heron. Snowy egret nesting ass not as for along as the night herons or blue herons. The congruence young snowy egret was found hatched. The nests from the congruence best was found hatched. The nests from the congruence.

Total nests found 63

Potal nests with eggs 61

Total nests under construction 2

The following combinations of eggs and young were found along with the number of nests with each.

* 1

Jan gladin

Charles and golden distance with the in the best

ang control of the c

LIBER SERVE TORREST THERE

is the ran interpolation in the second of th

f(x) is a sum of the property of x and y and y

ស្ត្រី មកទេ ១១ ខេត្ត និងស្ថិត ស្រឹក្សាសាស្ត្រី ខេត្ត ប្រធានាធិបតី ប្ធ

SURVEY OF HERON GOLDHY (Continued)

April 59, 2959

Snowy Erret continued

A TOTAL AND

eggs. This figure not entirely valid since some birds had not completed laying.

| 1 | egg | 10 nes | li del | | | |
|----------------|---------|--------|--|-------|-------------|-------|
| \$100 \$100 | 4 4 5 | | E C | | | |
| | No. 11. | | TE | | | |
| 10 | GERE | 22 | 88 | | | |
| | 0379 | | al. C | 7 | | |
| | | | and the state of t | Cotol | 的是一贯 | and m |

Average per clutch on bais of these figures Z. 197.

Olda Herra

eldereble recommende were large size (alay and sees section as some eldereble recommendate in the elder of the elder seeds. In a large seed the first seed of the elder seeds in the first seed of the elder seeds in the first seed of the elder seeds are the first seed of the elder seeds are the first seed of the elder seeds are the elder seed of the elder seeds of the elder

Total nests found - 73 Total ecoupied - 71 Total Empty 2

The following combinations of eggs and young were found. Sumber of name with each econinction also shown.

| 1 | erty les te er | * | | The state of the s |
|------|-------------------|---|---------------------------|--|
| | | v. v | B GAR and B Alle | 100 mg |
| | 0: | des | Le opposite and 21 proper | "Lord |
| 60 | 2000 | 6' 3' % | | · 4 |
| 3 | e: c | 5.0 | | 100 2 200 |
| 23 | 95,50 | and the second | | 1, |
| | OFFER | The same of | | The said |
| 344 | of the state of | 3 | | in the |
| 4(a) | VIMILA | 4 to the state of | | |
| | 713 23 | T | | |

or the country 6 . . . 5、13衛 泰 myse d The second of th op indi ₹. 2 mg . . . កក្រុមស្រែក ខេត្ត សំភាពនៅ ស (១០ 🚧 📆 🤇 east to the common cases the bold and speak towns of a 10020018

A STATE OF THE STA

> 11 4.00 634 /··· 8 5 1 to the

1. 1. 1

. . .

and the second

SURVEY (& HARRY COMMY (Jontinued)

a 211 19, 1950

Mino Horon Josti wad

Clutch figured on basis of on a sad/or your co bined:

| 1 | egg or | TO MAR | 1 | neals | | |
|---|--------|--------|--------------|-------|------------------|--|
| 2 | | - Ma | 1 27 | | 6 | |
| 3 | | | | | 2503 | |
| | | | 20 | | 100 | |
| 5 | | | × 2 | | 140 | |
| 6 | | | 43 | | ,4 | |
| | | | A CONTRACTOR | | HET PROGRAMMENTS | |

Avorage per cluber on basis of these timess 4.138

Clutch figured on basis of of costs which contained all jours or some

Average per clutch on besite of these filters 4.175

Aperiosu Areu

but contained little or no deport vegetation. Aggs second identical to those of the blue heren. They are respectfully. The lea number of american egret negles here is not surprickly, as trip bird is welder second on the area.

Shite-Thood Alpsey this

Biss Scoonsilessin

Some nests were missed in the count, as night heren and showy egret nests were especially well hidson. Leveral corners of the rock-cry were not reached. Thus the following estimates.

| | TO THE STATE OF TH | A processor and the contraction of the contraction | DOSGI GEVINO DE |
|-------------|--|---|-----------------|
| Might Heron | 423 | 70% | 600 60 |
| Blue Horon | 73 | 200 | |
| Ibls | | #3 | 67 |

| | · | | ני |
|---|---|---|----|
| | | | |
| | | | • |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| , | | • | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

TI IN I I MANUEL RANGEMENT TO A CHURNIEL OF

EURYSY OF HERES COLLUNY (Continued)

april 19, 1950

1. del Come by the continued

heren was found excume several of the newles one flap or night heren was found dead. It had a 7 inch carp protracing from its oill. Acres e of the rockety is a cut if and a last 6 light has a newle were in hologous actus. The cair ware in cattail on the border of the stand of beingus actus.

The colony is isolated from the mainland. It is recond via an intint adjoining island.

ě The second secon · ·

V. California eron

\[\begin{align*} & \frac{1234 - 1033 - 2034 - 2034 - 2134 - 2234 - 2234 - 2234 - 2234 - 2234 - 2234 - 223 - 236 - 237 - 244 - 189 - 189 - 191 - 192 - 193 - 199 - 209 - 21 - 212 - 218 - 214 - 223 - 236 - 237 - 244 - 249 - 312 - 314 - 351 - 364 - 365 - 364 - 365 - 364 - 365 - 364 - 365 - 364 - 365 Horse shoe lake H. F. il, i war a small tree + hanging from it till it. Camas Wash, Upper Facamas Lake 2 pm. Came in case to im with a majort. He was a visually a hing of the last years reg. when he had I had I had yours not no willacceloped, cleap ruly sed rown may in a the appeal agreed the rody color. He was at most right, this He was ited his Largewings, water + yet "by - blen with ling in a ray, we " bookerstored". ... It I'm is to every lie way I way - stended Dege airead of im as is and intella making Camas, Washington 3 - 7 p.m. Saw several making their way to the Columbia river from LaCamas lake. They went out over at different times, but May 20, 1944 they all seem to follow the same line of flight, coming and going. Leadbetter's Peninsula, near confluence of Willamette and Columbia Rivers. April 5, 1947: 10 a.m. I saw my first real heronry, and it was definitely worth the trip to see it, too! There were 150-200 nests in Cottonwoods growing along the north side of a small slough. Although there were acres and acres of Cottonwoods in the area, all the trees containing nests seemed to be in an area of less than an acre. Nests were from 50 feet height on up, with some up around 100' I would guess. Nesting trees didn't seem to be more than 100 feet from the slough. Some birds were still building nests, and carrying on with courship aisplays such as "cresting", and bowing, etc., from lofty perches. Other nests apparently are completed for there were a goodly number of birds on the nests, and probably incubating eggs at this time. Shot-like sounds produced by hitting a stick against a log resulted in the flushing of even those birds on the nests. In-mile Tuamp: Swerm seen, and one had over mill of the Thight Postrers Aug. 22, 1947 in the wings missing, making it woll a look 1 arg. They were all much note willing to give crothe of verious types then I flushed them. May 9, 1948 Leobetter Penin, Porthago: Pront come time location at the rockery. From the souma, we would judge some young tirds have notched. Eaults conjunct mosting material to some extent. The moonery is a slarge or larger than our past estimate of 175 to 100 nests. It a pears or the sens actional nests have been made at the west and of the classy, but take is not cartain. Sept. 28, 1948: Marposa Dam Site, Mariposa Co: feeding on young bullfrogs in pool still remaining in creek bed. Mar. 15, 1949 Leadbetter Penin, near confluence of Gillamette and Columbia Fivers. A few new nests have been built on the otherside of the slough. About 150 herons were present. Only one to a nest, the only exceltion being in one case shere two biras were perched at one nest. The Lever a bira came near a neighbor's nest or flew close by he was promptly challenged b, a nead throw acompanied by gutteral notes. With owny one bird to a nest it would seem that they were makes establishing nesting sites. . . . and then we say one carrying nesting materia. The new nests on the opposite side of the slough were not occupied while we were present. The surpled and watched the rookery for about 50 minutes around noon. The cottonwords were not get realed out. Only buds. July 23, 1952 Huckleberry, McCloud, Siskiyou Co. Calif. 1 p.m. along creek in yellowpine woods. Mar. 17, 1962. Canyon hanch, clime lanceh, win 6 Cliff. The 1907 for the top of redwoods on northwest slope of ridge just above reach bldgs. Why on nests, and one erg visible with cird standing beside it on nest.



Black legs - yellow bill.

- Fern Riage I am a total of a sere seen this and a sittement beaution August 8, 1947 the NE side and watched them fly to the stath em, and take or reversal miles, where we later somether gain.
- 12-mile : wamp fat low to been else non there in the analytic both August 9, 1947 ends. Three were seen at one time; gids ere token of a torother; one has observed with his mallet pritty well filled with the oler or allinogs from 2 to 4 inches lone, which I sutched it date, and swellow. It was funny to watch the bulbe clip own until it reserve the "to a the column" or predecestors. him satching two, a third came over, are made none bluff gives at the other two. The circu die "task" to the mestive, and to each other while I sutened. They are a bountimenty racelus site to watch as they Ily leisurery and rightly,
- Sept. 28, 1948: Mariposa Dam Site, Mariposa Co: feeding young bullfrogs, too. Oct. 2, 1948 Suisun State Refuge, Solano Co: these or snowys had nests from 30-75 feet up in Eucalyptus groves.
 - Jan. 8, 1949 Gray Lodge Refuge: several 100 in area. On way up I noted about 250 of these egrets perched on the bare earth of the west and north banks of the Sacramento River, whichever was exposed to the early morning sun. Blue Herons, too.
 - LaJolla, San Diego Co 3 seen perched 2-mile out to sea on kelp beas. Jan. 27, 1949 One was observed in trouble, with its legs tangled in the kelp.
 - July 12, 1949 Gray Loage Estinge, Butte C. P[Calif: seemen to be figure in an extractions out over the country-side from an enea on Butte Creek at 10 st 6 miles
 - July 24, 1949 Yosemite Valley, Mari coa Co. Unif. & p.m. circling over valley between al Capitan and Cathear a books, and it find by healed north over rid e just east of la Capitan.
- Nov. 16, 1949 8 miles N of Butte City, Calif. about 200 seen. 150 were seen in another field. These fields were recently cultivated - usually had been in rice. There were no egrets in the Gridley Egretry
 - Mar. 28, 1950 Bira Rock to LaJolla, San Diego Calif. 545 p.m. one seen 1 mile offshore on kelp beds.

March 24, 1940 - 2 mi Not deapprose, Oregon - 4 eggs in large nest of sticks & green branches. Nest ina copy. 75' up in a D. Fir. Cuttoos also 4 at 80' D. Fir. March 28,1942 -2 min NE of Scappoore, Oregon - 4 eggs in nest of large green fix branches. 70' you .: Bitis. June 13, 1954 Sacramento Weir, Yolo Co. Calif. Conspicuous by their absence -none seen. - Giff & Bis in Page 15 is it is not the state of Dec. 40, 1951 NIJA even, Marie



Number of field trip on which seen:

A flight of 1:-14, and they seemed much smaller than the blue heron,

but I could see no colors because of the time of the evening. All I saw was their silohuette against the fading light in the west. I did

notice that their, necks were crooked.

Aug. 16, 1949 S Butte County: Thurs are to 1.5 miles best of scheener Countries Road. Lat. 3 00 x : 11 N, and 1: 15-41 . Congity . (: E.T. 7 , . . 6 . -4 . 0 066)

There were objet :50 if the contract of the policy of the contract of of Car row to real throught, a mostly mentale and the respective


road, - ile and ! It be a line this laternoon, then the on the ut + 5-5 :

of them present - noisy takes a margian as it is a

.

Reddish Egret Dichromanassa rufescens

Feb. 22, 1951, Near Port Isabel, Texas - one perched on post in open water - dark phase



Louisiana Heron

Hydranassa tricolor ruficollis

Feb. 22, 1951 Laguna Maore, Port Isabel Texas wading in bay, near blue heron.

and the second of the second o •

2237-2243-

· Florida caerulea

Tittle Blue Heron

| | · | | | |
|---|---|--|--|--|
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| , | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

Lute Location and os av lion;

Fuccing river: The process in the knowledge, just the contract of the form of the contract of ugust 7. 154.

Procins liver: Of to area up to areas more in a transport or in a

The finals of adima form a scenar up the line objects in a district way of the very near the converse of the second than the converse of the second than the converse of the second than the s pril 26, 1941

1: -mile ruan; worth in porvening form mosph mealer of it. . It is Way 24, 1941 quittly to an abstract of the solution of the

more and living the time? Ju. 26. 194

Corvallis, Gravel Pits. In the muddy bottom lands I flushed a small March 19, 1946 (Little Green) Heron (or bittern). Body about 12" long. Wingspread 15" or so. Black and wings blue-grayish in color. Not too dark. Flew about 50 feet and Aanded again.

Stewart L., Corvallis O'A air were present, so I suspect they have a July 1, 1946 nest in the area.

Salem, Willamotte Slough Due perchee on log ruft i the railroad. 'uly 13, 1946 Stepart L, Corvallis, "I found the nest of the above pair this p.r., uly 181; 1946 but it had already been used. It was on the east side of the lake in a "lagoon" surrounced by veget time. It was at 15" over the water in the fork of a willow tree. The nest not ever healt" anameter (the main part) for a few misc. twigo more longer. Naturian was piled 4-5" ase, and formed a sauctr-cup thich was linea with finer twigs - all haid in the sime direction, parallel to out of in. The larger twigs agreemed woven.

"Poison Oan" Lake, Linn Co. Ca co to a pair that built a nost l' vover July 18, 1946 Later on the northwest side of an esh the on the firside of the lake. Make out of large tries and smaller ones inside it. I tion't climb to the nest, for the tree is to such a fewly and no journ in it.

July 20, 1946 - Poison Our Taxe of I climbed to the nest which I found on the little. but the kimbs weren't strong enough to permit my bettellay reching the nest. I could see in it, no that is apart all. The makene large downy to no with a Will over $\mathbf{l}_{\mathbb{S}^n}$ range. It has see like there might have been another downy young, but it could have been the bend of the sing of the other that I sav moving. There are also definitely at least : pole blue error in the most. The name less the most believe I appread ed within 25' this time (On the 18th, the took out the rest after I was under it). The agult stayed quite close all the while I was climbing to the nest.

to approach quite closely, so that detailed observations could be made even without binoculars. On Silver Creek there are always one or two to be found just above the forest guard station at Silver Lake as well as on Silver Lake Marsh. This heron should be looked for along the Chewaucan River and on the marsh near Paisley,

Oregon, and in suitable habitats near Lakeview, Oregon.

The Anthony green heron is usually very shy. Its liking for dense cover, where it is available, and its habit of "freezing" and permitting a person to walk by make it a very difficult bird to flush and to observe. It is definitely not a bird that can be found and observed by the casual methods of the "bird-walk" type. The observer will have to brave the dense alder and willow jungles along the streams in the western part of the state. Except on Warner Lake, the birds have always shown a definite preference for these jungle-like thickets along the streams and marshes where the vegetation crowds down to the very water's edge. Here in the shade and among the foliage its colors blend very well and its retiring habits make it most difficult to detect. Usually it will be seen only when it drops into the thicket or onto a bit of driftwood, or when it is flushed. At best it is a difficult and tedious task to find these birds. The best place to observe this bird is along the Willamette River along the stretch mentioned above, particularly below Peoria on the Benton County side of the river. By boat the backwaters are easily accessible. The birds are most often found perched about on the driftwood that piles up in back eddies and on the bars. Most of these points are not readily accessible from land. These birds undoubtedly nest here. There is no reason why this bird should not be as common along the Columbia River and its sloughs as it is on the Willamette River. There is certainly no reason for listing it as a "rare" form from the western and southern half of eastern Oregon.

In addition to the above records I have observed this heron at Coos Bay, July 4, 1942. Dr. K. L. Gordon of the Department of Zoology at Oregon State College says that he has always considered it quite common in the Willamette Valley. He has also observed it along the river on the highway between Grants Pass and Medford. The notes (unpublished) of Fred G. Evenden, Jr., show some interesting records in regard to its arrival in the spring and its late stay in the fall. The following records are from his notes:

Nov. 9, 1941, Toledo, Lincoln County, Oregon.

Nov. 5, Shelton, Washington (on Puget Sound).

Mar. 19, 1942, Corvallis, Benton County, Oregon.

March 19 and November 9 appear to be the earliest and latest dates on which this bird has been observed in the western part of Oregon. It is quite possible that along the coast and on the Willamette River it may be a year-round resident. In addition, Mr. Evenden has observed green herons as follows: "Aug. 7, and Sept. 21, 1940, Pudding River, Woodburn, Oregon; July 13, 1946, Croison Gulch, SW of Salem, Oregon. One was observed on July 27, 1946, on the cornice of the Commerce Building on the State College campus." His notes also show numerous records in Linn and Benton counties in the areas already mentioned, and his own comment is that "Anthony's green heron is more common than the great blue heron in this region." In view of his familiarity with this region and his thorough-going observations, this comment is worth noting.

While these notes are in no way complete as to distribution and habits, I hope that they will be of some help, particularly to the student, in dispelling the erroneous impression that the Anthony green heron is a rare or casual visitor in the state.

WM. GRAF

Arizona State College, Tempe, Arizona.

of the young.

July 22, 1946 - Poison Oak L., One abult was on the nest of local. when we approached, and as we watched another file and our hold by, the closely worked its vay from tree to tree in a chicle until it was in the nest tree, and the bird on the nest to defit and a which to the auge of the tree and file to a single bird to to away. The other bird then stayed at the nest. The bird to the been on the nest spent its time preening in the sun. I saw the single from the local way will expect an long as it takes the free mate to fine some food but the power. Hen we finally approached the rest, the full stay of take close and printted us to take some mivies of it in a murby oak tree. In just a flader up to the nest, and were I locate in this time, the power are perched around the cape of the nest. Also hence are 1 eyes. In the vote of robin's-eye-blue color. They were after into the activity.

bust time, only counce also be seen. It we quite that tipe, for it it is a deeper gray, and pin to there a to b/4" large on be seen growing on the wines, etc. On sine has the crop protated for a real could as if it has just been for. Fotal can the blood bill to to in might be 6-7" now. One bind is just a faith real, or the your would have also out by now. The leaves, branches, and pond rillies celes the real size thoroughly "whitetas as". The top off of the next is quite rilingy, and aerested, as you can be see below the next, and see the gains in the next OK. It is a solid mass of twiss under the center of the next, however. The years of these in the next it. It is a solid mass of these in the next it. It is a solid mass of these in the next it. It is a solid mass of these in the next it. It is a solid mass of the real and next it. It is a solid mass of the pairs under the center of the next, however. The years of these in the next it. It is a solid party of the pair of the right and needs held attraight to, also.

right and necks held offraight of, also.

July &c, 1946 - Stewart Lake of Young perched on a snag near treduck from the www.

July 25, 1946, Poison Oak Lake of 1 - U.S. Worked, clear and not.

Adult on the net at 1 poison Gordon worked around the lake fit camera

Adult on the not at 1 j.m. Loc Gordon worked around the lake for camera and tripod, and the bird, rathonal nervous, states of the last ath him only 15 feet away. As he approached, the young, hide has been at haing on the edge of the nest, on ear rice of the adult, the sallt rander the adult bird. For took jies, rever clopes, and took even a five return I worked up to and the nest to get action out of the bird. It rose from the nist, and at thed long the limb into the center of the trac, and then flew into an adjoining fir tree.

The young are juite active in the nest, and jeach upon the core frequently, sending a almost account their bullance, but somehas regularing it awkwardly. The jie leathers are ready coming out a st, and the down is almost a blue-gray not. The skip is that yell ich is a ces, and greenish in others. The legs are a yellow-green. The inter is a gelden-brown.

e took movies and stills at the nest, and then I reached for the

birds to bind the . The cheer of the two took on the opposite side of the nest, and walked out on a limb about a foot or to from the nist, and wouldn't come back to where I could reach it. The other one hard to the nist sticks tightly, and wouldn't release its hale unthe I let it ching to my fingers. There are still I have in the next. The eggs be no good, for they certainly could have betched by now. Poison Oak Lake? 8 to 11 a.m. Chimbed to the next again, but both young ran out on himbs away from the nist, where I couldn't reach them. The parent bird circled around close by over the like, and landed in a nearby oak, uttering its "unk" "unk" nates. I gove up and a turned to the nest later, and the joung were back in the nest, and the coult left the nest again. The claest one (the one I aidn't get to bard on the 23rd) went out the very same limb all three times. However, the younger of the two birds went out different limbs both times that it retreated. I nocked the limb up and down and eight the limb went

July 25, 1946

Reprinted from The Auk, Vol. 64, April, 1947, pp. 322, 323

Anthony's Green Heron nesting in western Oregon.—On July 18, 1946, a nest of the Anthony's Green Heron (Butorides virescens anthonyi) was found at the edge of a small lake in Linn Co., just three or four miles east of Corvallis, Oregon, in the heart of the Willamette Valley. The nest contained two downy young, of unequal size, and three eggs. Kodacolor stills and movies were taken of the adults, young, and nest on July 23 by Dr. K. L. Gordon, Head of the Dept. of Zoology at Oregon State College. On that date, none of the three eggs had hatched; as was proved later, they were failures.

The nest was situated about 12 feet over the water of the lake, in an ash tree (Fraxinus oregona). The nest itself had a dense shade canopy over it, even though the tree was in direct sunlight every afternoon.

Previous to this, on July 1, 1946, I found adult Green Herons carrying food to a nest (which I could not locate) on a small pond just two miles north of Corvallis, in Benton Co. However, on July 17 I did find this nest which was situated 15 feet over water in a willow (Salix lasiandra). This nest also was on the eastern side of the lake, but well shaded and not as exposed as the nest mentioned above. On July 23, I saw one half-grown young perched on a snag on this lake, evidently from the empty nest I found on July 1.

On July 27, 1946, the two young from the nest found July 18 were well feathered, and they left the nest by July 31. The nest was found torn up and with the three eggs floating in the water below where it had been, on August 2. The pond lilies and willows below the nest were thoroughly "white-washed" by the two growing young.

When I visited the lake on August 16, I watched both the young, now well developed, feeding in shallow water. The leg bands I had put on them were plainly visible when they perched on snags at the edge of the lake.

These two nests constitute the very first actual nesting records of this species in Oregon. At the date of publication of Jewett and Gabrielson's 'Birds of Oregon,' (1940), they stated that the bird was a decided rarity.

On May 25, 1944, a young Green Heron was found in the city of Portland (Jewett, Condor, 47, no. 5: 219, 1944)—the first published proof of the Green Heron breeding in Oregon. My unpublished notes show that on June 28, 1942, I found two half-grown young, barely able to fly and attended by the adult birds, on Horseshoe Lake, near St. Paul, in the northern part of Marion County. It is quite evident that in recent years the Green Heron has become much more common in Oregon. It is surprising that no one has found a nest before this time.—Fred G. Evenden, Jr., Dept. of Zoology, Oregon State College, Corvallis, Oregon.

July 25 continued: down, the young instinctively agreed its wing, undeveloped as they are. Finally I took a long stick and got it to perch of it, and then I lifted it in to me. I tack it to the round, banded it, and took gics of it. The wing pinions are chaing rapidly, as both primaries and second ries have aroun considerably. The primaries on the oldest bird are about a linches long not. And then I held it, it tried to peck at me with its long high. The counties turning carker and darker every day.

July 27, 1946 - OSC Campus. 10 a.m. I watched one circling over the compus, and it coon come to rest on the same of the Commice luin, in radial cornice.

Aug. 2, 1946 - Poison Oak Lake -0 Sal one acult carrying a 5" find in its bill, is it alit on an oak snace. I visited the late and a chapt hay and to any next I dutch, and then is away for a bille, it all the up, no the duding a to all was left. No sign is the year of a neglect to they have left the next before it was torn up by thoever, as whitever did it. All the vegetation below the next was thereup by "filte block" of the birds, companie to the up and the propert when last I visit a " the the dututs seem to have opecial traces that they habit, one has a levice they which they move then they conk their any around the lake.

August 15, 1946 - South Santiam I. Carcacia At Sweet home 4 p.m. Same need forcing on low water exposed became in migration.

August 16, 1946 - Poison Oak Lake 64 8-8 a.m. Sar both pound birds it at I bended. They had nearly full scult plumage, with one allohtly less developed than the other. Saw one saw to 0 d Greek, on the way both, the.

August 26, 1946, Poison Oak Lake, 8 a.m. In young one percheu for the ball small small small sticking out of the water. I approached within 50 fort, and while I stood there it became very nervous, shown by the flocking of the tail.

Sept. 11, 1946 - Every Park, Corvellis fluche and Store the at both to the river May 15, 1947 Stewart Lake, one perched on fallen oak tree with 4 Tood Fucks and 15-20 Mud Turtles

Sept. 7, 1947 Corvallis Gravel Fits for a temperature of the case of the five of the gravel bar. It other into the bire letter of its control of the five for the five form the five for the five form the

Sept. 16, 1947

Noting there, as is trive, to expect to be a considerable for its true of the line of

July 1947 - Ban Printle place at thurston, Lane Co. C. N. Nest at 15 'in alder on N side of alough off McKanje R. - Theft in July.

June 23, 1948 - Ban Printle place at thurston, Lane Co. 1 630 p.m.

Next on 5 side of alongh off McKenjie R., 100 feet from (W. of) last years next.

Next 8' over water in layer, on horizontal kimb to NW, about 8'out, too. 4 young, about 3'2-4" long with fill about 13," long.

July 25, 1948 Cottage Crove Preciveire Coult at 189 in . . in a princation or st area, but not over 1 years from a sum or , breather, or nest-inc. . to verseking root I have a returned to the content of the content o

Sept. 8, 1948 Sacramento Refuge: very common along the Logan River" or ditch, few being seen in the true marsh areas. This may because of the short willow growth, etc., available along the river or canal. No evident imms. seen.

May 14, 1949 Rocklin, Calif. I flew over hiway June 3, 1949 Corning, Calif. 2 pm. "" " Park July 5, 1950 Redains Calif. 12 moore, " " Park July 12, 1957 - Sev. along Cener. R. man carlon Party South, Calif. 1500 el. 7/39/62 - 1 ca with in a commerce. R.

5'cy og . Kins genge and a calleton meet of eggen College soliculate @ 32-24. ing Mangange and addition of the same of t

5/13/-19

Location and obrerve tions Lute

On hi heay to domes : " he care rec. 0, 1941

Sept. 8, 1948 Sacramento Refuge: exceedingly common, both adults and immatures. birds were commonly seen perched in trees, wherever available. Immatures werea quite uniform gray color, with a buffy curved line running out the wing, but no other distinctive markings.

Sept. 25, 1948 Grey Lodge Refuge: none seen over the refuge till 2 p.m. when I found a heavy concentration at the west end of the "loafing pond". I'd estimate there were between 150-200 individuals. I could see over 50 perched on the Scippus from one position. I flushed them by waving, shouting and clapping. Before I did this they sounded like a chicken-yard full of hens the way they chattered among themselves. This is the most I have ever seen. Contrasting these with the ones on the Sacramento Refuge where they perched in the taller willows, etc., at the edges of the ponds.

Grey Lodge Refuge: in the colong area where I saw 150-200 of them before, Oct. 23, 1948 I put up at least 3500-4000 of them! Adults and immatures. Iris red-brown.

Jan. 8, 1949 Gray Louge Refuge: just 50 or so seen. Big colony gone.

Jan. 29. 1949 Cyperus Point, 17-mile Drive, Monterey Peninsula - one on wave-washed rock at edge of tidepool just north of the golf course.

Humbolat Bay, Humbolat Colfone seen on bay mua flats at 430 p.m. April 5, 1949

June 22, 1949 Bishop Cr. or metropolis Per. 1 of Tells, 11ko Co. Tovalla: 4 just priched on cloating log on unstream tace of gam.

June 30, 1949 Stillwater "ilalife Mgmmt rea, Churchill Co. Fevana: Jane 1 10 ly

July 24, 1949 630 p. n - Yolv By - pass! Amall groupes flying 5 & to roost ?)

Nov. 22, 1949 Biggs - Butte City Book, Butte Co. H. Butte recht. Many Minch warms partled

Mughes Helew 14d. 7+ um. Six calling awashed typolad,



Least Bittern

Ixobrychus exilis hesperis

Sept. 8, 1948. Sacramento Wildlife Refuge: a new life-lister for me ! Brownish, with rusty wings, and not over 10" high. Would alight in tules, when flushed.
6/18/66 S and U of // Imokalee, Florida. 1 on Fire line whove sid

Observer Freu ivenden, Jr.

late Location and observations

May 18, 1941 It mile swamp south of what this colline to all like a made to May 24, 1941

If mile , want couth of torver his to not the control that the state of the

have no noted to interpreted to the process.

In mass of the first to the first transfer of the first transfer Long to which am of the first of the first the first terms.

May 12, 1942 May 31, 1141

12-mile Lump Co Nest w/ 4 aggs, mode fleed sedge, sloced in the water in a charp of catholic on World and

Leader Composition of the compos

Mc-factor to-mile or ofth cut to time. Linch 6, 1147

Militaria Con on the tries to prove of while level in the control of the year, at the Oct 10 1944 edge of the bank along a stretch of the river.

12-mile Swamp: one in the spike-rush within 25 feet of Tom McCallister, May 13, 1947 but it wouldn't show iteselt until have and I came up to the road toward the bridge. It flew to the NE side of the swamps where I mater Thushed it again.

July 24, 1947 12-mile (wamp, 550 a.m. s pair along trop the rath to the north end or the swamp, one behind the other.

12-mile Swamp: While eating lunch in the jeep on the road just east of the Feb. 14, 1948 Muday Cr. bridge I noticed something in the road which looked like appaper sack that was open. Checked it with the binoculars and found it to be a bittern standing still with neck pulled in. A minute later it began walking very, very slowly and deliberately bck into the swamp. The movement was so slow that the bird's camouflage was still effective even on the gravel. Eventually it went back into the swamp, but several minutes later I noticed it back in the road. This time it moved slowly toward the other side with neck stretched almost to a totally horizontal position. Just before it got back into the swamp, a car came along and at the last moment the bittern awkwardly took to the air - this ob by Dave Marshall.

12-mile Swamp. I also saw it at 330 p.m. just to the west of the second Feb. 15, 1948 bridge, at edge of the road, it flus ed to a 8' Ash "unloading" as it went, then flying 100 feet or so into another ash area. This was just a 1.0' from

where Dave saw it Saturday afternoon.

Secto Signal Depot - One in pool in field - 9a.m. april 4, 1949

July 11, 1949, Willows-tra Feyry, Glenn Colling one observed their with much wice open - not. Nov. 16, 1949 Butte Creek, along 1004 Levee, N of White Mallard Euch Club, near Gray Dodge Refuge - observed 2 at open eage of aitch, sputh//. They troze, but would turn head as I worked around them.

May 30,1953 SW Corner Lake Tahoe, topeSeach Marsh - 5 n.m. 2 seen, also eard pumping 31st June 27,1954 Pope Teach Larsh, whic Lohoc Wallife. A configurating could uplied to 15 miles and 1

٠-- ١ r .

Plegaci: quarauna

June 24, 1949 Fallon Irrigation area, Courchild Co. Nevada. First ones con-75-1.0 in mar shy lield 5% nines (of Fallon. On Februa, and fl. ing. About 250 others seen in area.

Scarlet This - Onara rubra

May 18, 1965, Greynolds Far., Teiari, Dad Co. Wig. Although the white-bbis is the Andmost common bird, there were 3 scarlet ibis seen, 2 notes of young.

6/19/66 Greynolds Tark. Nicki. Watched Whits-Jearlet Arbrid of rust. It was a Told mink

| | us columbianus |
|---------------------------------------|---|
| Date | Location and observations: |
| | Taft Bay, Taft, Ore. We saw from 8 to 12 of them on the far side of the bay, near the sand bar. |
| | 6 - Taft Buy 330pm. 9 of them at the south end of the buy, just a short distance from the highway. |
| | 47 - Lake Lytle, Tillamook Co: 3 birds feeding along shore at south end me calleted - Boca Reservoir, Nevada Co: probably a whistler - a single immature bird along the west side of the reservoir, in a wind-protected cove. Very little pure white, most of body appearing grayish. Head and neck appeared to be tinted a golden-brown color. Observed at 200 yards, so details are lacking. It did seem the head profile had much of the canvasback head profile. |
| | 8. Gray Lodge Refuge, SW Butte Co about 150 on the area. 9 Gray Lodge Refuge: close to 1500 of them on the refuge. They were a |
| Jan. 25, '49 Feb. 6, '49 Feb. 6, 1949 | beautiful sight flying against the strong wines. Many were calling. 12-mile Swamp (IBM) first saw four abults which came out from behind a seage clump in the channel next to the island south of the road. They didn't see us at first, but soon took to the dir, directing around behind us giving us a good view. Upon taking flight they uttered some notes similar to the honking of geese (canada). The found three others in behind some seage and spired in the big pond at the same location. These birds were purping their heads and necks up and down as other waterrowl do in courting. They flushed and I paced the distance from where they took to the air. Fach group flushed and I paced the distance from where they took to the air. Fach group flushed at about 100 feet from us. I have reason to believe that these were not the only swams on the swamp, because I later say what I thought to be a young one from the road. It was in the sedges southwest of the bridges. Grayish plurage. 12-mile Swamp of saw the 7 again (IPM) at the same location. To be of their calls were almost human-like. In addition to those given last time, the piras gave low whistles. They again pumped their heads up and down, can ing white doing so. All were adults. Sacramento-Gray Loage Pefuge is aw 2 flocks of 150 and 85 restine and feeding on bare fields, usually freshly proved. Gray Loage Refuge: perhaps close to 6-700 seen scattered thru refuge area, the other large flocks were seen outside the refuge in large open, plowed fields. They taxled out the water within 10 feet in this wind. Called often. Was |
| Feb. 27, 1949 | tough flying, even for them. |
| Jan. 30, 1949 | just south of the road south of the swamp. 12-mile Swamp! At dusk three adults circled in and lit at the edge of the water directly in front or our brind. The only sound they made was the loud rattling roar of the wind in their flight pinions as they landed. They spent their time feeding and preening. Then it was dark, we stepped quietly out in front of them, but they were not the least bit disturbed, neighter was a flock of nearby Canada Geese. Only the marriers become anarmed The |
| eb. 1, 1949 1 o e o t | 2-mile Swamp. Farmer ship they let appeared there Jan. 27, and had been there every day since then. The first few days, this was during the freeze, several of the birds came up to his house and were so take he at empted to approach them with a pan of grain. They were also eating shoots or rye grass. 19 - atilityater teture area, which is a cov shout in seen in NE section by Fred wright just the day before. |
| | walker take, mineral Co. Levada - small groups of about 6/group scattered over the labor. Use them of the shille and distance for a number seen was about 75. Occasionally a couple of the groups ould swim together, making groups of 12-14. |

Jan. 11, 1952 Hood - Franklin, Into 6, Caly . - 7 Coche of 10 - 25. 35-405, RY

768.13, (959 - 1 hoge the - 25 White.

208.15, 1859 - Denivitating much flat of fields of Common K, South to city. - 45.50 me.

2/4/13. Gene Tarm 10012 june of 12 deal copy, de 2, 1900 deal.

is a small construction of the first of the state of the

.u.b-r :: field to ...

Jan. 17, 1945 12-mile (want) sout 1 if corvainis/2 .

approximately 2 or 0,000 of these and 10- tesser bandues on the crea diverse toda, .

Wallula, Wash., along Columbia R., got up to within 30 feet of one bunch of 27 along the bank of the river. Dec. 3, 1943

Apr. 1, 1944 Glenn's Ferry to Hammett, Idaho - Floating down the river and on bars, etc. They all seemed to be traveling in pairs. In one instance a third party tried to join a couple and it was driven off right awayy.

Corvallis: ()*2 strings, totaling from 250-300 birds llew over at 3500 feet. Apr. 23, 1947

Apr. 24, 1947 Poison Oak Lake: 3 flights went over, heading north, all in long inverted "V". 275 in one flock, 150-160 in another and 225 in a third, which was above the clouds.

1)-mile Stamp: $^{0.7}$ I thought for the grass had left, but too, their Apr. 13, 1948 numbers probably run between 2,000 und 8,000 birds. Subspecies?" ""

es t side illy with Dake 12 b so h. July 27, 1948 July 27, 20, 1948 Tavis Lake: Card in tourn. 15 on bur it. of non-ruen on the balance a comment of the in-

12-mile swamp. at dusk 4 lone geese came over and circled the area mapy Jan. 30, 1949 times, and then after settling up a wild clamor ous call, settled down. Shortly after this the main body of the geese came in our the fields in small groups until well after cark, at which time the birds had all settled in this one small area of water. Apparently the linst lone goese which circled the area cautiously acted as scouts.

April 30, 1949 12-mile Swamp (IBM) in the time we were at the swamp (9:5-14.0) there was noted nine flocks of geese migrating. They came over throughout the time we were present. Flocks collosed of following numbers: 35, 20, 800, 250, 300, 120, 250, 60, 50; total: 2065 Rone of these binas stopped. There were, however, 7 Canadas at the far south end of the swamp.

. I tak in the - 5 in longitud Sept. 1, 1948 Tule L. Calif.

Nov. 29. 1950 Hogan Res. Calaveras Co. Calif. 225 (one flight of 36 came in from the valley while I was there. Mr. Ballard states that they heave the res. as late as 10 p.m. returning in the a.m. The goose le ve each day before the aucks. Lucks seem to leave in one large group

Abert Lake, SE side, Lake Co. Oregon. 3 pairs, with 2, 5, and 9 young. May 26, 1951 While on the bank, the young were mixed in a loose flock, but they segregated into their right groups as they moved towards the water at our approach.

Summer Lake, Oregon. many seen, and many with young, mostly of larger size May 26, 1951 than those seen at Abert Lake, but still with down, gray. 12 adults alone. Also 7 pairs in flock with 24 young. Young seen in family groups of 4, 4, 8, and 3, with pairs of adults.

June 11, 1951 Derby Dam Forebay, Truckee R., Washoe Co. Nevada - 2 pairs of acults with 4 young, 2/3 grown, and with 6 young.

June 5, 1952 Carson Valley, Douglas Co. Nevada - 38 adults and a total of 47 young, Cl. II, grouped as follows: 3, 2, 3, 4, 5, 7, 3, 3, 6, 5, 6.

June 27, 1954 ppc Peach Spit, 1 No inhec, Calif. Much sign around, and old nests of

this year found.

Aug. 7, 1963, Canyon Jct. wen, Well water eNFk, Ayo. pairs and family groups alone Wellowstone R. above canyons.

11, the second of the second o

Observer Fred Evenden, Jr.

1469-1485-1580-1617-1779-1819-1825-2035-2293-2657-

Late Location and charryctions

12-mile swamp south of Corvallis! Jan 17, 1945 Approximately 1 or 4,000 of these and the Com on Schaoli on the area covered today.

Klamath Lake, Ore.

Dec. 3, 1943 | Wallula, Wash, along Columbia R. - not many compared to the common Canada. Sept. 24, 1946, Toodburn, 10 a.m. the I inhte of them heading the mark:

- Jan. 29, 1947, 12-mile Swamp. in a flock of 3000& took off from north end of swamp in a mass, and split into 2 groups, one heading NW, while the other circled and landed in field east of swamp, near the hiway. Flushed them back & forth.
- Feb. 4, 1947 12-mile Swamp - approx. 1500 flushed from a field south of the swamp, when we were still 200-250 yards away, but all flew over us, as they flew to the north end, where we flushed them about a p hour later. It was a thrill to see them in flight, and still more of a thrill to hear the crescendo-like roar at the beginning of the flock taking to the air.
- 12-mile Swamp there never have been the numbers around this area that there Feb. 11, 1947 were today. There were between 6,000 and 10,000 altogether. "Triangle flight" They were very restless, and altho we did not flush any, a passing train did, for the geese were on both sides of the track in most of the fields, and a Cub plane flying in high from the north put them all to flight. No matter where you looked, near or far, strings of geese could be seen. It was a thrill to see them in such enormous numbers.
- Feb. 18, 1947 129mile Swamp: we flushed on single flock at the north end of the swamp. This was a thriller for there must have been at least 3000 to 3500 in the swamp, and then they circled and most of them came back over us at low altitude. We flushed them again after they landed in the next field north of the swamp. For the first time I had a really good chance to see how geese lose altitude when they come in to land. They roll over on one side until their ext nded wings are vertical, and they slop downward some 20-30 feet, before rolling their wings back horizontal, but they don't stop there if they plan on descending further, but continue to roll on over to the opposite side until the wings are again vertical.
- Feb. 25, 1947 12-mile Swamp: small flocks in shallow water at the north ena, with the bulk of 2 or 3,000?? at the south end, on grassy land, just seemed to be basking in the sun, for they weren't feeding. Just the sound of airplane engines on the flyway to the east will put some of them to flight.
- 120mile Swamp: 0 2500-3000 in flooded field north of the swamp. We did not Mar. 1. 1947 flush them. They were preening, and many were entending their wings and beating them vigorously, or running a short distance while beating them.
- April 8, 1947 12-mile Swamp: (approximately 2000-2500 were inone or two flocks at the south end of the swamp and an adjoining field. Most were Lessers with some Commons with them.
- April 15, 1947 12-mile Swamp: We though they had all gone, but we found one trying to sneak away thru the marsh vegetation at the N end of the swamp. Finally it flew. Whether or not it was sick, I don't know, but it did have trouble getting into the air, even against a strong N wind, and then it circled downwind, to
 - May 6, 1947, 12-mile Swamp 1 Tom flushed just one from the north part of the swamp. It may have been the one lave and I flushed about a month ago, but it seemed to be in apparent good health, while the one we saw was not????
 - 12-miles Swamp(still the one on the swamp. Flushed it 3X May 13, 1947

* */ . . Nov. 6, 1948 Hogan Res, Calaveres, R: 800 or so. 2 had white wing primaries.

apr. 13,1949 - Jacto Jignal Depat - 845 a.m. 3 large stongs flying north, calling lookly.

Dec. 3, 19 9, Failing Levate, near bruce. Lenton Co. Oregon. A. DB...
estimated 3,0.0 in all. They kept coming in and randing in a field directly north of rimeon Butte. Lere estimate shoots of rye grass. Trobably not one than 10 cacklers se was though many wore were undoubtedly present.

May 1,1959 — S. Corrollis, Organ, N. Jam 12 mile Swange, 1000 = in talls.

1/41/76 Nime Blade stormed that Army length.

1/41/76 Nime Blade stormed that Army length.



Observer Fred G. Evenden, Jr.

Cackling Goose

193-257-26/-266-373-440-486-501-516-575-599-605-688
1029a-1031-1043-1049-1052a-1069-1071-1000-1192-1217-1340-788-783-786-794-786-827-844-852-857-868-873-424-947-942-942-1919-1000-1191-1217-1340-7348-1350-7365-1367-1375-378-13794-1417-1429-1448-1527-1580-1617
Branta Canadensis minima

| Late | Location and observations |
|---------------|---|
| Apr 5 1946 | Corvallis, 830 a.m. A flock of 60-70 flew over the campus, heading due north. They ware calling loadly. |
| Feb. 11, 1947 | 12-mile Swamp of a few small flights of 7 to 15 present. Small size noticeable |
| Apr. 3, 1947 | Stewart Lake, Corvallis! A single one at the south end of the take in company with a single White-fronted Goose. |
| Apr. 24, 1947 | Corv, MacDonald Forest: a flock of 150-175 flew by about a 2 hour behind a flight of Lessers |
| August 8, 194 | 7 Fern Liage Lam: we flushed one from the PH side, and it layed about close, nover leaving the vicinity too far. |
| | Lavis Lake: "The cipin. |
| May 5, 1959 | Gray Lodge Refuge, Sk Butte Co. 25 flew over. They usually leave by now, with |
| Sept.21, 1950 | suragglers present until about May 15 according to Cowan and Cloya. Doodburn, Oregon. heading south at 930 p.m. Calling. |



Branta nigricans

Late | Location and observations



Black Brant, Cinnamon Teal and Yellow-headed Blackbird in the Willamette Valley, Oregon.—A rare sight, indeed, on October 23, 1941, was a solitary black brant (*Branta nigricans*) standing at the edge of a winter pond in a stubble field. The pond was about 100 feet east of highway 99W, just a few miles south of Monmouth, Oregon. Although I have seen many black brant at Newport, Oregon, on Yaquina Bay, this is the first, and only, one I have seen in the Willamette Valley.



Mar 26, 1947 Netarts Bay that night while camped on the edge of the bay, we heard large numbers of them coming into the bay shortly after dark (T. McCallister) Mar. 27, 1947 Netarts Bay: Afflock of 200 out on a mud flat in the bay. Several strays feeding on flats along shore (Tom McCallister).

May 18, 1947 Seal Rocks, Lincoln Co. - a flight of 130-140 flying fairly high on their way north.

Mar. 19, 1949 Tomales Bay, Marin Co. about 150 to 2000 feeding on eel grass in bay. Mar. 19, 1949 Bodega Bay, Sonoma Co. about 250 to 3000 in 3 or 4 small flocks.

Apr. 25, 1950 Humbolat Bay, Calif. few in north bay, but I made three counts on calm water in the South Bay early in the a.m. and the total averaged out to 35,000 birds. About 130 p.m. I was at the end of the north spit, while strong 30-40 mph winds were blowing from north and I observed several flocks of 150-200 or 300 brant heading north over the ocean to the west of the south spit, and then veering in sharply with the wind, as they reached the bar entrance channel. Perhaps these were north-bound migrants moving into Humboldt Bay from the south, and seeking haven from the winds.

Dec. 7, 1950 Morro Bay, S. Luis Obispo Co. Calif. a total of about 9,800 in 3 main bunches in east and central bay areas. Localities state the brant arrived suon after mid-November. Depart late Meb. and Larch.

April 18, 1951 Morro Bay, Calif. Mt. 610 a.m. Most in central portion of buy, some opposite fish piers downtown. Some fed at base of cliff in park. Others feeding over bay area. Some moving in from south and a few out to north. Total about 8,500 birds.

April 19, 1951 Pt. Pinos, Pacific Grove 1230PST - 74 on white-sand beach below Asilomar. Others flying and 25-30 floating offshore opposite rocky joints. Also feeding. Observed some territorial behavior with those birds on white sands.

April 20, 1951 Pt. Pinos, Pacific Grove - 15 on beach, 25 offshore, others in tide-pools.

April 19, 1951 Moss Landing - 5 in bay on mud bar.

April 21, 1951 Pt. Pinos, Pacific Grove, Calif. 10 in rock, tidepools.

Mar. 12, 1952 Moss Landing, Monterey Co.Calif. so oty brown, with neck ring incomplete, very white beelly behind the legs.

Mar. 22, 1952 Bodega Bay, Sonoma Co. Calif. 4,000 or so-as tide went out they moved to open water and grassed hills E of bay. Many people around bay this Sat.

Feb. 3, 1953 Morro Bay, Calif. in a 2-hour period this a.m. by Ralph Imler and I, a total of about 7,200 brant were counted.

Apr. 18, 1954 | Ft. Bragg-Westport, Mendocino Co. Calif. 4 north along coast



Branta bernicla hrota

Light-bellied Brant

Apr. 25, 1950, Humboldt Bay, Calif. Among the 1,000's of black brant were at least 7 that I feel certain were this species. They were seen in flight.



13269 111-148-192-467-516-578-598-747-848-957-984-992-1011-1031-1043-068 2035-2239-2434-2657

Anser all arrons abbitrons

Thate goose -+- 1043 - 1071 - 1432? -

Late

Location and observ tions

Nay C, 1948 | Francis transposition of the state of the s

noise on a little of the littl

Stewart Lake, Corvallis: a singe one at the south end of the lake in Apr. 3, 1947 company with a Cackling Goose. Both flew together to the north end of the lake. This peculiar association has been noted before at this lake.

12-mile Swamp: We were surprised to see 15 of them, not at all shy, Apr. 15, 1947 waddling and swimming around in the north half of the swamp. We later flushed them, and they flew to the south end.

Li-mile, $a_1:C_1$ involues it mentions in the substitute of the state of the substitute of the subst Oct. 7, 1947 They circulate even to a round the ground and eventually events to be stubbac in ac mean, this the factor of the best

Grey Lodge Refuge: 15-20 seen. Others heard. John says they just came Sept. 25, 1948 in the other day.

Gray Loage Ferrige Ma who S seer, and J In come says they are conjugate Aug. 16, 1949 that have been a decrete in the line and an algorithm to be a tag week with the freeding of the

Wingfield Park Bano, Nevade - One bird, evidently of this ageins with Canadas & Lease Canadas on ice at edge of open water in trulece R. -? subspecties - or hyprid? John dependshi took paic. Had gray back side of face + head, but otherwise looks like white fronted on tule. - Quite large. Jan. 10, 1950

Chen rossii

Ross's Goose

Dec. 29, 1948, Grey Lodge Refuge, Butte Co., Calif. 230 p.m. a new life-lister Saw a few, no evident grinning patch, and could be slightly smaller than the Lesser Snow; also a lighter call, and flight wing-beut.

7-08.4, 1953 Marced W. M. A. Marced Co, Caly. Rolph hales; Espaye Max. Gene crawford & of found to agreeine in excellent condition, but dead on the ramplants of Mr. W. M. A.

Dec. 11, 1075 - 3 of 10 doe format, Oalii. It is it a wing a light of the W. M. A.



| Chen hyperborea hyperborea | | | |
|--------------------------------|---|--|--|
| Date | Location and observations | | |
| Oct. 10, 1942 | Laurelhurst Park, Portland. It lands on the water with its feet down and forward. They hit first and the wings are set, but as the feet hit, the wings are "back-watered". | | |
| Jan. 17, 1943 | 12-mile swamp south of Corvan is:00° in with a flock of Canada Geese. Thains, various over a half rare asa,. It tay so and to beliand a so in the core of Canadas and at war canadas. Its canadas not unificate that of the Canadas. Tear section to rave time yellowish or brown on it, as the bire care chose. In act when tags. | | |
| Oct. 27, 1943 | Washougal, Washington, Partly cloudy and wind was strong. There were 50 Lesser Snow Geese, and they were flying wing— tip to wing—tip. The left flank went directly above us about 35' high. The Snow Geese winged steadily on up the river, against the strong wind, for they were evidently heading for the Eastern Oregon Flyway. Later there were 4 more that came up the river alone. | | |
| rec. 16, 2867 | | | |
| Aug. 27, 1948 Dec. 10, 1949 | Socramento, Calif - 7:45 p.m. Aleard them calling + soon spotled 38 of them at about 800'- 1000' elev. They were a some pinkish white at that altitude, reflecting the lights of the city - They really stood out + charges in formation readily seen | | |
| Feb. 4, 1950 | Southside Park, Sacramento: "still one present, watched it probe down in the mud alongside a small clump of bent-grass until the roots were exposed, pull up the plant, take it to the pond, and dabble it in its | | |

then it swallowed the plant, entire.

bill until all the mud was washen off and the roots shown a clean white,



Blue Goose

Chen caerulescens

January 8, 1949, Gray Lodge Refuge, Butte Co., Calif: one adult seen in small open area in tule growth in the SW part of the refuge, with Lesser Snows. This goose is very rare in the valley.



lem rocycna biconor (helv.)

Menaota Pool, Fresno Co. San Joaquin Liver: item Lowing in sauce Aug. 2, 1949 and marsh areas all day, one farely not us in most or in to too, and he asked me if that was a tros-ough since he had never seen one, and knew I was especially looking for them - it to - the thite than band showing prainty as it took of, one as it turned the ruce; use reid was plain, and it uttered distle-like notes in all ht, and hang its neck extra low when alighting (bout 75 for this lare it int; ea. 3 more flushed from the same force of sreat the bakers. ThenI clapped my hands to Elush some more, but only still to an excests rose, and then a minute or two later, I more I usless from the supersuman the first four tree-cuers no fluched. One flow a coldet circle around us at a citting of about 10 years, but the other 5 more only short flight. They appears a just slightly forces than a myllama. Oct. 25, 1951 Mendota Pool, Fresno Co. Calif. 145-210p.m. a total of 115 seen, / pop of which 18 were not yet half-grown. They were perched on muddy flat at water's edge and swimming in water. Mixed with shovellers - when flushed, the tree ducks and shovellers separated into their own flocks. The air was full of the tree duck's high-pitched calls. Oct. 22, 1952 Delta Lendota Canal 1 mi. from S end - Fresno Co At/- 5 on canal in open.

 ℓ

July 11, 1077, 1, 24-67



nen lama chan a and the Control of th ... C. Trini ELCE that my the first flate, plans, the age of the first flate, plans, from from the first flate, plans, the age of the first flate, plans, the age of the first flate, plans, the first flate, for the first flate, flate Oct. 15, 1942 Corvallis, Ore. McFaquen Fond. 7:30 to 8:30 a.m. Gizzard 16 c.c. volume of contents Black Bindweed, Polygonum Convolvulus 14 seeds Wheat , Triticum vulgare 3 seeds Grit. 4.0 c.c. Unknown 0.3 c.c. Wild Oats, Avena fatua 6.0 c.c. Oat awns and pulverized plant parts 6.0 c.c. Craw 73 c.c. volume of contents Theat, Triticum vulgare 11.0 c.c. Tila Oats, Avena fatua 60.0 c.c. Undetermined l seed Annual Cow Vetch, Vicia villosa 2 seeas Barley, Hordeum vulgare 2 seeds

> Female: Gizzard had 14.8 c.c. volume of contents Theat , Triticum vulgare

> > Wild Oats, Avena fatua

Mild Oats, Avena fatua

Craw: 81 cc. volume of contents Theat, Triticum vulgare

Bur-reed, Sparganium (eurycarpum)

Grit and undetermined

Annual Cow Vetch, Vicia villosa

9.5 c.c.

0.3 c.c.

4.5 c.c.

78.0 c.c.

0.5 c.c.

2 seeds

3 seeds

89-806-809-810-812-814-817-817-819-822-

the second of th



1552 - 1552 - 1552 - 1555 - 1558 - 1558 - 1558 - 1559 - 1560 - 1561 - 1566 - 1573 - 1576 - 1577 - 1578 - 1586 - 1586 - 1586 - 1587 - 1589 - 1689 - 16

Anas platyrhynchos platyrhynchos

| Date | Location and observations | |
|-----------------|---|--|
| Oct. 15, 1942 | Corvallis, Oregon, McFadden Pond 7:30 | to 8:30 a.m. |
| | Immature male. | |
| | Gizzard had 9 c.c. volume total | |
| | Wheat, Triticum vulgare | 1.0 e.c. |
| | Barley, <u>Hordeum vulgare</u> | 2.0 e.c. |
| | black Bindweed, Polygonum Co | |
| | Grit Grain pulp, etc. | 1.5 c.c. |
| | drain purp, ede. | 4.0 c.c. |
| | Proventriculus: Volume content | was 10 c.c. |
| | Barley , Hordeum vulgare | |
| | Wheat , Triticum vugare | |
| | Craw | |
| | file Oats, Avena fatua | 3 seeds |
| | Wheat, Triticum vulgare | 136.0 e.c. |
| | Summer Lake, Lake Co., Oregon November FeMale: | 1, 1042 Shot in a.m. |
| | Large gizzard nearly empty or abo | out a full 0.5 c.c. |
| | Grit | ≎ C.C. |
| | . <u>Scirpus</u> sp | 1.5 c.c. |
| | Uaren sp | .5 0.0. |
| | Undetermined Plant Laterial | L.5 C.C. |
| | Immature Hale: | |
| | Giazar contents nearly rule, by | ut mostly muscle at that 9 c.c. |
| | Large pieces of grit 1/8" a | |
| | Eroken up plant material cirpus sp | 5 c.c. |
| | Carex sp | .7 c.c. .5 c.e. |
| | | |
| 9-4-43 | Cloves Carele, Mr Chord Field, Wash. | 2 2 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 |
| | I mallards pieleing up gravel a | long the road going into me don't |
| | Field. The made flew) I who | a differences. |
| Oct. 5, 1943 | South Tacoma Swamp, Tacoma, Washington | |
| , i | Observed mallard drakes and hens | |
| Dec. 3, 1943 | Wallula, Wash., along Columbia A., 1000's of them! | |
| Apr. 24, 1946 | 12-mile Swamp - Corvallis. Saw several pairs and found the nests of two pair. One had D and the other 11 eggs in the set. | |
| Apr. 26, 1946. | | |
| July 23, 1946 S | tewart Lake, Corvallis. Ca, tured a femal its way out of the trap, and it was in i | e this a.m. that coulon't mina |
| | only la inches long. It could not fly. | |
| Aug. 16, 1946 | Stewart L., Corvallis A They were present | in large numbers, asons with |
| Tau 0 3045 | quite a few pintails, and shovellers. | |
| Jan. 2, 1947 | Stewart Lake, Corvallis, ice was ½" this lF were on open water, swam to the ice wateddled everywhere. | ck on the pond, and 3 M and, climbed up on it, and |
| | | |




Page 3. Mallard

Feb. 11, 1947 12-mile Swamp, a rough estimate of approx. 600 on the swamp - equal sexess

Feb. 18, 1947 12-mile Swamp of pretty well paired up, and the first to take flight as we approached. There were between 200 & 250 of them today.

- April 8, 1947 12-mile swamp 2 about 25 or 30 pairs scattered over the swamp. They seem to be mated up but we saw no sign of any having a nest at this time.
 - May 20, 1947 12-mile Swamp: a few pair were seen, and also single hens. Tom and lave saw one with young, and I found a recently vacated nest of dried rush stems, situated on the south side of a Spired bush, at the base, in a solid stand of Carex.
 - May 7, 1947, Over Little Muddy Creek, Shof Corvallis, 5 miles. Tred Freck reports a nest in an Alder on the East side of the creek on a horizontal limb running west. Nest of grass remains of high water, is feet from trunk of the tree, and is feet above the creek. Shaded except part of the alternoon.
 - June 5, 1947 Stewart Lake, 5 p.m. I pair present the hen "flopped" all over the water, but I couldn't locate any ness, or young.
- June 17, 1947 | 12-mile Swamp Of Thushea a different females and all acted as if they had nests, but no nests could be found in the cinity from which they were flushed, and so it is presumed that young were around somewhere, the none were to be seen.
- August 8, 1947 Fern Fiege Tam Pseveral Thushee, and in ec ipse parage, for no Tal arakes with seen.
 - Feb. 15, 1948 12-mile Swamp: perhaps 400 all told most in field at north end, and due to strong wind, others not on open water, but in smelter of ash and spirea brush on east side.
 - Mar. 30, 1948 12-mile Swamp: Of seemed to be well distributed over all the swamp area. At least 350 pairs, as they were mostly paired up, and about equal distribution of sches.
- April 13, 1948 12-mile Swamp! maybe 5 altorether, most paired up. Not with take that was already completed (273). Towar ha an erropened 1/2 of longth, shelf membrane still present, so without it was predation. Shother auch nest found completed, but with no sign of the birds around. Take of it she stems of Garex oregonensis, caose to the N sade of a large clum of it, with the "alley" to the north about 4 feet or so.
- April 21, 1946 Out Cr-ek, Liem Co. It makes I of Corvainis nest in pike of organase left on some branches by high water. Nest about 15" over but if or horize ontal sing that had two connections to the north bunk of the creek. Cavity about 5-9" in diameter, and about 1,-2" deep. Quite a bit of shall pieces of down lining and ground the most. The nest had been ridged, polarly by a coon (or skunk), and the empty eggs (6 or 7) were in a pine on the grass about a foot from the nest cavity. The eggs core a greenish-thite, averaging 2" x 1 5/8" in size. They were all opened on the sic., near one end, and a few at both erds. That tip makes could be seen indice the shell.
- May 1, 1948 Fern Fidge Received: 1 p.m. a few, and found rest of 1 plan of T plan of dike, just 4' below the road atop the dike. Surrounded by tall grasses and weeds. 12-eggs. Took pic.
- June 13, h.48. lh-miles comple of the most care, and other not in the plumage. It lest a half as a resume surpure, and once town over note having families. I flushed 3 different crosses of journain various of the swamp, that a use not ify. These groups combained 7,0, and 11 totals. Some hour triangto use year among.
- June 1., 1948 Cottage Grove Leserveir: What flight of lo, also a pair with electric which we could not locate.



- July 8, 1948 12-mile Scamp: pair of acults are about 1. , up not yet a le to 11y.
- July 16, 1948 Fern Flago Reservoir: Ppair and 4 joung, the more control is line very little down 8 degr, at base of rose can in determine. East about 50 deep and 7" across.
- July 27, 1948 Fest side Klamath Marches ? pair with downy yours seen.
- Sept. 25, 1948 Grey Lodge Refuge: not many seen in the a.m., but this p.m. I flushed 100's in waves as I did the pintails in the a.m. 2000 in area.
- Oct. 23, 1948 Grey Lodge Refuge: 25-30,000 of them. Wave after wave would take off.

 Mixed with baldpates and pintails. Before they were flushed you could hear a continual hum over the refuge area wike a huge swarm of bees, from the chattering of the ducks.
- Dec. 29, 1948 Gray Lodge Refuge, 230 p.m. Over 1,000,000 ducks in the area, principally of this species, pintails, and baldpates.
 - Jan. 8, 1949 Gray Loage Leiuge: Over 1,500,000 ducks in the area today, principally Mallaras, Balapates, and Pintails (1).
- Mar. 7, 1949 12-mile Swamp Vall paired and scattered throughout the swamp, including the flooded ash woods.
- April 23, 1949 Grey loage Refuge, Butte Co. 130 p.m. 1 hen on nest of 8 eggs within 22 fect of passing cars on gravel county road leading into the reluge. Rest well-lined with down. No brush, or tall herbaceous vegetation near it.
- May 5, 1949 Gray Loage Fefuge. The April 23 mest had hatched out. John Cowan told me only 3 successfully left the nest. There had been 8 eggs.
 - July 31, 1949 Grane Luke, Calify, 17, ol. in semile pumage
- Aug. 16, 1949 Gra, bouge here restricted business the fine has been morth, and old females as the early part of the most, but the rettine canalizes, of same and morters.
- Sept. 1, 1949 Tile Lake, Cali. 1900's in arch. The only to pintails in bothlish hospital. Tany for makes that I have, the just be in its to gain pack for a of ital markets. Tone towers or meet on the interest of
 - Mar. 2, 1950 Bear Creek, Colusa Co. Calif. 3 pair seen one pair flew upstream ahead of the car at 35% m.p.h.
 - Mar.
 - Apr. 1, 1950. Southside Park, Sacramento, Calif. observed 2 hens on the park lake, with 10 and 13 3 or 4-day old chicks accompanying them.
 - May 8, 1950 Stillwater Wildlife Engmnt Area, Nevada seemed to be always a mare and female together. Total of 12 pairs of this kind, plus 3 mares and 4 unidentified. Total 31 mallards.

July 6,1950 Trivity R, S of Stringtown Trivity Co. 19 a.m. 19 in pool wear,

- Oct. 9, 1950, Wingfield Park, Reno Nevada 1250 PPT. 48M and 19F or immatures I watched one pair "neck bobbing" while a second female or immature was swimming nearby. All 3 were just above the dam spill sill.
- April 27, 1951 Southside Park, S acramento, Calif. 7 pairs of adults, some males inpartial molt. 3 unattached F or imms. or /?males and 5 unattached males. 6 broods seen: 5 were 8"; 4, $3\frac{1}{2}$ "; 5, 3"; 1, 9"; 3, 6"; and 4, 7" in length.
- June 12, 1951 Silver State Gun Club, Churchill Co. Nevada: 1F w/ 10-2/3-grown young another with 8 1/3-grown young.
- Aug. 10, 1951 Southside Park, Sacramento, Calif: 33 mallards, no full males, but many males with gray of wings and body showing.
- Sept 7, 1951 Southside Park, Sacto. Calif. No full males, but all with M color patches.17:15

 Jan. 9, 1952 contheside Per's, Danie. Calif. 2 pr. nec's del de
- June 5, 1952 Carson Valley, Douglas Co. Nev. 20M, & ad. and 6 pr. Class III broods: 8 & I4 July 23, 1952 Huckleberry, McCloud, Sikkiyou Co. Calif. 9 full males on pool in meadow
- Oct. 9, 1952 Lake Washington, Yolo Co. Calif. observed 2 M's neck-bobbing w/ female.

along creek in yellow-pine woodland area.



Nov. 7, 1952, Southside Park, Sacto, Calif: M's and F's grouped as follows: 182, 1 & 1, 2& 5, 2 & 2, 4 & 7, 2& 2, - total - 31.

June 27, 1954, Pope Beach Lake Talion Calif. 5 ... Dest Louis acon. For although white new of Legan. But how her one of C. . . lest Louis 10-124 dia. To provide all all for all formations of the formation of th

Popo Deach, Taito Intoe, Celif. 31, 1 poir, and I'm formation most with Nay 14,195/

12 eggs in sine clum about 3! hiel.

Nav 20,1955, Tope Boach, Into the could be made to another publication of the July 9, 1955 Aright's Take, 7000' ch. Ederado to Takif. The publication of the July 1, 1958, Aright's Take and the first of the second of the July and the July 17 and 11 on May 17, 1958, Pope Reach, Take Taboe, Calif. 25-30 pair, 10 cgrs on May 17 and 11 on May 17, 1958, Pope Reach, Take Taboe, Calif. 25-30 pair, 10 cgrs on May 17 and 11 on

May 16 in dense lodgenole pine woods, without any surrounding cover on bare ground. Neartest clearing by and 60 feet away. (odachrove bics)

May 30, 1958 - 8 pair seen, nests with 11 pale line-proce eros under signs. I also in willow thicket of otions and twins with toff-s adjetop some, and a semalcolored sens in set of Joinsus in dense six in 1990 the Act with a set

June 7, 1759, Yolo Byrass, Yolo Co. Calif. 1F with nine 2/3 grown young.

Jamely, 1869 Carlford, won als of danny going. lid if the to your you



New Mexican Duck
Anas diazi novimexicana

Feb. 20, 1951 Acala to McNary, Hudspeth Co. Texas. 12 on small reservoir with other ducks



Anas rubripes

Black Dude

Feb. 21, 1953 Aransas Refuge, Texas. Finally got to see my first black ducks - while they may be hard to pin down on the water, once they take to the air, the light under-wings are a dead giveaway. There were also mottled ducks there.



mas fulvigula

Mottled Duck

,

,

•

.

..te Location and observations

March 5, 1.40 4 miles et poi convelii de la careta

Oct. 5, 1943

South Tacoma Swamp, Tacoma Washington

One male flew in front of me while I was watching the G-w. Teal.

The gray-speckled breast and light abdomen showed up plainly.

Nov. 17, 1943 Walla Walla, Wash. 3-5p.m. _ a pair of Jadwallo with the matterds + Baldyates.


Black under tail overto, white wing patch in flight. White sally. What + sives appeared graziah.

Head has two brown shades to it. Took of directly into the iir.

Dec. 3, 1943 Wallula, Wash. - along Columbia R., _ saw 4 of them.

Aug. 29, 1847, Form liege lam. To flucted at least of the, milesot to the instruct, with lenty of place in the reculum.

May 12, 1950, Stillwater Wighife Mngant Area, Churchill Co. Nevaga: flushed gadwall hen from nest in solid stand of ungrazed saltgrass. 10 eggs. LBM.



Balapate

.... ii : . , _ ...

lante of Hield trig. " 84-110-111-112-126-141-152-187-191-192-193-197-206-210-211-212-253-257-260-261-262-263-264-265-266-267-268-369-270-273-274-275-

mithin at a stone me. I count to the to come a conthe still cover. In or a train of the very very line is not be trained was inconsticuous.

ocivalia: Office le como o le crime, a proposition de la como o la parato, whente, it is a company of the property.

corver his Officer of the our in paint of paint of the contraction of Maril a , and

Portland, East Moreland Casting Pool. Saw a raft of 500 to 600 Oct. 19, 1942 Baldpates on the water this a.m. They were mostly all adults.

Horseshoe Lake, St. Paul, Ore. 3-23-43 One brilliantly colored male that accomed quite time 4) . - 2 early to to levit lead, in a nowboat.

Walla Walla, Wash. - 8a. m. when + 30 mphylvinds from su Jan 17 1944 as I was coming home from the see, flight of ter right on the sound of wester mallards + a few Hald pates, were winging their way overhead in a su of wester direction. They were in small disorganized flocks + also in large v- formation. There must have been 2-3,000 in all. about 10-1032 u.m. They all seemed to do coming back flying in a NE & easterly direction. Perhaps they had all your too gravel red along some river, to get heir just require to the are the wanth to me two ray revisors.

Snake R. - Glemia Form to Hammett. 1500, about equal make & family acre in pairs + small flocks + large bunches all over the river + along the edges candes brush apr. 1, 1944 on the gravel bars.

Tillamook Bay Entrance: small flock feeding on rocky beach eage. March 26, 1947 Upper Tillamook Bay: large flocks feeding in grass at mouth of Miami and Tillamook Rivers (Tom Mc Callister)

Grey Lodge Refuge: 15-20,000 seen. Oct. 23, 1948

12- mile Livery, Beston 6. 40 seen in I compact flock. Just coming out of fall plumage. No calls. Oct. 15, 1948 Lover Alamath a condit. I row ray over harring to may than when Sept. 1, 1949 with potulish rug. 27- , but up to Makey box.

April 9, 1958 - about a dozen seen on a bond 1 mi. W of Plymouth, Amador CoCalif.




ť

- Feb. 26, 1949 Toledo, Lincoln Co, Oregon (DBM) One bright colored male in with flock of about 50 baldpates west of Toledo. Brick red head slowed up well. Was along marshy stream bottom with flooded fields and green grass bank. This bird, which McAllister spotted, was under gur observation for about 15 min.
 - Nov. 22, 1949 Gray Lodge Refuge, Butte Co. Calif. new-life-lister! Saw three of them today, in with the many flocks of baldpates on the open waters of the refuge. The first I have seen in the open. The orange-tinged crown and brownish wash to the head were distinctive. There may have been others on the refuge, but it would be purely accidental if they were seen. These 3 I saw were all within a 50-foot radius of each other, among the balapates.

10/71/14 - E-Comme



1757 -1762 a - 1758-1766-4772-1779-1795-1794-1797-1797-1799-1804-1819-9827-1822-1821-1828-1830 990-2002-2004-2005-2006-2007-2014-2028-2028-2029-2033-2035-2040-2049-23472-2387-2458-2542-1.830-1831-1832-1834-1551.1862.1862.1366 1 374. 873-1879-Obseiver Fred Evenuen, Jr. 75-17-81-102-111-191-192-193-199-206-210-212-218-244-252-260-261-American Pintail Female, immature: Craw: Volume of contents, 26 c.c. 8 kernels of Barley, Hordeum vulgare The rest was wild oats, Avena fatua Gizz ard: 10 c.c. volume of contents black Bindweed (Polygonum Convolvulus) 23 seeds Wild Oats, Avena fatua 2.5 c.c. Barley, Hordeum vulgare 9 kernels Grit and Undetermined 6.5 c.c. Summer Lake, Lake Co., Oregon: November 1, 1942 Immature Male: dizzare as large at malkara and volume 11.5 c.c. Cyperus sp 4.5 C.C. Ecirpus sp i.i c.c. Craver and grit 4.0 c.c. Broken up plant material 3. U. C. ... Adraecen 11-mile .m.Odl 100 ree reg Lrch 6, 1842 The Libum area of the pair coeff reads and Leader and provident to apply The what-plane to the inch of the his months are not the is great cirtane a own the aramo. ang 21,1943 11a.M. PWT - Weroma Beach Oregon 8 Pintails going south over the surf - flying iow, it any 16, 1946 - Stewart L. O. storm followed that afternoon 12-mile Swamp or pretty close to 1000 of them, equally M & F, in the grass Jan. 29, 1947 and shallow water at north end of the swamp, flushed, circled, and landed again farther on. Feb. 4. 1947 12-mile Swamp 2 2 flights of 70 and 30 went over us at the south end, and then we flushed about 200 more at the north end of the swall, and these left the swamp area altogether. Apr. 15, 1947 12-mile Swamp or perhaps 100-150 birds, pretty well-divided as to sex. July 24, 1847 Form Fiage lam: 0 5 in golige - plumage. Moodburn Golf Course: Ost a.m. a might or he care in at her! in me we not th, July 30, 1947 and they veered as they ame over us, one went on to the state state . Limit seggregate by sex. Thunderstorm in the area, see a missay rain allows. August 8, 1847 Fr rn liage Lam: again I rev reversi or there bitter. Jome makes her in fur. plumage, tair and arl. March 30, 1948 12-mile Swamp: There were about 125 or so birus, and all scened to be about paired up. Some pairs were even resting side by sie with their bills tucked underwing. July 21, 1948 (inlamette 4., belo preven City / / alli inter in the world a very court Sept. 8, 1948 Sacramento Refuge: 10-15,000 of them ! A sight to see them as they fulush! Sept. 25, 1948 Grey Lodge Refuge: 1000's present, and flushed this a.m. Few males in full plumage. Must have been 5000 in the area. Oct. 23, 1948 Grey Lodge Refuge: 35-40,000 seen. Jan. 24, 1949 Tulare Lake Basin at least 10,000 in flooded stubble field.



Feb. 1, 1949 Long Tom E, 3 mi. N of onroe, renton Co. (TH A) - the most abundant cabbles in this unusual concentration. 3 pintails to 2 mallaras. Retween the bridge at this point, for 3 miles up the river (south) to konne, the river was a mass of cucks. This concentration began just south of the bridge and consisted entirely of cabblers - mallaras, pintails, and a few scattered, rech-winged teal. At this point there the river bends just south of the bridge, I counted 2,000 birds by comparative groups of 50. 10,000 wirds would be a conservative estimate for the entire concentration, as I only covered a small portion of the river south of the bridge.

The bridge marks a definite point of change in the river. South of the bridge, where the dabblers concentrated, the river suddenly becomes shallow and narrows with many mud banks and bars extending into the river. North of the bridge it was deep and wide with abrupt banks. This point was come out interestingly enuf by the duck distribution, for all the diving ducks, mergansers, and wood ducks we e found north of the bridge. The pied-billed grebe, white-fronted goose, and a few dabblers were also found on the north side.

July 31, 1949 Grass Lake, Calif - all in Ferale Lumane.

Sept. 1, 1949. Tule name, Calif. 1... however in the area. Forther on between patient. Probability 75%. There is result a but in the first area.

Sept. 24,1849 Vallejo obtuat, Epal. to some cr., common. opin: releven a 56 sexed birds, in order it as obtained, in a some issues, in a some income and a some income income income income i

October 5, 1949 South of Yachats, Lincoln Co. Oregon. 8 of them on pools of rocky coast in tidewaters. On pools of salt water on sand beaches, and rocky headlands, too!

June 9, 1950 Putah Sinks, Yolo Bypass, olo Co. To p.m. about 25 seen in one bunch, and 9 more in anot er. 1 F with 7 young

bunch, and 9 more in anot er. 1 F with 7 young
Oct. 16, 1952 Southside Park, Sacto. Calif. 1 Amale. Plumage just turning for last time
it was almost all F plumage. Back graying at base of neck and black and
white under rump turning.

Apr. 9, 1958 - Pair on pond near Plymouth, Amador Co. Calif.

Oct. 10, 1959, Bay Færm Island and Oakport Road, Alameda Co. Calif. 150-200 ointails, seen, but only a few full-plumaged males among them.

788, 1966 - 100's over lie hade Ruding South to 3.5' + 201.



mas falenta

Feb. 6, 1954 - Stow Lake, Golden Gate Park, San Francsco, Callif. 5 p.m. The bird presumed to be a Falcated Teal was feeding on the mrass adjacent to Stow Lake with about 35 or 50 Baldrates. It is assured to be a male. When walked towards, it moved away the factest, entered the water first, and swam the farthest away, but came back ashere with bildrates as soon as we reentered the car. Fen flushed rapidly, it flow perfectly well to the middle of the lake area, along with a conformation. Sheculum not noted. About baldbate-size, with generally gravish a coarmac of back and wings. Rear flank, and under tail somewhat like a green-winged Teal male, with light rich buffy tones, bordered by black. Chest distinctly speckled with unique pattern. Brownish-gray on lighter hase. The head, usually pulled in with neck extended only when feeding. Dides of head an irridescent green, including the very long plumes at the back, which usually rested upon the back of the bird. Gave head an odd shape. Lever saw the crest erected. In addition to the green on sides, a plain white spot at top base of bill, with ruddy forehead and top of head. Throat and upper chest, and part of neck sides white, with heavy black ban i across white.

| | | | | , |
|---|---|--|---|---|
| | | | | |
| r | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| , | | | | |
| | | | | |
| | | | • | |
| | | | | |
| | | | | |
| | • | | | |

112/821-1836-1839-1840-1847-1853-1888-1982-2002-2007-2024-2025-2035-2049-2106-2110-2132-2516-2018-2606-

Observer Fred Evenden, Jr.

cumber of liels this.

252-257-264-165-266-268-269-281-298-307-321-328-337-

boats in the colonial three colonials

Oct. 15, 1942 Corvailis, Oreg. McFaaden Pond 7:30 to 8:30 a.m.

Male Immature:

Gizzard nearly empty, 0.7 c.c.

Black Binaweea Polygonum Convolvulus 21 seeds and many more were broken up

0.3 c.c.

Bur-reed, Sparganium (eurycarpum)
Spikerush, Eleocharis sp.
Grit (quartz and brown-rea agate)

0.1 c.c.
0.2 c.c.

Female Immature:

Gizzard content volume came to 1 c.c.

Amimal matter 0.3 c.c.

Hemiptera (Notonectique or Corixidae) Coleoptera (Dytiscique)
Homoptera (Leaf Hoppers)

Plant Natter 0.4 c.c. (Highly broken up) Black Bindweed, Polygonum Convolvulus Spikerush, Eleocharis sp.?

Gravel and silica or quartz . 0.3 c.c.

Female Immature:

Gizzard total volume 1.75 c.c.

Black Bindweed, Polygonum Co: volvulus Spikerush, Eleocharis sp.? White pulpy plant tissue, undeterminable Grit 0.3 c.c. a few fine barbs of white feathers.

Homoptera (Leaf Hoppers)

Gizzard nearly empty | 1 c.c. volume total Male: Black Bindweed Polygonum Convolvulus

Spikerush, Eleocharis sp.?

Bulrush, Scirpus sp.?

Nov. 1, 1942

Summer Lake, Lake Co., Oregon

Male: Gizzaro hao 3.5 c.c.

0.5 c.c. Potamogeton sp. 0.5 c.c. Ecurpus sp Carex sp. U.2 c.c. Cyperus s . 0.5 c.c. 1.0 c.c. Fine Grit

Plant laterial undetermined 0.8 c.c.

Female: Gizzard nearly empty -- 1.5 c.c.

Scirpus nevagensis

(Nevada Club-rush) 0.66 c.c.

Other Carex seed and

unaetermined plant matter -- .4 c.c. Grit .3 c.c.



Green-winged Teal
page 2
Nettich carolinense

| D ate | Location and observations | | | | | | | |
|-------------------------------|---|--|--|--|--|--|--|--|
| Nov. 1, 1942 | Summer Lake, Lake Co., Oregon shot in the a.m. Female: Gizzard contents l.c.c. Scirpus cernuus Vahl Carex alerta Contentia seage Scirpus sp | | | | | | | |
| Jan. 17, 1943 | la-swamp south of Corvainis: Orling to talk about, yet sin! Boy !This was really substained to talk about, yet sin! I would say we say well over 1000 or these toul and the almost solid flock cirching and of life and squarking as they flow overhead. Then they were now the usually silent air sounded like a miniature gale. | | | | | | | |
| Oct. 5, 1943 | South Tacoma Swamp, Tacoma Wash. Watched 5 or 6 of these teal feeding on duckweed colonies. They would put their bill horizontal and open and close it as they moved forward into the patch of cuckweed. True dabbling. | | | | | | | |
| Feb. 4, 1947 | 12-mile Swamp, between 200 and 300 were seen in the grass and shallow water at the north end of the swamp. There has been a decided reduction in the number of this species the past couple of weeks. | | | | | | | |
| Dec. 7, 1947 → Feb. 8, 1948 | Item liege lead to the to hant, but to the first of the state of the swamp today, and what a thrilling sight it was to watch them wheel and turn in unison. One flock picked up some male Shovellers, and after a few banks, the Shovellers ended up at the head of the line, and they retained that position for some time! "t was indeed a wonderful thing to watch, and it is too bad that it all couldn't beeseen at the same moment. | | | | | | | |
| Feb. 14, 1848 | 12-mile Swamp probably 4000 or more all together. Luring the morning a markh hawk female kept them in the air for some time, as it kept making passes at them. The passes were lazy-like and the hawk appeared to be playing with the teal. | | | | | | | |
| Sept. 1, 1949 | fule i. Calin. 1884. most es on patient in sos it a nest to salvares | | | | | | | |
| Oct. 16, 1952 Nov. 7, 1952 | Southside Fark, Sacto. Calif. 1 male with full-colored speculum, but still in F plumage. SouthsidePark male getting full plumage now. | | | | | | | |
| | | | | | | | | |

querquecula discorp

iate | Location and observations

March 5, Ave. nest of Corventies and a little to a make in act, allowed a floc of 16 Ameswing Seal From a make in act, vicinity. They estimate a titues and file over a march wis.

Makes had their olding on a sure thit or cents.

Jan. 14, 1947

12-mile Swamp. 2 F and 1 M. this small group of 3 flew by at 50 yards, and from all appearances, the one male could be nothing else. It is questionable, inasmuch as this is out of season for any migrants.

Jan. 29, 1947, 12-mile Swamp took off from the center of the swamp, and went into the south end of the swamp, as they did Jan. 14th.

May 13, 1947

12-mile Swamp? Tom found a female and two 5 or 6 day old young swimming in shallow water among Eleocharis at the south end of the swamp, in an area where we had not visited before. The "talked" to them as she flew, and they evidently "disappeared", for we did not find them later. Later Tom and Foc Gordon saw the male bird, and still later, we did not to see it when it took off from this same vicinity with S or 4 fallards. This is a doggone good record, and will be hard to make some of the Fird-boys believe

August 8, 1947 Fern Ridge Lam[©] "Blue linged Teal" app?? or either there two. 5 were observed resting and preening near the fixing road, and in the obscure plumage, although the wins-patches showed us word, no other calor differentiation could be made.

Aug. 13, 1947 | 12-mile (wamp? flushed a young, of non-descript plushed, and then 10 feet away, I flushed the S adults in eckipse plumage. The list and getting quite colorful, and one bird has the check patch beginning to develop so this verifies fact week's.

Sept. 2, 1949 la-mine avang@ wegon: e true sinc. north e. true since.

color amost vaite, indicating the readily care-ray considerate.

cinnaron teals.

. (4. (4. - x * ², +g+ -- 14, 1 , r T.

Observer Fred Evenden, Jr.

Number of field trip on which observed:

Location and Observations



On May 23, 1942, a group of students from the Field Zoology class at Oregon State College had an opportunity to observe a male cinnamon teal (Querquedula cyanoptera) in full breeding plumage, while on a field trip to a swamp some twelve miles south of Corvallis.

The bird was observed at a distance of 25 feet for several minutes. It was the first cinnamon teal most of us had ever seen, and it was certainly a sight to remember. It seemed rather tame, and when I approached, it flew just a short distance and settled down into the reeds again.

The cinnamon teal in the Willamette Valley seems to be a rare occurrence.



May 28, 1944

Caldwell, west and east to Boise, Idaho. Saw female on nest and male was close by.

Mar. 20, 1947 Colo. R. Indian Res., south of Parker, Ariz: over 50 M and F present. Males are strikingly colored.

12-mile Swamp, Corvallis: Or one male was seen for sure, and McCarlister April 8, 1947 believes he may have seen another one as well. We all had satisfactory chances to study it.

12-mile Swamp, Corvallis: again we saw a beautiful male, perhaps the Apr. 22, 1947 same one, and it seemed to be along, for when flying, or feeding, it was always by itself.

Sacramento Refuge: perhaps a total of 750 or so seen over the area. Sept. 8, 1948 Few full males, many molting. About equal M & F.

Grey Lodge Refuge: still many present. Must have seen over 750 today. Sept. 25, 1948 Many males in full plumage.

July 31, 1949 Grass Lake, Calif. - all in ferale plumage. Menaota Pool, Calif. En in Temant permage. Aug. 2, 1949

Tule L. Califfords. rote pulse as a non-non-standar mere- and a Sept. 1, 1949

Gray Lodge: some males in full plumage. Dec. 29, 1950

SWMA, Falton, Churchill Co. Nevada: females with 4 and 6 young.
Mandota Pool, Frence Co, Cal. 15 - 911 + 5F on swall pend. Mandota Pool, 197 June 12, 1951 March 11, 1952 both do + ga.

Gray Lodge Refuge, Butte Co. Calif. many pairs, w/ extra males in May 1, 1952 full plumage, common.

June 4, 1952 SWMA, Lead Lake-Milan Units, Churchill Co. Nevada: common, many adult males alone on "pastures" - young broods too. Males perched on old muddied muskrat feeding platforms in old Typha & Scirpus areas at edge of HOH

it. Joe, lacific Fraye, Calif. Watched at least 7 ring in courts in activities.

Brusille Rd, Nof Elk grow Rd. 8 m + 2 For imm on slough new cattails.



hay 14, 1.41 in-mile evant scott of Carvani, of Cienton and one line to the tand and this is the execution Fine: in view and induce and which on the course of the course

iny 0, 11.7 Lo-hills for protein strong of the 12 of the

Oct. 15, 1942 Corvallis, Oregon. McFadden Pond (same as above) 7:30 to 8:30 a.m.

Female: Gizzard was very small 2 c.c. volume of contents

Black Bindweed, Polygonum Convolvulus

Bur-reed, Sparganium (eurycarpum)

Grit is quartz and reddish brown agate and basalt (?)

Nov. 19, 1942 Corvallis, McFaquen Duck Trap / Pond 2:30 to 3:50 p.m.

A bout 20 of them, many soult plumage males in the group.

Red legs showed up well on Tem les. They really "set-in" to a lake.

Oct. 5, 1943 South Eacoma Swamp, Tacoma, Washington
One female seen, and then a little later I watched a male in full
plumage preening itself on the water. While I watched it stood up
on its legs, stretching and beating its wings. It flew soon after that.

Oct. 27, '43. Washougal, Washington

Two males flew low overhead on the Columbia River bottoms.

Apr. 26, 1946 //ry's Miver/ Saw several males and Female s. Seemed to be paired anyl6,1446 - Standig Paile Swamp.

Jan. 14, 1947 - 12-mile Swamp, about 30 M and 15F (or juveniles). The males had as brilliant and beautiful a plumage as I have ever seen on them.

Feb. 25, 1947 - 12-mile Swamp: about equally male and females; perhaps 250-300 of them.

Apr. 8, 1947 - 12-mile Swamp 0 2 2-300 were present, the ratio being about 3M to 1F.

March 30, 1948 12-mile Swamp. There was a precominance of males and let year males, but there was a limited amt. of pairing up. Perhaps 50 pairs, and he single birds seen. All these ducks were scattered over the starp area.

Jan. 20, 1949 Signal Depet, Jawanento, Calif. - 430 p.m. - Pair on guddle infield across road from Depot. Itill present thru 7 et. 16th, 1949

Sept. 1, 1948 Tule Lake, willif. 1... about a come into the has analyseed and result part of this period. It to amberous us. 17-18. The brack to other bird on the come and filt one.

Mar. 28, 1950 Gust south of Oceanside, San Diego Co. Calif. observed several pairs of shovellers of which the males of two pairs were the courtship antics of neck and head bobbing. Males in full plumage.

Dec. 29, 1950 Gray Lodge Refuge: Quite a few males in full plumage.

2-20-52 Marshes NW of \$6/2406 Valleyo, Solano Co.Calif. most abundant duck, and most were males, saw two males "neck-bobbing" in front of one female.

Oct. 10,1959 Oakport Raod, Alameda Cr. Calif. several M and F, but no males in full oluma ge yet.

10/2/ 14 Obigantine NWA N' GERRY - Partial a Chino - Ted rettet , growie



Oct. 17, 1942 Covrallis, Ore. Mary's River 2 miles south. 3.pm

> Male, mature: Gizzard has 7 c.c. total volume of grit and highly broken up plant material. It was partially empty.

> > At the junction of the bronchi into the trachea was a holow, stilf, transparent growth about 3/4" in aiameter that has the consistency of a pingpong ball. At least that is what it reminded me of.

Feb. 19, 1943 Corvallia, Brook Lane of them on a slough of the Mary's River.

3-23-43 Horisina The, Stout, "re.

I ger. I one very large flocker of is.

Walla Walla, Washington 3-5p. m. & slightly somaller toran a Coldgrate. Head clark brownick gray, with white use patch, pointed to the rear. Wing tipe white, & white bar near back of folked wing. Head crested, prointed downward. Short, steep-sloped ball. High forched. Broad tall, rint. birest frown. Side receive was 4 wing white. Nov. 20,1943

Mar. 15, 1944 Pudding R., Woodburn, Oregon concentrations of 20-30 were found ina pond of backwater in the deep woods along the river.

Stewart Lake, Convenies. Flow to pair of them. One pair has a poung July 1, 1946

that can't fly yet, but they get up and run weis on the obser. "Poison Oak" Take, linn Coll Pair with raming or Il young. can't fly. July 4, 1946

- Hazelaur lake, Falen Od adults and young here is in this July 13, 1946 - "Poison Can" L., Linn Co Or Tutched one o bb. in in the color July 18, 1946 ater as rechere I sat. It would leap up one pull of a sevent cu

Soft Mana-grass heads.

later) and that was all I saw of them.

April 24, 1947 "Poison Oak Lake" Linn Co. watched a pair. female swimling and leading the M and giving continuous notes; while M preened, F flew up into an Ash by the water's edge, and about 10 feet high. Stayed a while, aropied back to the water and proceeded on around the lake, with the M following, both just looking, not feeding. All the while he was pushing his head way forward and pulling it way back, as if doing some phase of a courting. They'a whistle to each other every so often. They worked by way around one side of the lake, and another pair swam out and headed directly for them from the other side. As the 2nd pair approached the 1st pair retreated slowly, and in 5 minutes had been retreated some 50-60 feet along the west side. They whistled etc at each other, and after a time, the 1st pair advanced again, and this time the 2nd pair aid not chase them, but retreated, instead, finally flying to the east side where they came from, and the first pair advanced on another 70-100 feet Later this 1st pair flew up and around and lit in two oaks, the M 25-30 feet up on the NE side of one that had a cavity in it. He hung on the outside, looking in, and then the F flew over from the other tree, and displacing the M. she hung on the side of the oak, by the hole, and he lit on another limb,

> 15 feet away, bobbing his head ahead and back. After about 3 minutes, she dropped to a perch a few feet from the hole, and just sat there, and the M flew back to the oak she had 1st lit in. I scared her about then (15 minutes)

.

May 20, 1947 12-mile Swamp: flushed one pair, and they had nearly hegrown young that coulon't fly yet. Length was about 10" or so. La er, I chught one of equal size, after a hard chare. The bird would run on the water, and if I got very close it would give, and swim underwater for distances up to 15 feet without coming to the surface. The water was shallow and clear and I could see it using its nearly featherless wings as an aid in swimming, and it could see well under water, for it doaged vegetation growing wheat of it. Finally, it neared exhaustich, and I almost caught it, but it dove, and swam on, and they stayed down, and I tried to pick it up, but 100nd that it was holding itself down by using the birl, and in order to pick it out of the water I had to break off the seare it was nolding in its This bird was also about 10" long, and had well-developed claws on the toes, and the toe (hind) was partially lobed. Brownish caring in on the breast and shoulders, and the tail already had that "squarish" iorm, with the metallic coloration.

May 24, 1947 Peoria Ferry Load, March 5 miles for . Corvallis: Fat feature and 5 young swimming from Ponu Lily emergent-leaf area into spirea and

willow thicket in the sater. Fern Riage 1 am: many were seen at various location, at the tracery ir. August 8, 1947


lk-mile Swamp flushed / or 4, and saw one that was in the minuted stage. Augl 13, 1947 iem lie de more de la companie de la ·1ec. 7, _967 June 18, 1948

Thushed a family that had from help young, 1/2 croth - run val, I t

on the vaca. If -mise than problems is a least problems. July 8, 1948

Beaver Fond, 3 mi. of Pruce Ita., Penton Coleton mater and two readles Apr. 30, 1945 together playing about in the on pond about 5 a.m. They librate any attention to me. (IFM).

June 9, 1957 Natomas Dist, along Elverta Road, Sacto. Co. Calif. 2 pair, with 1 F in rice fields in open country, without woody cover nearby. Lay 8000, Fred 6. And. for in gracust in ing open paid type of the Interior Cte. and I get then the war named



•

- Canead 1026-1027-1033-1036-1043-1046-1047-1062-1234-1243-1249-1303-1313-325-331-374-378-367-957-974-975-984-988-7011

1383-1385-1397-1472-1445-1457-1560-14950-15524-15524-1562-1234-1243-1249-1303-1313-325-1336-1341-348-7349-1365-1329-1385-1339-1385-1349-1393-1345-1349-1393-1345-1349-1365-1348-

Late

Location and oppervations

Sept. 25, 1948 Grey Lodge Refuge: 4 immatures wwimming close to coots in open pond, at edge close to Sagittaria.

- June 30, 1949 [tillwater 'ilcline Parmit. mus, Ostrchil Collevada II. 400 musc of mostly males. Not as far slors in their moult as the classes. In communication of signs of molting. 17 breads. Treads were as solutions: 6, 8, 15, 17, 4, 5, 10, 5, 10, 5, 11, 9, 14, 11, 4, 5, 6, 7, and 5. The incommunity energy was tone of the broads of 4 is which 8 whe partry reathered. The desire it is the 4 must have been subpted of the 3 were adopted. Them of the broads of 7.76 young. This say 8 pairs, separate, alle and Female together.
- the respectant although the respect to the respectant of the respectant to the respect to the re
- Feb. 27, 1950. Doerfler's Pond, 2; miles SE of Colorado Lake, Linn Co. Ore. IBE. 3 p.m. First redheads I have ever seen in the valley. Jay Long says he has never seen them in the valley before either. Jay spotted this first one. Later we spotted others and looked at them through Jay's 19.5% spotting scope. They were in with ring-neds and cans. Still later three of us, in walling around the pond had a flock of redheads and cans alight right in front of us at about 100 feet in the bright sumlight. Saw up to 24 redheads at one time. Perhals this was the total for the pond.
 - June 12, 1951 SWMA, Fallon Churchill Co. Nevada. Fs with 8 2/week-old yng. & 6 yng.



Date Location a de Fivations

gril S., 1945 | Colv. Lide: "Tempere Le Sie Sie Sie Le Sie to the size is noticed to the second included the off male in the con-

Portland, Laurelhurst Saw one male on the pond. He had his Oct. 10, 1942 head tucked under a wing, and didn't want to be disturbed

Oct. 15, 1942 Corvallis, Ore. McFadden Ponu 7:30 to 8:50 a.m. Female immature:

Gizzard volume 6.5 c.c.

Insects: Black Ants (Formicicae) The mandible of another insect.

Digestea or pulverized plant bulk 2 c.c. Cruciferae, pod or some species Black Binaweea, Polygonum Convolvulus 2.5 c.c. Bur-reed, Sparganium (eurycarpum) 1.5 c.c. 0.5 c.c.

November 7, 1942 Wagner Butte, 10 miles suth or Corvallis Shot at 8:8 a.m. Immature Male: Gizzara Contents 4.0 c.c.

Crit 1.0 c.c. Unaetermined , lant material 0.5 c.c. Insect body and ing parts .E c.c. Unidentified minute arthropod ### eggs (?)Purple, black, 2.5 c.c.

said brown and tan or gray.

March 9, 1943 Corvallis Duck Frage Work Ord. There were about 50 of them on the good & of all the ducked prevent, they were translat & payed atte attention is me. I cause in exight used strong direct wing beater

Jan. 14, 1947, 12-mile Swamp - I thought I saw a flight of 4 or 5 land in a part of the swamp on the far side, but nothing further was seen of them.

Nov. 25, 1948 East Moreland Pond, Portland, Oregon: about 12 on the east pand, mostly males. No scaups!

March 11, 1950, Southside Park, Sacramento, Calif. 28M and 16F seen, many under shad of overhanging island vegetation, in . .ter, with several mallarus.

Feb. 27, 1950. Doerfler's Pona, 22 mi. SE of Colorado Lake, Line Co. Jregon .DB. .. about 100 of them. None of them appeared to be scaups although Charlie Traub thought he saw 2 scaups. They also performed beautifully by taking off and landing again often on and whist ing in for landings, too.

Nov. 7, 1952 | Southside Park, Sacto. Calif. M's and F's as follows: 1&1, 1F, 11&7, 3M - 24tota

Feb 24 1963 : Ed Convollie, in him & Oregon - 15 on a pand w/ coole Nov. 21, 1965, accoracteur 15. Vas's D. C. 2111 and 70 or inc.

Dec. 23, 1965 " ". 2511 and For inc.

3/k/30 C+O Cand, The Fully (48 15M 1 Fine



12/21/2 / 10/2 / 371 1 2 2 1 12/27/-19

| t | | | |
|---|--|---|--|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | • | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

Loc vion and observations

November 1, 142 Summer Lake, Lake Co. oregon | Inot in a.c.

Wale: Gizzald contents only partly rime - 3.5 c.c.

Grit otamogeton sp 2.75 c.c. 0.4 c.c.

Ecirpus sp.

Feb. 21,1943 Warrant - in a contract of the there is way of the to the second

Jan. 25, 1947, Echman Slough, Waldport, two bunches of 6M and 4F, and 4M and 2F. After taking off, legs hang awkwardly for some time

Nov. 25, 1948 Madrone Reservoir, Butte Co. Calif. 1975' elev. 230 p.m. adults, M & F and immatures - 25-30 present.

Bass Lake, El Lorado Co. about 90 resting with bills tucked underwing this noon, but as wind came up from the SE, they began to spread out, and feed, March 5, 1949 showing activity.

Feb. 24, 1949 Long Tom R., 3 mi. N or Monroe, Oregon - 60, some frequently gave carls that could be described only as shrill growl or like the banking of a ting sug. Some of the males giving the near-throw. W outnumbered F 2 to 1.

July 31, 1949 Grand Land, Camille 1 in Formula plumage. Aug. 2, 1949 Me naota Pool, C if. o in it. Local lungo.

,

•

Mr. David P. Olson, Research Asst. Univ. of Minnesota, St. Paul 1, Minnesota.

Dear Mr. Olson:

I hope some of the following data will be of use to you in your study of the Canvasback:

Breakdown of localities and months for data used in j of wlm article in 172:

October - Tomales Bay area - 1:5

November - Sacramento - 1:2

December - Sacto. - 0:2

January - Sierra foothills E of Sacto - 1:3 Sacramento - 123:98

February - Clear L, Take Co. Calif - 2:6 Sierra fthls E of Sacto - 25:32

Sacto - 40:27

March - Sierra Fthls E of Sacto - 1:3

Sacto - 3:0

April - Sierra fhtls E ofSacto - 1:1

Western Oregon

Lincoln County - Dec. - 2:0 Jan. - 10:5 Feb. - 2:28 Var. - 2:2 Apr. - 3:2 (before 1953 records)

Benton County: Dec. 1:0 Jan. -2:0 Feb. 52:28 Mar. -2:2 Apr. 3:2.

Churchill Co. Nevada - Apr. -2:2.

March 17, 1962 - California: Bolinas Lagoon, Marin Co. - 6:5F or imm. Salano Co. marshes - 20:32F or imm.

In addition: Counts without sex ratios (there before your request care in

Mexico: 3/6/51 - 1 at Terminal, NE Zacatecas, Mexico.

3/9/51 - 1 at roadside pond, Jimenez to Chihuahua, Chi. Mex.

11/6/61 - SE of Tos Mochis, Sinaloa, Mex.-4

11/12/61- Guadalajara to Lake Chapala, Jalisco, Mex - L

11/14/61 - N of Tagos, Jalisco, -Mex. 2

Sincerely,

.

 $(\mu_{ij})_{ij}$, $(\mu_{ij})_{ij}$, $(\mu_{ij})_{ij}$, $(\mu_{ij})_{ij}$, $(\mu_{ij})_{ij}$, $(\mu_{ij})_{ij}$, $(\mu_{ij})_{ij}$

the state of the s

16 21 97 8 21 77 8 7. 7.7. 1. 1. C. 615. 1:5 l/ C. . T 20, 100 Citarian 11 to the second of the second 50,00 7,7188 the state of the s 13:71 714 7 Sind while Barrier Commence 11.5 11.2 11.0 1200 1200 1200 121 many of the 1 - 1 ---MEXICOSE OF LOS MANCHES, SHELLES IN GUPPHARRIE L. C. M. J. J. C. C. C. C. NOF 201685, JAN 400, 2 ON 1/19 01 JIMENEZ TO CHIMBERT, 3/9/51 TERMINAL, ME THER CONS - I CA CONDER, WILLES

• • •

| 1961 | | | BALANCE OF 1961 - 1962 SESSION SCHEDULE | | | | | | | | | | | 1962 | 1962 | | |
|-------------------|------------|--------------|---|-------|-------------|-------------|-------------|-------------|--------------|-------------|------|----|-----------------|------------|------------|--------|--|
| ELDER | 9/20 SM | 9 / 24 CT | ' | 1 1 | 10/18 SM | 10/29 SG | 10/29 CT | 11/15 SM | 11 /26 CT | 12/17 CT | 1 | | 1/7 CMM 2 | 1/15 CG | 1/17 SM | | |
| веам | X | Х | X | | Х | Х | | X | | X | X | | X | X | X | 1 | |
| BURKHARD | X | X | X | | Х | X | | X | | X | X | ļ | X | X | X | | |
| CASTLE | X | X | X | | X | X | | X | | X | X | | | X | X | | |
| CRITES | X | X | X | | X | X | | X | | X | X | | X | X | X | | |
| CUNNINGHAM | X | X | | Х | Х | X | | X | / | X | Х | X | | Х | X | | |
| DAVIS | X | X | Х | | Х | X | | X | | X | X | | X | х | X | | |
| DUNN | X | X | X | | Х | X | | х | | X | X | | х | X | X | | |
| EAST | X | X | X | | Х | X | | X | | X | X | | X | X | X | | |
| EVENDEN | X | | | | Х | X | X | X | | X | X | | Х | Х | X | | |
| FREEMAN | X | X | X | | X | Х | | X | | X | Х | | Х | Х | X | | |
| GUETLING | X | X | X | | X | X | | X | | | Х | | X | Х | X | | |
| KENDRICK | X | | X | | X | X | X | Х | | | Х | | Х | X | X | | |
| KIPP | × | | Х | | Х | Х | X | Х | | | Х | | х | X | X | | |
| KROGH | X | | | | X | X | X | X | | | X | Х | | Х | X | | |
| LARSEN | X | | + | X | X | X | X | Х | | | Х | X | | Х | Х | | |
| LAUGENOUR | X | | | X | Х | X | X | Х | | | Х | | | Х | X | | |
| | X | | | | X | X | X | Х | | | Х | Х | | X | х | | |
| MC CURRY | X | + | | X | X | X | X | X | | | X | | | X | X | | |
| | | | - | | | | | | | | | | | | | | |
| MEMERING | X | | X | | X | X | X | X | | | X | X | | X | X | , have | |
| OGDEN | X | | | X | X | X | X | X | | | X | - | | X | X | | |
| ORANGE | | | + | | | | | | X | | X | X | | X | X | | |
| PAPE | X | | | | X | X | | X | X | | X | X | | X | X | | |
| REED | X | ' | | X | X | X | | X | X | | X | - | X | X | X | | |
| ROHWER | X | | - | | X | X | | X | Х | | X | X_ | - | X | X | +; | |
| SCHUNKE | X | - | | X | X | X | - | X | X | | X | - | | _X | X | | |
| SIEFKIN | X | | | X | X | X | | X | X | | X | X | 1 | X | X | - | |
| SPARKS | X | - | | X | X | X | | X | X | | X | X | 1 | _X | X | _ | |
| SPEARMAN | X | - | | X | X | X | | X | X | | X | X | | X | X | - | |
| STAINBROOK | X | | | X | X | X | | x | X | | X | 1 | 4 | X | X | | |
| YEATES | X | | | X | X | X | | X | X | | Х | X | | X | Х | | |
| TOTAL NEE DE D | 30 | 10 | 12 | 12 | 30 | 30 | 10 | 30 | 10 | 10 | 30 | 12 | 12 | 30 | 30 | 0 | |
| NEEDED | 30 | 10 | 1 6- | A See | 1 | 30 | | 30 | 10 | 10 | . 50 | 14 | 16 | 30 | 30 | | |

SM - REGULAR SESSION MEETING 7:30 P.M.

CT - SPECIAL SESSION MEETING TO

SG - SPECIAL CONGREGATION MEETING CG - ANNUAL CONGREGATIONAL MEETING

RECEIVE COMMUNICANTS 5:30 P.M.

CMM- COMMUNION SUNDAY

"1" - 9:30 SERVICE - "2" - 11 A.M. SERVICE

IF YOU CANNOT MEET THIS SCHEDULE OBLIGATION, PLEASE SECURE YOUR OWN NOTE: SUBSTITUTE WHERE SUBSTITUTE IS NEEDED TO MEET A QUORUM OR FILL OUT A FUNCTION OF SERVICE.

University of Minnesota

Institute of Agriculture
St. Paul 1. Minnesota

DEPARTMENT OF ENTOMOLOGY AND ECONOMIC ZOOLOGY

22 February 1962

Mr. Fred G. Evenden 1336 Fitch Way Sacramento 25, California

Dear Fred,

From the enclosed cooperator letter you will note that I'm studying canvasback sex ratios as one phase of my canvasback research program.

I'm very much interested in your sex ratio observations given in the Journal of Wildlife Management in 1952. Could you tell me something about the dates of the California observations, also the type of habitat involved? That is, were the canvasbacks seen on large lakes or bays such as San Francisco Bay or smaller lakes and marshes?

Your data and that given me by Morton Smith in Louisiana suggest that there is a tendency for females to winter further south than males. If you have a chance to observe any canvasbacks this winter and spring or know of anyone who could, I'd be grateful for the information.

Sincerely,

David P. Olson

DPO/jeo

Encl.



Observer lied livencen, Jr.

Greator : cata, .uc.

124-366-1107-1429-1433-1532-1559-1578-2242-

Tyroc. Marila

DATE

Location and our row time.



Sept. 5, 1942

Pudding River, Woodburn: This afternoon there was a flock of about 35 Lesser Scaups feeding in the river around and just below the outlet to the sewer from the Ray-Waling cannery. The river is filthy below this point, and someday I hope that someone will be appointed with the power to clean up and halt such damages to our rivers in the State.

Oct. 24, 1942

Pudding River, Woodburn Flushed a few of these from the river below the sewage outlet and found one dead one. Branches and logs along the river are coated with the filthyest slime you ever saw.

apr. 1, 1944

Inake R. Idaho - several were of served dead along the ranks of the . iver.

Feb. 11, 1948

Dec. 27.1954

near Owl Creek, E of Corvallis, 2 p.m. - 3 males and 2 females were on a small pond, beside a fence row within 15 feet of the Albany-Corv. hiway! When I stopped the car near them, they swam rather leisurely away.

nug. 31, 1949 Tule Lake, Calif. one journ one has botulien. I ask at the a season

5. San Francisco Par, Son Cateo Paidre, several landre to the groups and scottere to ter have onen visible.

*

.

.

Observer Fred G. Evenden, Jr.

| | And production of the Control of the |
|----------------|--|
| Date | Location and observations: |
| Dec. 3, 1943 | Wallula, Wash., along Columbia R I saw 10-12 individually and one flock |
| | of 8 flying together downstream. The black head, and back, and white |
| | sides show up at a distance, and from 50' to 100' distance the black |
| | head turns a dark green of the nearly round spot on the cheek becomes evident. |
| | Under parts and sides all white. White inner wings. White on sides when bird is on |
| | the set, with faint streaking of black on higher parts. |
| Apr. 1, 1944 | Glenn's Ferry to Hammett, Idaho. There were singles, pairs and large flocks |
| | I observed one female jump cirectly into the air when frightened suddenly. |
| | They are a very wary bird, and are hardto get near. |
| April 13, 1940 | the state of the s |
| | of Goots along the wort side. Light was wrong/ so we clun't make a certain identity, other than "Golden-eye" until we furshed it, and it liew north, |
| | to the east of us, when the greenish-glint to the head, that the remaish |
| | white spot were plainly seen. |
| Dec. 2, 1948 | |
| | full males, and another flock of 3, or which there was one female or immature. Also 3 other males seen. |
| | |
| Feb. 22, 1949 | 20 seen on Rogue River between Rogue River and Medford at 1330 noon. |
| Jan. 5, 1951 | |
| Feb. 6, 1950 | Bay Farm Island, Alameda Co. Calif. Few. |
|) | in the latest in place. |
| Mar. 17, 19 | 262 Bolinas Lagoon, Marin Go. Chlif. Moonish. 1 ad. mole per For mim. 25-30sec |
| 3/7/04 | West granting landy fitch, lid, 20-25gir, many courting, twill gracent mate |



Darrowin Colera-

Lucinatt i lamaic.

__te

August 5, 1946

Lin: Co. 4000' elev. This p.m. a group of us watched a female feeding in a shallow lake, near the scout camp lake. Le were quiet, and it carried on normally and gave us all a treat, as it went by us slowly, not over 30 fast away.

Mar 26, 1947

May 16, 1947

Tillamook Bay entrance by Tom McCallister 1 pair.

Lost Lake, S. Santiam Pass 2 pair were seen in the center of the lake, and they swam for a while, as we approached, and then a dived, and 2 ran for a take off

Sept. 11, 1947

Big Lake, Linn to - and victor instatures and to a refer to by or a change munage. I got within the toutton as tic Yearn a pure our set the in the or the rime. Constitute will see connect once, in it is not blosse times that I shedded to choose to them, because on the are type her hours not them add me move, socione, not alime, or alime notes are lively ut they clian't may, and as in all a up on them the jist san recapilled or me, and tiru the inact into one main part of the first of including that off (one not beam to be to be to rounded of privation), but it has low years trying it. The of pay not out the fator is 76! in . o. The, were in group train in in all highly popular, they may rare to [5 others in mic-lake - z_{ij} - z_{ij} - z_{ij} body stantold a took z_{ij} c_{ij} - 1415 of TLost Lake, I. Santiam Pass, Lim Co, Nogon - 2 pell M & I F or inn. Living in mid-lake

May 27, 1951

Apr. 9, 1967, NE of St. Maries, Benewah Co. Idaho. 8 M&F. I more melanistic F in breeding plumage?? Bill appeared more slender, and was orange-red not yellow. No whitish area below head. Whole body more or less uniformly brown. Body seemed slightly smaller than F. Golden-eyes.



Late Location and observations

Lec. 10, and a Representation of the state o

Dec. 3, 1943 Wallula, Wash. - along columbia R., - saw 5 of them today.

Apr. 7, 1944 Mtn. Home Reservoir, Idaho. Male and female. The male was brightantly white and the remale was quite drab, with a white spot on the side of the head. The male would dive and when he broke the surface the female would dive and when she came up they would both stay up a few moments, and then they would repeat the above procedure.

August 9, 1946 Red Butte, lake, Linn Co. 9 a.m. We had a swell view of a male and a female Bufflehead, and their 4 young swimming in the deeper water off the south end of the lake. The young evidently couldn't fly, for all they did was get up and run on the water. Reither of the adults flew or dove, either.

Feb. 11, 1948 Stewart L, la mi. N of Corvallis: 3 males in beautiful plumage, and 3 females 2 of which were last years. Flushed them 3 times, but they just flew to the other end each time! A Lesser Scaup male was with them. The males and older female took off with a noticeably shorter run (about 12), than did the two young females, which took almost 20-25! to take off.

)Feb. 28, 1948 Stewart Lake - 230 p.m. 3 males and 3 females present (one of which was the joung mentioned before. The 3 males were really beauties, and flushed back and forth on the lake several times. Take-off runs averaged about 15.

March 30, 1948 12-mile Swamp: 4 immatures, or females, probabily the latter. Take-off distance, 5-7 feet.

June 15, 1040

1 -mine (vample of). Is obtained in its translation of the series of (probably 5/4 size) in consequent the frame of incitive his inclines of the without pond. It was probably an impating, for the wines; control in model of the pot, jet a time of the series in the frame of the point of the families of the only divide with I had been and the first translation of the pot, etc. I finally flushed there bive, etc. escans from fit of interest the rection, and the influence to be first-cot, one. The protected the series of the first-cot, one. The protected the series to be first the expectation of the potential of the series to be a first three one the sound of the first thin, the company is the first test of the first three company in the left believed to be a first through this part of the first and from first place the translation of the first pond on the first than, the three himself the first pond on the I had been one got that the bives return the first pond on the I had been one got that the bives return to the first pond on the I had been one got the set by the series of the first pond on the I had been one got the . Surface of the first pond on the I had been one got the . Surface of the first pond on the I had been one got the . Surface of the first pond on the I had been one got the . Surface of the first pond on the I had been one got the . Surface of the first pond on the I had been one got the . Surface of the first pond on the first pond of the first pond of

Dec. 2, 1948: Truckee R., just below Lake Tahoe of I F or imm. on river with coots & redhead.

Feb 24 1960 Hairry Frebry, Nothing to Carly - 500 6 painted - 7 Juliots.

. Mar. 17, 1962 Kani Bolinas Lagoon, Varin Go. Calif. Moonish. 1 ad male to 3 For im -40 seen.



EDWARD GREY INSTITUTE OF FIELD ORNITHOLOGY

(Dept. of Zoological Field Studies, Oxford University)

BOTANIC GARDEN

OXFORD

Director: David Lack, M.A., SC.D., F.R.S.

Telephone: Oxford 47826

Dr. F.G. Evenden, 1717 Professional Drive, Sacramento 25, California, USA.

January 5, 1960.

Dear Dr. Evenden,

Thank you so much for your letter in reply to my request about Harlequin Ducks. You have a nice little group of records, naturally I would like as much detail as possible upon them, since this increases the value of each record. You did state what sex and ages were involved in the records of August 6,1942 and August 7,1942 but not in the cases of the records of April 12,1941, and 14,1942, march 19,1943 and June 15-14,1946. It would greatly a 13 to the value of these records if you had information to fill in how many birds of each sex were involved; I also wonder if the June 15-14,1946 record may not have included young birds. Another puzzling record is that of August 7,1942 since I know of no record previously in which a male was accompanying a female with young. It should have been molting on the sea.

If you could find time to add further details to these interesting reports of yours I would be very grateful. So as to save having to list all such desirable details I am enclosing a copy of the current questionnaire and wonder if you could complete the details in that perhaps.

Thank you so much for writing.

rours Sincerely

Timothy Myrel

December 14, 1059

M. T. Myres, Edward Grey Institute of Field Ornithology, Botanic Garden, Oxford, England.

Dear Dr. Myres:

I noted with interest your request for information on surmer records of the Harlequin Duck. This is in response to that request in the Condor.

April 12, 1941 - Mollala River, N. Fork, Clackamas County, Cregon.

April 27, 1941 - Facific Cean, Newport, Tincoln County, Gregon.

March 14, 1942 - Mary's River, Corvallis, Benton Co. Cregon.

August 6, 1942 - Big Lake, Hastern Tinn Co. Cregot, (top of Cascade Pange) Female

August 7, 1942 - Big Lake, again. Male, Female and 4 young on the lake.

March 19, 1943 - Mollala River, near Canby, Tlackamas Co. Cregon.

June 13-14, 1946 - Breitenbush River, eastern Marion Co. Cregon.

Sept. 7, 1946 - Boola Park and Also Short Sand Beach State Darks, Pacific Cean, Clatsop and Tillamock Counties, Oregon.

April 21, 1951 - Point Pinos, Monterey Co. Calif. Pacific Ocean.

I hope that some of these records may be of interest to you in your compilatory work on the Tarlequin Duck.

Sincerely,

Dr. Fred G. Evenden,

FGE:me



Funder of field origin on lies of May 1.57-62-20-102 -123- -220-366-403-446-489-566-567-1004-1904-

estern Harle uin Luca

histronicus histrionicus pacificus

Lite | Location and observations

Aug. 6, 1942 Big Lake: Female swimming non-chalantly in a small road of the north side of the lake. Larger & heavier heathered around the head. Henevally a dark brown color. Notary larger than the Wallards.

ang. 7, 1942 Big Lake: Watched the male & female & 4 young in the lake near the grown which I watched watched (I made this change in identification because of proints of Cearned in identifying I ducks in Jame Manage of Cease.)


June 13,1946 Breitenbush R., Forest Camp. 7 p.m. I caught a glimpse of a female flying upstream. Sometime later one, probably the same one, flew upstream and set its wings opposite me, barely lit, and dove below the surface. It soon came up, and shook water a little. Then it dove under and didn't come up till it had swum a out 30' across the current. It is almost unbelievable, for the water was very swift and strong in its course, and the stream is strewn with huge boulders. A little later the bird appeared on a rock 50 feet below me, and proceeded to preen itself for a few minutes. The male was not seen.

6/12/67 Clover PT, Victoria B.C.

525 valio Py/ For imn: 1/0 2/0 4/2 8/3 5/, 4/1 5/0

6/13/67 " Relias - 3/0 5/3 2/, = 16/4 or 4/1

8/26-27/74 Beauty Kell-Clover JT, Whiching the 14M + 17 min 2 200 4/1



Fred G. Evenden, Jr., Observer

Old Squaw

Number of field trips on which observed: 192-566-1001-1381-2312-

Clangula hymealis

Date

Location and observations

Mar 26, 1947

Dec. 6, 1949

May 16,1955

Neahkahnie Mtn, Tillamook Co by Tom McCallister: one bira taken to be of this species in with a northward moving flock of Surf Scoters.

Hoss Landing, Lonterey Co. Calif. 1.0 . 1. 0 served just one female in winter plumage swimming in the lagoon with other tucks. Tope Beach, Take Takes, I. of Upper Proches ., 1911. 31.0. 1 is threatening weather, but end of seve I second of seving storms. I could be rule and invenile male riding low to an en - or cryphy granis - line to common. I could be single and back. The cides and being and back. The cides and being a line of the short and a for join. On so at 75' w/ 74 binos in sumstine. The short are also for join. On so at 75' w/ 74 binos in sumstine. The white from an island, what also, many brown on one, and brown belong with the from an island, many also, and brown belong the land and a could be common be seen in the same make the short of an and base of held cuto needs behind. In equal so only a land with the could be seen in the same and the streets.

3/24/79

Exposition fash, Toronto and 15:10. 6 por, uf some "world lighting"-courting?

•

.

Tumber of field trips of lie of this of the section of the section

Location and observ tions

Apral (to, assis to board is the formula continue to the part of the part of the state of the st The both is should be enjoy as are;

nimin () solveil i, for an analysis of the solveil war war and a live and a

 $\frac{1}{2} = \frac{1}{2} = \frac{1}$ Jan. 25, 1947, Yaquina Bay, south shore. about 8 of them out on the sand, some 15 feet from the edge of the water. Perched on wood debris, or just on the sand. Just seemed to be resting. Didn't pay much attention to us. When we flushed them, they got up and "ran" on the sand, as they would on the water, and took off in a distance of about 15 feet (helped b, the 25-30 mile winds today). All took off quartering into the wind, although not all facing that way when flushed. This is the first time I have seen them out of water, perched on Land.

•

r

.

Late Location and observations

Ecola State Park, Clatsop Co by Tom McCallister: large rafts offshore. March 26, 1947 March 26, 1947 Neahkahnie Mtn/ steady lines of this species in flocks of about 50

oct. 3, 1949 Huckleberry Hill, Lincoln Co. Oregon - 20-25 feeding 100 yards beyond the breakers for at least 3 hours in a.m., and in approximately the same spot. Oct. 5, 1949 Seal Rocks, Oregon. white forehead only, and feeding in shallow, rocky

Oct.16,1960 Davis Sewer Pond, Road 102, Yolo Co. Calif. 1:15 n.m. PST.2 immatures basically shades of brown, with head profile similar to that of Canvasback and with two large buffy-white areas on sides of head, fore and aft. Alice Williams saw one on Oct. 15th, 30X soope used today. Petty Kimball.



Observer ired Lvenden, Jr.


.mericum .coter

70-118-446-566-1079-110-1110-1877-2068-2347-2465-266-61-

Didemia Lmericana.

Late Location and opserv tions

reb. . o, i.e. or the torres. The rest is a very inc.



i illustro ele telire e delve e e e e e e e bediene e oril 30, 194; ouglacements-make aware anapara energy March 6, 1945 4 of today the second of the second in the second beauty to the second of the second o

Klamatk Lake Or

bent min-con-oucher

There were 100's, f. them in were of the wife in my adjusting March 27, 1947 Golden Gate Park. 1800 PST. 2 pair seen. males in full plumage and bill light blue color.

Sept. 20, 1847 Fern Fidge land i mule de translation de translation non-de l'étal de translation de la company non-de la company de la company

Oct. 30, 1948 Bayshore Drive, N of Bay Bridge: 930 a.m. rafts and rafts of them. 1000's of them on the bay just north of the bay bridge.

Nov. 13, 1948 Bayshore Drive, Berkeley-Oakland 7 10,000's of them on the bay area just west of the hiway, and north of the bay bridge.

Feb. 26, 1949 Toledo, Lincoln Co, Oregon (IBM) Jay Long stotted this one in wide spot of river east of Tolqeo.

Feb. 26, 1949 Yaquina Bay, Oregon (IBM) 6 just outside town in bay. ion-5,1249 Bay shore Drive, Berkeley - Oakland (\$350 + 600 +? + too far out to count + net. 2,1956 Autorograph, toper with. Wes serve, and a mile or will be a more

AND THE CONTRACT OF THE CONTRA

Location and observations

relli . The land to be discuss to chear, and a second this time of the second to the s _pril &5, 194;

200 Ol. 19 21 . 20 1 . 27 1 -

ruseing 1, Corveis Texas assumed in a room the bridge. In the years upsurant, lit, crecus on to no give edition of dealer, in our . chockish call note. That I this head it a uncold at the case that r cione tamine our. It circles and care ples events, aspara it ver, rajia dingbear y see satum one a jedin on a "in ling and a a", probably because a the sub-buck sines, I state store.

12-mile Swamp % a female seen with 4 or 5 young by lave, and later I saw May 13, 1947 the auult biras, but not the young. These were also eported by bb Storm last Saturday (May 10th).

May 20, 1947 12-mile Swamp: Orat least 2, and possibly 3 different females were been in the area. One had 10 young, and another had at least 4 that I know of. I found myself in the liable of them, and the F was trying to accoy me away. Then that wouldn't work she swam about in the open, with crest rully erected, and uttering a heron-like series of notes. As she swam back and forth, first one, and then another of the young a peared at fer side, and followed her closely as she swam away.

May 27, 1947 Faling Estate Swamp, east of Danile Swamy: saw 4 young (or this

species?) in open water, just above the dam. Form Ridge lam area - saw adults and joung (I didn't have a chance to count June 25, 1947 them, but at least 5 were seen).

Nov. 29, 1945 Camp Jeco Fond, Calaveras Co. Calif. a concentration of this duck on the pond this a.m. - 2714 & 56F or imm.

Jan. 10, 1 the Camp Seco Land. 15H and 23 F or imm.

- 12/12 Chinestague NWA, Va. Ps Courter, a a.m.

Widewater C+O Caul, MD. I faid W/ 1 Floof Cheed

lumber of field trips on hidle bestven:

2402-6- 2411-2448-2458-2460-2478-2498-2506-2523-2591-2665-2673-

Location and observations

Lec. 10, 1941 Mar 's liver, corville: " screen no a conjunt vivor a ville and a constant vill

alsea River mear head of tidewater Pair of brilliant matio in full sumage with red will and wit.

February 28, 48 Campe, delineton, Viesmo Wiles Wiles on the legal Link India of the Company of the transfer of the Company of bid. Tanda tad med ist had a cook.

Washingal River - Camas, Wash. Female catching a 5" trout in the shallow water

July 26 1944 McCall, Idaho. Saw male and iemale and 5 young on Payette river this evening.

McCall Idaho. Saw male and female and young (5) pretty well along in 29 1944 size, on this Payette R. today. Aug.

Avery Park, Corv. 4 p.m. 1 female in an eddy focket off the main part Jan. 11, 1947. of the Mary's River. When we flushed it, it used up 55' to 60' of water for its takeoff, and then flew only 100 feet or so farther on before it banked into another "cove" for a landing.

Apr. 27,1947 Rogue R., 6 mi about Illahe, Curry Co: 3 not quite in full plumage were on a sand bar by the river at 2 p.m. and had to enter the water to take off when we flushed them.

Aug. 24, 1947 :illamette F., Corv. What almost exactly 1215 noon I saw a group of towers" come around a benu in the river loss No make ustram fr a lose I put the glasses on them we exented habitor, as they are not us not many across the water to the insiec a the rant beneat the river. It to a treeapproximately 17 minutes to one their say countries, the length of the Arride of the curve, and into an entury rone and proposed and research the river from me. The pires term in the finale naumore, the all term of him but an size and combrities that seems like the country and country to the like the we distinguished the applitude of the remains it is to go incollect. There exists a while a in of the bittle button thing to and strated their regions are significant. displaying the bright white this pubch some unserest ruther. The evidently busy flicking, for they bid not receive, the mist called a shallow that Tree receives the shallow with a crease to other, adving, about an about that weem to be drawing such which with they prided over the rater both their head "cocked" to king or made the she countries but a retired; a site that bill and front part of the head supported the orch of the appropriation. nearly submerged with early the top anchor the formula to such the vent swimming entirony sub-rega, at unice time a definite increase in a cit a cou was noted. All i or them wid a , wer job of "Mi instal end" the try tendese to turn the rush, etc., into the shallow which who she will be accounted got a couple of ret from the later, charle to market, in our our time couling roll. Some of the birds, after even ing under a certifical annial the become to actually come up, jum installed of the statement and anely, the lot algebra at alr. They were not seen from 1245 to 1430, but at that time they sere evicently flushed from some judge further down them, for 5 eg list by Albert and landed opposite we, on the inside of the river curve again, the true legan "feeding" upstreum again. They found one on so, this care in a liver, und all of them hesitated there a few minutes and here observed to do some speciing, and just of in sitting.



- Feb. 1, 1949 Long Tom R, 3 mi. N of Monroe, Benton Co (THMCA)-flock of 40 together north of the bridge. Very wary. Cans and ring-necked were easily approached and flushed only after all the other ducks. 25M & 15F mergansers (?F) Males in full breeding plumage.
- June 21, 1949 S. Fk. Bumbolat River, Hylton lam Site, Elko Co. Nevada El goulg with female young would give and swim 20 feet under water.
 - Nov. 6, 1949 Bolinas Lagoon, Marin Co. 1030 a.m. a fairly compact group of 35 swimming and diving together on the lagoon. Only 3 full-plumaged males seen rest were "redheads".
 - Dec. 15, 1949, Stillwater Refuge Area, Chrochill Co. .evada 10 a.m. a F or imm in open ditch, with 2"-deep, 5" long Centrarchid fish in its bill. It drop ed the fish when we ap reached.
 - Jan. 4,1950 Feather K, Sutter Co, at Verona, Calif 6 on river, near sand bar. 5 p.m. Waters very murky, But Feb. 25, 1950 Sucramento R, Shaeta Retreat to Castle Cray, Siskingon County Calif 5 p.m. Waters very murky, But observed a single 9 or imm. and a pair drifting downstream along the edge.
 - June 12, 1951 Pyramid Lake, Nevada 1F seen with 6 young
 - May 31,1953 Lake Tahoe, Unper Truckee R. Adorado Co. Calif. 15-18mostly males, with
 - July 27-2, 1 16, Stony illustable, 70 tol. thorach to. while the plant in the half-groun paner residing a manage white lake, the half of wore seen at writing the ces about the lake and a till a we were there. It is that they all most of a new 10 feet all there.
 - Sept. 7-9, 1958 Zephyr Point, I. Tahoe, Mevada up to 13 at one time, all in F plumage.

.umber of fiel trib to line or .rv-1: = co-brountie lie sens or 1107-1110-1410-1533-1567-1810-1854-1888-1899-1901-1903-1904-2068-2289-2243-2259-2243-2259-22591

Lergus serr tor

Location and observ tions

..., i.e.. Lemmelmungt of main to be year in the contract of t

pril 80, 1948 | Corvan in: 0° and comment of the first of Viving Value of Value of

7-25-43 Washangal K., Januar, W.n. 150- J.W.T.

Netarts Bay, Til amook Co. 3 M and 6F. First noted early that morning

Mar. 27, 1947 Netarts Bay, Tillamook Co. 3 M and 6F. First noted early that morning which were going thru mating antics. M would swim toward F snap head down low against breast and then straight up in the air with a quick jerk. Sometimes they would raise the whole body from the water and then bringing the neck back down would prostrate themselves flat, the head and neck stretched parallel to the surface of the water. Occasionally the M and F would make a low cou hing sound. (Tom McCallister)

Feb. 26, 1949 Yaquina Bay, Oregon (IBM) saw perhaps 20. Never more than two at a time, and almost all were males.

And almost all wer e males.

Navarro R., man Allicon, Minderino Co. Calif. 5:15 pm.

Mellaru, Va. 1 Mello weet lotane, Kying other muster.

511cm 976, 2611- See Links Scientell 4176. 00/156.

٠ There was a second of the seco

igman gypominii i farthini d

The first one, 2 were shapered on the of region of the lead of the leading of the

Mar.12,1 6h, N of Ventura, Jentura Co.Cabif. somewhere over 5 m lafael its. and I of Fadres Forest - locked down from jet on 3 Condons circling ventura ring cumulus claud - shout 5:h/-(p.a. 7. Thank at 1/, 10) the large inventure.



May 11, 1040 Poyz it's it souther that the third on the Line of the its pour a. . a we at i . . . il il . . . Letween Jelianson and Alban Prote of the Commission in small and or the commission of the commission o Feb. 15, 1941 ween Jeneure on the first of the flyin very notice enching in small protes must be a shake the chief. iarch 11, 1941 were noise of them nor- than a core to cove the country. COTVERED COURSE CONTRACT OF A WAR TO BE STORED THE STATE OF THE STATE 1.2.ch 27, 1940 Reoria Ferry San 5 of them, 1 st of the year. Mar. 10, 1943 Corvallis. 0325 p.m. I was looking out the window of my room when my eye March 17 1946 caught two "specks" soaring a mile or so to the south. As they approached rapidly, I soon could see they were Vultures, first for this year. Dixon Creek Ridge, Corvallis Peveral of them were feeding on a dead Apr. 25, 1946 Black-tailed Jackrabbit. It was completely gutted, and most of the meat was still on the legs, etc. Just north of Ferguson, Lane Co. Came upon il of them, elch perchea on May 25, 1946 a fence post in a continuous group. Quite an oac sight to see. July 12, 1946 On gravel rosa near Sianey aitchnorth of Talbot? ** One was perched on a short electric wire pole as I drove up, so I coasted to a stop op osite it, and turned the radio music quite loud, and the bird, looked all around below it, trying to determine the scund, and then I whistled passages of the tune, and the viro almost real of its perch each time I aid, for it would cock its head to one side and then the other with rapid jerks. Finally it raised its wings, stretched then horizontally, and leaned off the pole top and flew away Indian Ford, leschutes Co. o young circled close around me as I stood Aug. 12, 1946 on a rocky outcrop near here. Theye were very curious, indeed. No redaish color was seen on their grayish helds. tewart Like 2 2 groups of 7, and 10 slowly cirching so the own the Sept 11, 146 country side. lelso, ashington - 2 or 7 reen Sept 24. 140 o dieno asmington - just one seen over the hills, a crea-Sept 25, 146 Mar. 7. 1947 Salem to Corvallis: two pair were seen, first of the year. durant Ly dordon, in the word that come in in no. Mar. 19, 1947 March 23, 1947 Joshua Tree N. Mon, Calif. 5 p.m. circling over the mountains NE of camp in a flock of about 45 birds. Joshua Tree N. Mon. Calif. 615 a.m. circling in grous of 35-45 from 0645 March 24, 1947 to 0800, and all moved off together, flapping their wings.

Oak-Fir belt, 3 miles W of 12-mile Swamp - one on the ground feeding on a

Lewisburg, Benton Co. It was so calm tousy that they had to flap their

12-mile Swamp several circled over the area. Crows were chasing one, and

quite old skeleton of a sheep.

July 24, 1947 Fern Riage Lam 9 of them feeding on Je succentrate sheet at a contract

wings continuously to keep going.

Redwings were chasing another one.

old highway.

April 8, 1947

April 11, 1947

May 6, 1947




Vulture Payre L.

- Maclonala lorest 2 young observed perched on the lower is or lines August 7, 1 947 of r. Fir in the heart of a Fir rore, ton a FF singe. The heads were barely colored. They would fly only a short distance before all fitting.
 - Peoria Ferry sailing over, and & seer in tops of talk snags, with Aug. 9, 1947 win's extended insum-pose.
 - Belifountain-Philomath I saw a few circling over the crea, and the 27 Aug. 22, 1947 were perched either on the ground or on lend. Josts, and lined up thoug the fence line up the hill.
- Themile tham one circles los over us. In ithatum the the the me Oct. 18, 1947
- head. Very late micront late cate record for the state of the one later.

 12-mile Swamp 1st or the year 3 birds seen, all going north, slowly. Mar. 9, 1948
- None over 50 feet above ground.
 Oak Ridge, 2 mi. 3 of Corvallis: 5 circling together, and 1 leutains March 27, 1948 circling above them give-bombed them as the approached the vicinity of the Lectail nest site.
- li-mile wamp down seen all over this fact, to down while, July 8, 1040 circling in small mades of a to 7 cm; .
 North Umpque 1, by how inches to make - for the common of
- July 23, 1945 in 1 at Punc. Sept. 25, 1948
- Grey Lodge Refuge: hone seen on way up here, and none seen till about 930 a.m. Many seen over area. Two feeding on dead jack rabbit. Sept. 26, 1948 Folsom Dam Site: scattered individuals that soon bunched in one thermal with others - 10 of them and one Redtail. The redtail made passes at a couple of them, so they headed due south in a broad front. About 2 miles south they hit another thermal where they joined more vultures.
- There were 27 of them in this thermal. Other individuals seen over area all morning. S. of Placerville: 460-70 circling in 2 groups, moving south
- Oct. 3, 1948 Oct. 4, 1948 Nevada - not a single one seen all day.
- Oct. 15, 1948 Nigger Hill, El Dorado Co: 726 circling in tall spiral to the south over open ridge at 1130 a.m.
 - Gray Lodge Refuge Tone circling over range area just SE of the refuge Jan. 8, 1949 at 930 a.m.
- Gray Lodge Refuge. Tat least 10 or 12 in area. Plenty of Took in dead ducks Feb. 6, 1949 and geese (mostly snows)
- Sacramento River south of Sacramento: 730 or so circling in 2 groups northward March 12, 1949 Sefferson Jet. New 4.5. 99E, Nof alkany, Orgon. - Many vultures circling over april 11, 1949 This area at 1:45 p.m.
- U.S. 99, N of Pitt a. Bridge over Sharta Lake, Shesta Co, Calif. 5- pm. 40-50 august 21,1949 Lest. 2, 1949 - N. of Chiloguin, Klamath G., Oregon - 7 seen at 9 30 a.m.
- Syst-11,1949 43 mi N. of K. Falla Oregon 25-30 at 11a.m.
- Sept. 21, 1949 Dr. Perry Reiche, U.S.B.R. Deologist told me that Aug. 16-20 he observed vulture mating activity on the flate west of Mendocino Mendocino Co, Calf, on the N w coast. Me saw them such day, once at only 75' dietance. One groups of 7 were on the ground in a 30'- circle, 2 were mating, with 9 17 completely exceed eagled. Some sort of dancing was occasionally lone by the bibba. Could they be nesting for a southern nesting.

 Detober 13, 1949. Lower Button Sinks N of Maria.
- October 13, 1949, Lower Butte Sinks, N of Meridian, Sutter Co. at least 250 circling over thermal of large brush fire.
- Nov. 16, 1949 Black Butte Reservoir Area, S tony Cr., Tehama Co. none seen until after 930 a.m. and then a couple. By noon I found several perched in dead tops of cottonwoods and later in p.m. about 15 were circling over the area.
- Nov. 16, 1949 N of Butte City, Glenn Co. Calif. about 45 circling in column of smoke from burning brush, at 4 p.m.



ą

Jan. 3 & 4, 1950, Black Butte Res. area, N & S Fks, Stony Gr., Glenn & Tehama Cos, Calif. Observed singles, and up to 5 in small groups circling over the nearby countryside, and then about 430 I discovered about 160 of them gothered or mathering in the upper half of several very tall Premont Cottonwoods; they could be seen coming from all directions in almost straight lines to this area of river bottom habitat along the D. Pk. of Stony Creek. The sun had left the volley by this time. They approached singly, not in grow s. watched two squabble and one chase the other for a couple of sinutes through the cottonwoods, with no regard for other perching vultures, of which several were knocked or over-balanced on their perches. At 5 p.n. I returned for a final count as it was nearly dusk, and saw no more coming in - the total was about 2x0, only 60 having come in during the half hour period of dusk. Checked the vultures at the roost at 9 a.m. (the 4th), and found almost all were still perched in cottonwoods, with one or two moving to other trees. Those perchang weren't all in the trees they had been in last night, but the reason for their listurbance was that a rancher was cultivating the field next to the original roosting trees. That the tractor noise disturbed the birds was shown by them moving whenever the tractor came close to the perches, whereas those birds perched in trees farther from the field were not disturbed, and remained on their perch. Some of the disturbed birds were perchad in a small group on a grassy tholl about 1/8 mile west of the roast trees at 930 a.m. This Fholl was in the oak woodland. At noon, another check was made of the roost area, and only a few birds were seen ercbing in the cottonwoods, and a few others were circlist over this and nearby areas. Evidentally the bulk of the 220 Lai left the area for the day, about 11 a.c. we saw vultures east and south of Orland, after noon. low far t is reast draws the birds from, I would hesitate to state. May 17, 1950 Black Butte Res. area, Tehama Co. Calif. 7 pairs, mostly perched one

May 17, 1950 Black Butte Res. area, Tehama Co. Calif. 7 pairs, mostly perched one pair/cottonwood or sycamore. Observed one bird fully spread-eagled in the early morning sublight, atop a sycamore.

Jan. 50, 1951 Near Thornton, S. Joaquin Jo Colifeveral seen - N15-50, 5503.

Dec. 18, 1952 8:30a. Softmaklin, Sacto. So, Skip. 2013 flying around - cell- 40° E

Apr. 19, 1954 Geysers Loop- Cloverdale-SE of Geyserville, Sonoma Co. Calif. I/ gopher snake that was run overon road.

Oct. 17, 195h, Sacramento Pyrass, Yolo Cr. Tolif. Posons moving sout of verying littless. Nay 16-17,1955 arounte's fine allow, while of the little horizon ach office.


Feb. 21, 1957 Several along american R. bluff Carmichael-Folson, Leets G. Celif. after storing night.
Many sean et Kedding, California.

Feb. 25, 1059 U.S.Hiway 50, near Strawberry, Eldorado Co. Calif. 1 circling high with 2 white pelicans at same time in same direction 1130 a.m.PST. to W.

June 22,1961 Bass Lake, Eldorado Co. Calif. 615 a.m. PDST. 2 in snag sunning selves with wings outspread.

4-18 cy - 100's our hack Oech area near should what we legen

June 28,1962 Bear 7. mouth at Feather R, ST Yuba Go.Galif. 6-1 am.12 s in cottonwoods perched and sunning. It ers on beach feeding on s awned out shad. Few circling over area.



1851-1852-1853-2226-2-228-2229-2235-2236-2237-224/-2242-2244-2609-

Black Vulture Coragyps atratus atratus

Feb. 21, 1951 Caterina, Dimmit Co. Texas - we knew these 6 souring birds the moment we saw them. Sawed-off tail, and graying patches near wing tips.

10-17-69 - Cowain Gupth, la - I hi, civeling t NE !! in mid-form.

Swallow-t iled Tite The noides forficatus

Matacumbe * Yer Targo below The noides May 20, 1965, Tear/Tomostead Florida - one chased by Gray Mingbird.

May 21, 1965, Everglades MPk Tla - as many as 20 at 1 time. A thing of true beauty in flight.

6/9/76 in a forthery the will all the



757-1783-1804-1815-1819-1898-1068-1068-1083-1183-1203-1256-1284-1290-1318-1321, 1355-1371-1382-1390-1421-1544-1552-1554-1617-1783-1804-1815-1819-1898-1901-1981-1888-2001-2041-2056-2158-2166-2168-2168-2169-2170-2206-2266-2273-2310-2312-2323-2323-2345-2363-2312-2396-2199-2491-2493-2580-2581-2633-2312-2396-2396-2491-2493-2580-2581-2633-Elanus leucurus majusculus White-tailed Kite

Sept. 25, 1948 Grey Lodge Refuge, Gridley first ones I have ever seen John Cowan took me to Porter's Pond where he knew they hung out. Seems they are quite sedentary in habits. The 1st one I saw was perched in the dead top of a $\frac{1}{2}$ -live willow with a Crow just 10 feet away, in the same tree. This bird seon flew low ower the open area and served upward to perch on a dead willow tree, where the

mate was already perched. Their habitat, here, seems to be scattered willows and cottonwoods, with mostly open marshy, or dry land in which it hunts. John says that from food pellet analysis they eat about 95% Microtus. Their nesting site this year was at a height of 11 feet on the SE side of one small willow adjacent to another willow, these two then being in a large open area. The nest was of flimsy structure. John banded 5 young birds, the adults would dive at him, tho never hitting. In the afternoon, both birds were present in the same area, but soon left when I walked thru their area. I did not see them again. They look just like the Black-winged Kite of India, that I saw at Karachi in 1945.

Oct. 2, 1948: Suisum to Grizzly Island Ferry: Pair flying over tule area below the hiway, as well as across the hiway over near barren hillsides south of the hiway. Watched a black-bird chasing one, and later saw them bantering with

each other. Also watched them hover like the sparrow hawk.

Oct. 2, 1948: Suisun State Refuge, Joyce Island: 'I hovering high over tules, but slowly lowering itself, while still hovering. Could have been one of pair seen along hiway in a.m. for it is only a mile or so away from the refuge.

Nov. 20, 1949 - 3. m. N. of Rio Vicia, toleno vo. Calif - 2 , Rovering over fieldest 3 Nov. 5, 1949, Marshes northwest of Vallejo, Solano Co. one perched on a post in one of the salt marshes, there. Another nearby, hovering above the marsh.

Dec. 6, 1949 Wear Moss Landing, Monterey Co. Calif. about 2 p.m. we observed one hovering for 2-3 minutes without pause - observed for 5 minutes. Kept dropping lower all the time but made no passes. Would lose altitude be extending wings in a vertical "V" over back, thus slipping downward. Within a-mile of new Steam Plant.

Dec. 7, 1949 - Calinas R. between Ring City and Met 5 p. . - watched one hovering, and making passes in company with a marsh hawk.

Lec 19, 1949 - Cottonwood Creek, Tohana- Sharta Co, Calif - 1600 P.S.T. Ion willow any downstream below hiway

Apr. 8, 1950, Lake Tashington, Yolo Co. Ill a.m. - one hovering at 75', held altitude, and I saw that the center tail feathers were gray.

July 18, 1950 - Waldon Krack, El Face Greek OF May han Joquin Villy of on Call I recommend to in cottonwood willow area.

Nov. 29, 1950, Near Lambert, Sacto. Co. Carif. I perched on wire lines 5 times, near market road. Gave "mex" call, too.

Sept. 7, 1951 Lake Washington, Yolo Co. Calif. 4 seen this a.m. -flying, and perched in willows at edge of lake. All seemed to be in ful! plugage. Jan. 29, 1953 - about 5 miles due west of Tudor, Sutter Co. Calif. 9:30 a.m. PST. Clear,

calm, and not cold. Along a dirt-gravel county road paralleled by willows Ralph Imler and I found 4 kites - evidently 2 pairs perched atop the willows or flying about and over them. Observed 2 of birds to head off from tree-top level, one behind and lower apparently chasing the other-the two klimbed steadily to altitude of 150-200 feet where "trailer" finally caught up to "leader" and I observed "trailer" to mount the back of "leader" for a few moments, perhaps only 1-3 seconds and then they separated, calling, and I watched each pull wings in close to body (tho not dlosed) and dive at about a 25-degree angle from perpendicular to tree level where they alit. During 10-15 minutes we also observed individuals to "float" slowly downward to treetop level from heights of 75-100 feet. Wings spread in broad "V" over back and with feed extended straight downward. Drop was not exactly vertical, but somewhat on angle. They also used hovering flight sometimes. Most interesting flight was horizontal over field at 60-80 feet with wings in "V" over body, With only very tips of wings beating rapidly - no inner wing movement.

17 mi Not lasto along rive- 1 ad. long chased by crown

•

~~

March 3, 1.56, Excelsion Road, Danie. On. 0.011. in the life of a land included its or mind by marchy being teal. I would be a land included in the land in the land.

Nov. 22, 1958 - 12 min N of Res Oar Colifornia Jet 25, 1960 - Skellen, Facto Co, Colifornia Excelsion Rd. Feb. 18, 196: - 12 min W. Sent. From C. Colifornia C. Colifornia Feb. 18, 196: - 12 min W. Sent. From C. Colifornia C. Colifornia

Oct. 4, 1961, S. Fork, Mokelumne R., opposite Sycamore Slough, 2 miles above Terminous, San Joaquin Co. Calif. Dale Grane saw and counted 17 white-tailed kites on a small island at this location.

December 9, 1961, 6 mi. W of Yuba City, SutterSo. Calif. along S.S.H.20. 3 kites over stubble field.

U.S.40 Jct. with Fairfield, Solano Co. Calif. 2 over orchard at 8:20p.m.PDST May 23, 1962 to might most, This Re or one in wire during our of do in field in tele such Sherrough Courties. Jan. 30, 1966,

5/70/14 along 49101, Thomas 5. of the Nobert + And Cloverbale at "pass".

| • | | | | |
|---|---|--|--|--|
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | • | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

Goshawk

istur atricopianus

Late Location and openivations

August 12, 1946 Indian Ford, leschutes Co. Saw one come in through the camp area fast, and it caught a Callo just like that. It was a young Callo.

Aug. 3, 1949 Bass Lake, Lacera Co. C lif. One charca a gray circum, but misson. Forest was very still afterwice.


May 30,1953 5 p.m. I full male took a yello-wheaded blackbird at the one leach Carsh Si cornerof Take Tahoe /tin view of allof us. It and otlers set up terrific commotion. The hawk carried the bird directly west across (right to oines. ctober 24-28,1954 - 140 Co. Nub. Willet, I follow, I follow, percent in the rest of the cost?

Lat



Observer Fred Ivenden, Jr. unbor of Field trink I Sharp-shinned Hank 11-36-65-75- 42-94-164-131-134-160-161-168-178-184-197-199-200-202-205-210-212 Location and ob. -ry tion Lept. 17, 1041 nako et melnakan di bibli bilan di bibli bilan di dikibirah, . Mangalan alam timbir bilan di bibli bibli bir pendangan di bibli bibli bir bir bir bir bir overtane a Vana - I to a pier it to a practite moon to a partial a lloca ol' nobina. U readum, Tregon, inc 83% in. Oct 21,1943 a sharp shinned Hawk captureda Robin + carried it away from our yard. Walla Walla, Wash. 8a. m. temp. 320- very forgy Nov. 14, 1943 One frew to the ground under a likacoush and stayed there a few seconds refore flying up into the bush. He was not ofterery, but might have been. : Wallula, Wash - wary Columbia K . - saw one longing to catch a fit House Finch. Dec. 3, 1943 Walla Wulla, Wash. 845 a.m. a skarge skin lit in the top branches of a 50' tree of watched Jan. 18, 1944 Kim fromous window, cost 125'amy. It agen wereste, he make a way for a forty thing. get them He just not in the con of womanie in it must a to me to me with in hat proverent. Last part & var! I wrote this to wing out the wester a in secure & inte to a thit interes! July 29 1944. McCall, Idaho, along Payette river. Saw one carrying a young robin away. Marion Oregon - saw one carrying a Vaux Swift, and b ing chased by 5 others. May 17, 1947 June 5, 1947 - Stewart Lake of taw one chasing and diving at a circling leated. Hawk by 20, 1947 - 12-mile Evamp Of one lie over the country tries, controlly of the second secon July 20, 1947 aug. 1,1948 Hogan Res., Calaveras Co. . made passes at violet-green swallows several Oct. 19, 1949 times, but failed to make a catch. Swallows gave alarm notes at its approa Sept. 9, 1951 Manzanita Lake, Lassen N. P. Calif. About 9 a.m. I saw a white-headed woodpecker fly out from the top of a very tall Jeffrey Pine and chase a sharp-shinned hawk, even tho the hawk had not come close to the tree in which the woodpecker was working. 1 4/27/14 Man Gassimille "4d. Elasal de Maser "buck Fret Gogly, Cherry Hitting, This it Treat Great below 13. Carry grand widge Need in Museus Place 9 & the Cooke of the Buy in congre Color to they 1 th Colinate broken

children with the grade of which the same is the



Cooper's Hawk

-- 1-17-11-24-31-44-45-50-64-46-142-143-144-146-149-161-192-199-296-297-352-435794-799-804-808-817-12-485-491-501-585-603-629-649-657-665-697-719-720-734-743-766-779...CCIDITOR COONTI 1076-1115-1132-1163-1164-1180-1194-1281-1352-1359-1385-1342-1419-1480-1508-1542-1542-1643-1770-1836-1839-1840-1850-2000-2013-2006-2013-200 2-7-19-22-29-31-44-45-50-64-96-142-143-144-146-149-161-192-199-296-297-352-435-

Late | Location and observation.

April 86, 184 (hive as very him. or no note to the second beautiful as very the second beautiful as ver

to contract the top of the contract of the con

12-mile Swamp, Bruce S tation: flushed one bird from the ground in the Jan. 18, 48 midst of a Sweetbriar thicket - its prey was a hen pheasant with the head and entrails gone.

Sept. 25, 1948 Grey Lodge Refuge: one adult flushed from the ground at the eage of the road. We could find no prey, but we did flush a covey of Valley Quail, which it may have been after.

Folsom Res. Site, Calif: 3 seen. One persistently chasing flicker, but .Oct. 13, 1948 the hawk lost out in the oak brush area.

Oct 19, 1949 Whiskey Creek, Halavers Rt, Calaveras Co, Calif, 1145 a.m. - one in the oak -D. pine woodland Dec. 11, 1949 Foisom Kee area, ElaDorado G. S. 3 m. 4 mi 5 of Corning otalama Co, Caly- 8 35 am. One observed in open country near a small row of Encalyptua trees.

Oct. 1,1253 Santa Rosa, Colif, to De at 1030 a.m. 3 tempers eiroli. o Logother.

Jec. 5, 195h Fotom Nos. 2003, clacur Co. Mili. Lat her on carmy 2000-4-117701 714 or away.

Sept. 10, 1961 Lassen Peak Trail, Shasta Co. Calif. - 1 along trail at 0,800 ft. elev. 5/24/29 Nogales Country Class Pous, Carry Calulton were to to be united as



opduing Paysent's part to the pain a men and a re-March lo, 194% Feb. 20, 1943 Corenthia, Seoria 7 my 1. nd OR. dan one pair maning was ther amind and a western in a consent tout tout I all 6-28-43 Camas W.n in a wong time Walla Walla, Wash. - Two of them hovering over a grantised brillaide & occasion 7eb. 27, 1944 diving at something on the ground, where by is. Albany-Corvallis. Saw an adult circling at 80-100 feet above a sence Feb. 13, 1946 row and all of a sudden pulmet earthward with folded wings, but ata checked speed. It pulled out at fence top level and wid not go to the ground after any animal, but lit on a fence post instead. The bus was going too fast to see what happened next. Stewart L, Corvallis, or Saw one being chased by two angry rouins. August 1, 1946 Indian Ford, Teschutes ? Saw a young one on the ground in a large August 12, 1946 meadow. I could not tell if it was feeding, or not. Vildword Camp to Prineville, Crook 0. 1030 a.m. August 15, 1946 Saw b young ones cirching over a mountain meadow west of the summit. They were acting the way that vultures do in carcling around. Feb. 18, 1947, Oak-fir belt between Beave r Cr. and Bellfountain: one perched on very top of D. Fir. Another perched on snag in more or less open area. Later circling up on cylinder and then diving off. Feb. 25, 1947 Oak-fir belt 3 miles west of 12-mile Swamp: flushed a female from an old oak, rather deep in the fir woods. It flew out over an open field and continued on. As we watched it get farther away, we heard a crescendo roar of air directly over our heads, but couldn't figure it out until about 300 yards off, we saw this "bolt from the blue" giving, past the bird we flushed, and with folded wings. Hardly had we finished seeing that and we heard another loud roar of air, and a second bird came diving at the area where the flushed bird was. The angle of descent of both these birds was about 60° from the horizontal. All three then chased each other off to another area, flying rather low. Later, when we reached the ridge-top, I spotted a red-tail sailing very high, in a straight line to the west. After about & mile it aropped its angle of flight a little, and gained speed, swooping upward, then, toward another bird that we hadn't seen circling before. After gaining sufficient altitude near this 2nd bird, it folded its wings and plummeted nearly straight down, pulled out, and climbed straight, almost, in a short loop, and from what we could see, it winged clear on over in the loop, folded its wings, and went into another dive, and loop - this was done a total of 8 times, before the distance away was too great for us to see. 3-20-47 Colo. R. Indian Res., south of Parker Ariz: 1 perched near a new nest in a tall, dead cottonwood 200 yards from the hiway. One souring will 3 Harristantes Mar. 26, 1947. Tilden Park Hills, Berkeley: several pair seen over Berkeley-Oakland hill area, soaring and courting during the time we were out this p.m. N side Rogue R., 4 mi about Illahe, Curry Co: nest on east side of D. Fir Apr. 27, 1947 against the trunk, at 60' in the tree which was on top of a hill. 2 mi. N. of Corvellis. Corgan hant ong its 1 2- max old you in a nict. May 4, 1947



- May 10, 1947 2 miles N of Corvallis: a pair were located circling over the hill area, and their nest was found at 75 feet in a more or less isolated cotton—wood, on an open southeast-sloping hillside. One young was reported in the nest last week, with a garter snake and 2 moles were in the nest, too. One dead egg, partly along.
- May 27, 1947 Faling Estate, west of If-mile Swamp: one pair circled constantly over one butte-like hall, and were observed in courting acts, once.
- June 5, 1947 Stewart Lake, latched Frewer Dlackbirds and Charp-chirned Lauk both chase a high-circling Rectail Hawk.
- August 7, 1947 Coffin Futte, near Camp agair, il a.m.: several circlin low over the summit of the butte. Lian't look skactly like reatail, and had immature plumage, characterized by prominent dark held and throat, with sharp
- demarcation between nech and chest, which was might.

 Nov. 18, 1947 Link south on Talomath of the median control of the median contr
- Oak Bidge, 2 mi. For Corvallis a pair circling over the range above me, March 26, 1948 crying at short intervals. They kept this up all the time I was clambering over the riage, and simally they came closer as I went down an open slope, one of the birds setting its wangs and swooping in from at reast _-mile away, but pulled out and hovered about 50 feet above me, screaming all the time. I suspected a nearby nest, but coolan't spot one, so sat down and waitea, watching the two hawks "talk to each other" with a set of notes (barking-line) entirely new to me, as they made partial contact, but they eventually circled way up and went ver the riage, and then I happened to look below me, and there was the nest. In a clump or ash and alders on a rlat leage at the very base of this steep of en slope was the nest, cout 45-50' up in an alder (about 22" PBH). As I was above the nest I could see something white, but it was so far awa. I couldn't tell if it was a downy young, or eggs. I will check later with more powerful glasses. The parents were not in the area all the rest of the time I was there.
- March 27, 1948

 Same: the two chased off the vultures, and soon a third Redtail joined them and they circled and sailed around, but no conflicts ensued they all moved west over the reage, and we cli bed the hill above the nest, but even with SEX magnification we club detect no eggs, or young. Inasmuch as courting activities were observed yesterda, perhaps nosting has not begun for this pair.
- Mpril 5, 1948 a-mile i. of adair Vil are, Donton Co. checked a nest at a -55 feet in a maple on a north slope, at the lower edge of the oak and brush belt, g-way up the slope. No sign of life, yet. In this clump of haples was one dead snag, riddled ofth large lares. In the very top, some is feet up, was a nest of light squirmis. Fraw 8 or them.
- Sept. 19, 1948 Natoma, below Folsom: immature seen. Nest at 30' in live oak.
- Sept. 25, 1948 Grey Lodge Refuge: Quite a few in the area, mostly immatures. John shot one last week that was preying on pheasants. We flushed an immature that had just killed a hen of the year (bursa still prominent). Talon puncture in thigh and back. It had begun to feed on the neck, just behind the head.
- Sept. 28, 1948: Ber Dam Site, Mariposa Co: Wa couple seen. Nest in Populus fremontii at 35' or so. Isolated tree near creek bed.
- Oct. 3, 1948 12-mile Swamp, Bruce Sta, Benton Co, Oregon: one was perched on a telephone pole along the railroad track. Around this bird on the wires were 150 goldfinches apparently unconcerned about the hawk.
- Net.2, 1948 Boca Res. Nevada Co 2 observed perched on very low sagebrush on north side of Prosser Creek valley below Hobart Mills. Birds were within 100' of each other. A strong wind was blowing.

•

-

•

- March 5, 1949 Bass Lake, El Torado Co. several seem, utilizing up-currents along N-S ridges, hovering in one spot. Saw one perched on top of an oak, whiche another was high above, doing the looping love gives described other year s
 - another was high above, doing the looping love gives described other year.s.

 May 21, 1949 Coast S. of rt. Lobos, Monterey Co. Feveral pairs flushed one second-year bird from the ground. Its "crop" was full and burging we found the rear 1/3 of a gopher snake near alpaso Creek, which it had left on the ground. It seemed to be a fresh kill.
- June 11, 1949 Corvalis, Oregon adult hovering over field of small hay stacks, like a sparrow hawk, and then plumged about 35 feet to the ground, and when I next saw it, it was just sitting there, and I don't know if it caught anything, or not.
- June 21, 1949 S. Fk. bumbolat h., neur 10-mile Creek, hako Co. at 350 p.m. I found a F circling close to us over a high rock obtero just has real from the nouth of 10-mile Creck. & nest sites on Al supper of pinnacle outcrop, within 6 feet of each other. Took pics. Lower n at like 3' high and used this Jear. I forma 2 joung piras perchea on a small sagebush a out 50 years uphark from the nest sit s. They were on a west slore, exposed to strong to winds this p.m. They were both open-mouthed and "ari] ing" from their mouths down the tip of the Lower bile. As lan Slater moved in to take a pic, one joung liew, ending up in tall sage growth at the edge of the river 4 leet below and over --mile away from the perch it left. The other young dien't c re to 12y, and pictur's were taken as close as l'away. I captured and balded to is young (#54-632488) size 7b.) One tresit, killed young jack musit, still gara, was found about 20 feet from the joins. Hearp, perches on mick outero s had many pellets scattered about, indicating that cottontail and jack abouts make up a good share of the adult's food in this area. I gave the young rabbit to the remaining ounghawk, but it payed no attention to it. Towns had rich, plann tawny brown area or chest, and was heavily by to: so below with dark brown.
 - Feb. 2, 1950, Folsom Dam Site, Sacramento Co. Calif. pair doing courting dives at 130p Feb. 8, 1950, Eugen-eMilton, Calaveras Co. Calif. several seen, one of which was nearly white-headed, and very light-bodied, but with dark wings, and a
 - red tail.

 Feb. 20, 1950 Lincoln, Placer Co. Wone flicked wings a bit, and another strongly when I honked the horn at it. Saw another carrying a huge stick into a large oak.
 - Mar. 27, 1951 SE of Lodoga, Colusa Co. Calif. 5 seen one on nest at 25' in blue oak on steep east slope.
- Oct. 17, 1951, Eugene-Milton, Calaveras Co. Calif. 1 adult with light golden-tan head. See ob for Feb. 8, 1950, just above.
- Nov. 7, 1952 Lake Washington, Yolo Co. Calif. Watched one carrying a large, but dead rabbit from road. Had difficulty, flew low, and alit frequently, and as I pressed him inthe car he dropped the load and circled away to return after I passed the quarry. Rabbit had been dead some time.

March 24, 1940 - Lauries Island O regar 2 eggs in next of dead streke, makings large attention, 45 lugarin continued. F.

April 6, 1941 Lauries Island, Oregon. 2 eggs in large of the next of 5 cf et ... a . thousand - Cultures

April 6, 1941 Lauries Island, Oregon. 2 eggs in large of the next of 5 cf et ... a . thousand - Cultures

March 26, 1942 faceries Island, Oregon - 2 aggs in large state at 1 2 11 crecks of a thorough Catters and the freshly killed face rat in nest. Next 70' in collowood. If on next. of soating above. Catters.

Nov. 7, 1952, Lake Washington, Yolo Co. Calif. immature carrying dead rabbit from road.

May 1, 1954 Sacto. River, below Knight's Landing, Yolo Co. Calif. 1F on nest in cottonwood. July 5, 1955 Sacramento Bypass, Yolo Co. Calif. Chased by crow.

Jet. 15, 1960 ge sate & Colf Shelder Welter Elligne and Saved paint conting Enter This part lar. 10, 1960 It. Tamplicas, Town D. 2011. actolog one horowing to the mon State win

Feb.18,1961 Pt. Reyes area, Marin Co.Calif. pair courting.
June 2, 1962 NE of Granite Bay, Felson Res. Placer Co.Calif. 1 with gopher snake

1.44

•

Krider's Red-tailed Hawk Buteo borealis krideri

Feb. 21, 1951 Carrizo Springs, Dimmit Co. Texas - head and shoulders almost white.

May 5-8, 1975 Monsons. Tueson. One name to telepho e the with mune man't.

Farlier it had brought a conher son o.

May 70, 1979. 57h. Each, Calp. About (1) on wise line. "Jellar capad" of public biggs,

body - beautiful! Then we talk and horizontal line should then sugar a fine light,

5/3/30 - Plumen La, (10/11) Mand by S.D. orienthad 18h.



Red- fellied Hawls

1100-1111-1133-1191-1251-1256-1293-1367-1375-1468-1978-1043-1048-1058-1069-1071
Buteo lineatus elegana

2226-2224E-2235-2234-2241-2242-2310-2410-2429-2436-24574.

Folsom Res. Site, Calif: one being chased by 5 crows. It lit in cottonwood, and wasn't bothered any more.

Jan. 3, 1949 - W of Bitterwater, San Benito Co. - one using small earth hummock for perch, on barren hillside near summit.of pass.


Jan.27, 1949 - Santa Margarita R., Camp Penaleton on erched in lower limbs of sycamore on west side of DeLuz Creek - was a rich rusty-brown on chest, tail distinctly and heavily barred with black and white, and rusty-chestnut on back between wings; Wings seemed light brown, with white spots the length of the trailing edge.

Feb. 17, 1949 Redding-Red Bluff at 5 p.m. I seen along the road.

Jan 14, 1953 - Saila Carlon, Carl. One met and was I saw.

Feb. 4, 1953 - along C.S.H. 120 just New of Stanislans R. near Oakdale, perched in an all walnut the laid livray.

Feb. 21, 1953 - Serval can along and R. Gluff - Carmithael to Folson, Suito 6, cold.



Buteo swainsonl

Late Location and observations

Dec. 3, 1943

Wallula, Wash. - along Col. R. - there were 3 of them that agent their time olying town over the sucks + geese on the river, but I saw no allempt to catch a sucks.

April 24, 1949 Franklin Ranch Floughs, E of Courtland, Facto Co. pair with newly-constructed nest in willow about 55' over slough.

June 18, 1849 Lake "ashington, Yolo Co. 8 a.m. one perched on ground in recently-cut alialia field where it captured a 50" popher shake. Then I wanted out to flush it, it carried the wreathing shake away. It lit if o yards away, and when I flushed it again, Freyer Plackbings pursued it Folyards, thile it still held on to the shake. It's flight was never over 6' above the fields.

Dec. 4, 1950 Pixley tulare Co., Salif. 1235, nron.

May 14, 1952 Pahranagat Valley nerth of Alamo, Lincoln Co. Nevada - 10 a.m. PDST. a pair with a nest in dead cottonwood less than \(\frac{1}{4} \) mile from highway. Nest a t height of 15-20 feet. One adult observed chasing a sparrow hawk.

May 12, 1952 - Carson River - 5 miles above Fort Churchill, Lyon Co. Nevada - adult observed on nest in living cottonwood at elev. of about 40 feet.

Oug 22, 1952 New Clay Factor Co. Celf.

7/26/54-Ed U.S. 395200 pet, SE Jussen G. (DP) 6 news small every I serve junipera in which were found 7 nests of all eyes, from 18' above ground to 16/est above, which was to take sof the Junipera.

Buteo albonotatus

March 18, 1947, Bill Williams River Delta - Arizona:

l seen. Not a foraging hawk, but seems to stay put in trees a great deal of the time. Average arrival date. Old nest in dead cottonwood at about 40'. Thin nest. Call or scream thin and whining whistle. Did quite a bit of soaring when we flushed it. Up to 500' high. The bill gleams at a distance. Nearly black. Tail bands show well. Could be confused with a vulture if you couldn't see the tail.

Dec. 1, 1947 - Letter from gale Monson
Renesting failed, for reasons unknown.

Coffee to a series of the seri

Reprinted from The Wilson Bulletin, Vol. 59, No. 3, Sept. 1947, p. 172

Zone-tailed Hawk breeding along Colorado River .- Since the Zone-tailed Hawk (Buteo albonotatus) nests principally in canyons of mountainous country from central Arizona southward, a report of its breeding in the immediate valley of the lower Colorado River on the western border of Arizona is of interest. On July 13, 1946, I found a juvenile Zone-tail, obviously not long out of the nest, perched on a branch of dead mesquite in the Bill Williams River delta, Yuma County, Arizona. At my close approach it made no effort to escape, and when I reached to pick it up, its only defense was to spread its wings and open its bill. I tested its ability to fly by tossing it into the air. It flapped clumsily and came to an awkward rest on a near-by bush. During this time the parent birds were flying boldly overhead, uttering a shrill screaming, and frequently alighting in a large dead cottonwood near by. Upon scanning the cottonwood, I discerned a moderate-sized nest of twigs about 40 feet from the ground in one of the main crotches of the tree. Since there were no other suitable nest trees close by, this was undoubtedly the birds' nest. A thorough search (as thorough as the denseness of the brush would permit) failed to reveal the presence of more than the one juvenile, and from the actions of the parent birds I judged that there were no more.

The general area is an almost impenetrable jungle of dead mesquite (*Prosopis juliflora*) and rank arrowweed (*Pluchea sericea*), from which protrudes an occasional stark, dead cottonwood. Close at hand is a small beaver-dammed stream. flowing through the dead mesquite and arrowweed; further away are willow-cottonwood woodlands and towering cliffs. The nest tree is about half a mile from where the small stream of the Bill Williams River joins the waters of Havasu Lake (impounded by Parker Dam).

In this same restricted area, I had seen Zone-tailed Hawks before: on August 14, 1943, three (including one I thought to be a juvenile); on August 25, 1943, two; on September 7, 1943, two; and on May 29, 1946, one. (I did not visit the area in 1944 or 1945.) My latest fall record for the Colorado River valley is September 16 (1946), when I saw one along the Colorado River, about two miles southwest of Parker Dam, in San Bernardino County, California.—Gale Monson, Fish and Wildlife Service, Parker, Arizona.

Observer Fred Evenden, Jr.

. American how h-lenge for k

48!-55!-199-418-435-481-735-999-1011-2008-

Dateo Lagorus gancti-johannis

i.te

Location and ob. Arv. thomas

eb. 22, 1941

hear Peoria: Ore It was a steel of coar 1.1 to only the You-bill have. color prown with a graph of the upper of this. It must have had a wing-spread of fire feet. Corval is, "if-mile awarp south of town.

Jan. 17, 1943

One flew over the mursh and enother was weaving in and out along a fence row trying to capture a trewer's blackbing out of a flock.

25 Aug '46' Avery Park, Corvallis. Oftaw this one sitting on a spag over the ary's River with Chickagees working the same tree, and under it was an Anthony's Green Leron perched on a log at the ease of the latter. Foon the hawk flew downstream, and immediately after the peron took off upstream. It must have been scared to death.

Sept. 20, 147 Greenberg Sta., bertan Co:02 and enjar of ortific to enterior a disc of late Let or and year parts of perched on a react port as a substitution area, this a.m. her it them, so calling about 40 rest up and to min verse marky notificate against the ine, have explication, a car a too is no in roughts below it.

Observer trea Feathern St.

renruminous soumeles **2651** 27019- 011-9-1021-2242-2332-2550-2554-2621-2622-

Luieo Populi:

Lute Loc tion and observ tions

Feb. 4, 1953 3 mi E of Merced W. MA., Morced Co, Calif, NW of El Nido. 1030 a.m. Very Eight individual.

July 7, 1957 11 Mer of Booker. Whate Sine Co November - Mar of the Comments.

Short-tailed Howk Bute brackyrus

May 21, 1965, N of Royal Falm turn in, Everglades MPRENE. I desh these adult chased by Master: Finebird. - with Sandy Corunt.



Harris Hawk

Parabuteo unicinctus harrisi

March 20, 1947

Colo. R. Indian Res. south of Parker Ariz: Watched one perched in a tree. Flusted as we approached and circled up to nearly 2000' and then another, the mate, before not seen, took off and circled up the same "cylinder" but not quite so high, and took off after the 1st hawk, to the north, and then, out of nowhere, came a 3rd one high and followed the other two. Field markings were visible up to a p-mile away, at least. The bird is generally a sooty black color with a white-tripped tail, and a wide white band at the base of the tail, above and below. A coppery tint showed on the upper and reddish brown on the lower surface of the wings. Tips deeply fingered, and wings had a rather "shaggy" appearance. One Redlail souring away with the 3 Harris Hawks.

apr. 28, 1947 . Letter from yale Monein gound a nest of these at the browny A. of towards on the wast.

3/14/63 Ef Rofageo, Ca Tep. 10-12/1 seen



Feb. 21, 1951 Rio Grande S of Lardo perched in tall willow beside the river.

Telly 12, 1977, to the control of th

. * •

umoni u riek tii : e li ol lvee:

Aquila chrysaetos canadensis

June 28, 1940

Late

Loc tion an obs statute:

.ay 17, 1941

from ramelia take Pricincles as over the last. There will be incorpress. Tygh Varley: "Female flow from not in v on in one top as about on a Level with the rock enounce the circulate willing.

lebruary 28, 1943

U-mas, distington: Prune Hill: All to the Amount very large Golden lighe them even und police value a temp nethre. As was only about Evise to bove the minner.

Mar 31, 1944

Mtn. Home to Boise, Idaho Saw one flying slowly over a sage-covered hill and drop, ing suddenly to the ground. He flapped his wings after hitting the ground, but we didn't observe whether he got any prey or not. May have been a jack rabbit, as there are plenty in the area.

Mar. 14, 1947

Winne are ... for your, with from our or

Apr. 5, 1947

Leadbetter Penin Willamette-Col Sloughs: immature (could also possibly by an immature Amer. Rough-leg Hawk). Dark brown with mottling below. Black tip to tail, whitish band inside it. Whitish underwings. Bail broad and fanned widely. Wings deep. Head appeared to be elongate and rather pointed.

Sept. 25, 1948

Grey Lodge Refuge: 2 immatures (windows in wings, and with basal 2/3 of tail white, and prominent black band near tip). They soared over fairly low, one held onto the tail of the other for a moment. These are the first for the year, says John. They usually winter on Marysville Buttes a few miles due south of the refuge, and spread out over the area for food, which usually consists of Jack Rabbits.

Sept. 29, 1948

Lingard to 10 miles SW of Merced, Merced Co. Colf. 1 2nd year immature perched on fence post. Another found dead beside the gravel road 300 yards from the live one. The dead one was pretty light, with all flesh, etc. cleaned off, and bones quite dry. I'd estimate it had been dead at least a week or 10 days. Mate still staying closs??

Oct. 23, 1948


Grey Lodge Refuge: a pair cruising back and forth over the area previousy occupied by the kites, which were not seen there today. The eagles would perch in low willows in open areas, or fly low to the ground, making passes at birds or mammals on the ground or in the air. I saw it pass at pheasants, coots, redwings, and mallards. None of the attempts were successful. They appeared to work as a team, one often being on the ground, with the other one working toward it in the air.

Dec. 30, 1949 Hogan Res. Calaveras to. Jalif. 1 i mature perched in cottonwood snag near L arm of recervoir - had I the patches on cheeks Lithin 200 yards of it I found where a white-beenster Canada Goose had been taken, wit' some wind an' many cobs of breast feathers scattered around.

april 23, 1952 Feb. 4, 1953 9-28-55 3/12/57

W Find Africand Refugo, Marcello, Calif. 10 a.m. I young of lect year on telephono pole - deary for this a.m. it was southern 5 of Gross Lake US 97, 5 is bigon to Calif. Hornbrook, Sishiyon Co, Calif. I seem man thinking.

Homing to NM - one philodes 100's of Ruche on Bolack to plose



97; -- 6 194.

Haliaeetus leucocochalum larcamus

Location and out iv ion.

Tort Bay: Who lit, the it was, in low to the riret I had ever and the get. In the total to the little New kt, Ivil autitudit voing ing inly

Mar. 7, 1944 Hot-Lake Oregon. Heavy bill. Head and shoulders mostly white, some bboth. He was on the ground in a pasture that had some brush. Maybe

it had a jack rabbit or other form of rodent.

February 4, 1947 Faling Estate, Benton Co. 12 miles S. of Corvallis. Two local men reliably report that a very large Gold Eagle has been present from time to time on the dill between The Faling + Mc 7 adden Swangs.

Philomather Wor without the Laster think they now me meating things in

Feb. 28, 1947 Fern Ridge Dam, Lane Co; perched in top of a dead tree, on a point at the south end of the reservoir. While we watched a solitary Raven approached it, and flew at it, as if to chase it away, but the Eagle just say there, and soon, the raven, which had lit next to it, flew away. The eagle was a male in clean dark brown and white plumage.

Jan. 5, 1951 Parcee Res. Amador-Calaveras Co. Calif. at 1 p.m. I watched one full adult eagle harassing a flock of 18 coots in mid-reservoir. Only those coots on the pathway of the eagle's flight low over the water dove, and the eagle, missing, would wheel and make another run. Finally it flew a mile to shore and alit.

March 19, 1951 Woodward Res. Stanislaus Co. Calif. 1 pair of adults, male in full plumage. F an immature?

Dec. 18, 1952 Hogan Res, Calaveras Co. Calif. 1245PST. 2 adults in tall dead will ww to east of dam, across reservoir. Jan. 30, 195h Hogan Reservoir. 1:15 p.m. 2, in two trees, with rost below one, a of dam.



Observer Fred Lvenden, Ji.

Marsh Hawk

at bir d'ains bluje a labor of sivet; 48-75-147-199-205-210-212-266-271-274-277-281-287-288-290-291-293-294-295-296-297-298-299-300-301-302-303-304-305-306-307-308-309-310-317-318-

la-mile swamp south of Corvallis ? Jan. 17, 1945

> 'e saw several of there toda, , and both of up saw his many is at least 4 of them at one time. They were a rather resty rea on the back and the white rump putch showed , wdimuy. I consider the dive and chase pheasants in an adjoining arela. It even chased a small bird (Song Sparrow). Then it will ove the close it dove and tried to capture mallerer and Finterer, but at monotonest enough. They now recove iros very mich as, a saturd being over 10 feet up.

Fortaguen La-mile eventy; The property of arch 6, 1948

inouting mean the creation to the rest in the good of a work age. nout i minima.

Wallula, wash, along col. K. - was a ., one was leeding on head attention trans of road. Dec. 3, 1943

Glenns Ferry to Hammett, Idaho. Three of these came soaring in, overhead. Apr. 1, 1944 th t I could not tell. They seemed to be almost white under wings and body, with black wing tips. No banding or barring of the tail. Top side appeared grayish. About 24-30" wingspread. One was observed to dive repeatedly and intensely at smaller birds, recwings, etc., in the brush on a river island. They could be White-tailed Kites, although I wonder !

Apr. 7, 1944 Mtn. Home, Reservoir, Idaho. Marsh hawks were in this area mand were constantly harassing the pairs of Magpies. As many as 4 were seen at one time. Two being white below and gray above; with black wing tips, and the other two being dark brown with some buffy striping, but all four had the white rump patch. A lot of the time was spent in chasing each other thru the skies. One pair appeared to almost come to "blows" which evidently is part of the courting procedure. They emitted their "squeaks" quite often. They were loud and plain. I observed, simultaneously, both males (grey) set their wings in a glide, let their legs hang limp below their body, and watched them glide to the ground, where they stayed for several minutes, apparently doing nothing, except for the occasional "squeak" note of theirs. These birds lit in o, en spots about 100 feet apart. Also these male birds would dive from about 200' straight down and when within a few feet of the ground, they reversed direction, and swooped vertically into the air to about 1001. winged over and dove again, and swooped up again to about 501, winged over and dove again, swooped up to about 25' or 30' and aid at least 5 more of these vertical dives and climbs at lesser heights. While doing this they occasionally emitted "squeaks".

- Mtn. Home Reservoir. Saw them doing more fancy diving, alighting on the Apr. 12, 1944 ground, and just sitting there. The male bird is always the one that does it
- Mtn. Home Reservoir saw some of them doing perfect loop-the-loops. Apr 17, 1944 -Mtn. Home, Area II, south of town. observed dark and light phases doing loop-the-loop dives. Flushed a female from the brush. Also, Joan and I located a site where a pair are constructing a nest. It was the male, light colored phase doing the building. We watched it carry nesting material to it.



May 8, 1944

Circus hudsonius

Mtn Mtn. Home, Area II, On Apr 17 we first found this marsh hawk nest, led Apr. 21, 1944 to it by the male, observed carrying nesting material to it. The nest is in a clump of old "reed or jointed" stemmed weed or grass. It is about 25 feet from a seldom-used farm lane and is about 150 feet fro. a gravel country road. When the hawk lands at the mest, it does so in plain view of the surrounding area. This site is located in an old abandoned pasture containing clumps of briar and the former described grass. Nearest water is over 4 mile away, and it is a nearly stagnant ditch, about two feet wide. The hawks broke down and crushed the center of this clump for a nest site. The nest itself is about 0 inches in diameter and from 0" to 1" deep. Construction is sloppy or Laphazard, and consists of dried stems of weeds and grass. Mtn. Home, Area II, Joan and I approached to within 8 reet of the nest, at Apr 24 1944 which point the female (dark) explosed from the nest site and went away. The nest had been completed since last observed on Apr 21. The nest cup was grass lined. About 1" deep and 7-8" across. Quite neat, now. There was one egg in the nest (took picture) Egg was white, and bluntly oval. $1_{\frac{1}{2}} - 1_{\frac{1}{2}}$ " x 2" in size.530 p.m. Apr 26, 1944 Mtn. Home, Area II, Today the marsh hawk nest had 2 eggs in it. Apr 28 1944 Mtn. Home, AreaII, Nest now has 3 eggs. Apr 29, 1944 Mtn. Home, Area II, nest has 4 eggs. Flushed female from nest twice ina half hour (took a picture). It seems that an egg is haid every other day, Up to today it has been laid on the day between the days we have ugs. 24 ups 26 ups 25 yas. 29 may 1 may 2 may 4 may 6 may 8 may 10 may 12 1egg 2 eggs 3 eggs 4 eggs 5 eggs 6 eggs 6 eggs 6 eggs 6 eggs 6 eggs 6 eggs may 17 may 18 may 19 6 egg2 6 egg2 6 eggs Mtn. Home, Area II, nest has 5 eggs in it (took Picture) May 1, 1944 May 2, 1944 Mtn. Home - 5 eggs in the nest. The male was on the nest and stayed longer than the female would have, I as I approached. He made a lot of scolding notes as he took off and as long as I was near the mest or within 50 feet of it. All this time the female was observed to be perched on a fence post 200' away, while the male was taking over incubation outies. May 4, 1944 Mtn. Home 530 p.m. Flushed female from nest containing 6 eggs. She stayed close and when I stooped to take picture, she set her wings and dove directly for me. Mtn. Home. Flushed female from the May 6, 1944 nest again, this time there were still 6 eggs. Evidently all that are to be laid. Each day, she stays

May 10 1944 Mtn. Home 4 p.m. The female stayed about 10 feet from me all the time I was at the nest and never over 15 feet high. She followed me all the way

longer on the nest, as I approach.

near the nest.

Mtn. Home Female stayed on the nest longer this time and then never went over 10 feet from me while I was

from the nest and never over 15 lest high. She ioliowed me all the way from the nest to 500-600' away. "Keeking" at me constantly; Finally she left me and went directly back to the nest. Evidently eggs will hatch in a few days. (??????).

GENERAL NOTES

Nesting of the Marsh Hawk at Mountain Home, Southern Idaho.—On April 17, 1944, we spotted a male marsh hawk carrying nesting materials into a dense clump of weeds in an abandoned pasture a short distance from the south end of town. The site was in plain view of a well-traveled gravel road leading to some farms near by. The nearest water is more than a quarter of a mile away.

Upon further investigation we found a nesting site, partially completed, in this clump of weeds. When we visited it four days later, it seemed to be completed.

When we visited the site on April 24 there was one egg in the nest. The female was on the nest, as she was nearly every time the nest was later visited. Once, on May 2, the male was on the nest. At that time it contained five eggs. Several times when I approached the nest, both during the period of incubation and after the young were hatched out, neither adult was present on the nest.

The dates on which the nest was visited and the number of eggs found were as follows:

| April 241 | egg | May | 15 | eggs |
|-----------|------|-----|-----|------|
| April 262 | eggs | May | 25 | eggs |
| April 283 | eggs | May | 46 | eggs |
| April 294 | eggs | May | 196 | eggs |

During the period from May 20 to May 27, I was unable to visit the nest at all, but upon my return on May 28, I found three young and three eggs in the nest.

Remaining dates that the nest was visited, and the number of eggs and young found, follow:

| May 294 young; 2 eggs | June 15 young; 1 egg |
|-----------------------|----------------------|
| May 304 young; 2 eggs | June 26 young |
| May 315 young: 1 egg | |

From all indications, it would seem that the eggs were laid on alternate days, beginning April 23; i.e., April 23, 25, 27, 29, May 1, and May 3.

When I returned on May 28 and found three young and three eggs, the young were of varying sizes, thus indicating that they must have hatched as they were laid, with incubation beginning with the laying of the first egg on April 23. As four young were found on the 29th, it seems almost certain that the three other young had hatched on May 23, 25, and 27, in the same order as they were laid. The last three eggs to hatch substantiated this assumption. The incubation period for each of the six eggs was exactly 30 days, as can be gathered from the above facts.

By June 15 the oldest young had reached a size of 10 to 11 inches in overall length, with the wing primaries about 5 inches in length. Most of the body was still covered with down, however. By this date the entire nesting site and all of the weeds surrounding it had been thoroughly beaten down, and the birds would be found anywhere within 5 or 6 feet of the nesting cup proper.

I am afraid that both the young and adult marsh hawks met an untimely end, for when I returned to the nest on June 18, no sign of the young or of either of the adults could be found. Ranchers had been working in a neighboring field during the previous few days.—Fred G. Evenden, Jr., Oregon State College, Corvallis.

Circus hudsonius

May 12, 1944 Mtn. Home, Area II, Idaho. Still 6 eggs in nest. The female was not on the nest when we approached, but came soon thereafter. She stayed within 15' or so of the two of us all the time we were at the nest. May 14 1944 Mtn. Home, Area II, The female was on the nest. Just 6 eggs yet. None hatched. She stayed near all the time we were near the nest. No sign of the male. May 16 1944 Mtn. Home, Female still on nest. 6 eggs, not hatched yet. Female did repeated and continuous dives at us, pulling out within 2 or 3 feet of us. May 17, 1944 Mtn. Home, Idaho - the female was really fussy today. She dove at me repeatedly and came within a foot or two of my head, each time ! May 18, 1944 Mtn. Home, Hawk stayed on nest till I was right at the nest. She dived on my head and circled over me calling and scolding. Eggs still not hatched - she staged in the air scolding till I left the territory. (Joan) May 19, 1944 Mtn. Home 9 a.m. Hawks nest still with six eggs. hawk reacted the same as on the 18th. (Joan). May 28, 1944 Mtn. Home, Area II, Idaho. 5 p.m. The female was on the nest and when she took off, it was at me. She came uncomfortably chase all the time I was at the nest. There were 3 eggs, and 5 young in the nest. The young seem to have hatched at intervals as variations in their size was evident. The body of the longest was $3\frac{1}{2}$ " in length. Eyes are open. Bill black, mouth pink. Natal down covers the entire body and wings. It is a buily or pinkish cream in color. May 29, 1944 Mtn. Home. Today there were 4 young and 2 eggs. The newly hatched was about the size of the largest young. It could be they are hatching 2 days apart, as a result of being laid 2 days apart at the beginning. If this is so, the first hatched on May 23, the second on May 25th, and the third on May 27th, and the 4th definitely on the 29th. It so happens that egg one was laid Apr. 23, egg 2 on Apr. 25, etc., making exactly 30 days from laying to hatching. We'll see. The female was not at the nest, but came soon after I got to the nest. The young were peeping loudly, with mouths open. It was very hot and they may have been dry or thirsty. The male and female were 200 or so yarus away when I got to the nest. May 30, 1944 Mun. Home. The female was on the nest again at 3 p.m. as I expected, because we had intermittent thundershowers all day. One instance, we had .17 inch of rain in 30 minutes. This was 2 hours before I visited the nest. Again, she came straight at me, as I approached the nest. Still 4 young and 2 eggs in the nest, which helps to prove conclusions I made yesterday. apr. 17 apr. 21 apr. 24 apr. 26 apr. 28 apr. 29 8nay 1 May 2 May 4 May 6 may 8 may 10 May 12 May 14 Ming 6 or Belg. Complete 1 egg 2 opp 3 oggs 4 eggs 5 eggs 6 eggs ? ASSUMED?

May 17 May 18 May 19 May 23 May 25 May 27 May 28 May 29 May 30 May 31 June 1 June 2

6 eggs 6 eggs 6 eggs 5 eggs 4 eggs 3 eggs 3 eggs 2 eggs 2 eggs 1 egg 1 egg 1 egg

May 31, 1944 Mtn. Home. 11 a.m. Female on nest. 5 are hatched. One egg. The largest young is now about $4\frac{1}{2}$ long. No feathers or quills. Still thoroughly covered with down that is a distinctly gray color. No sign of the male. Two of the females breast feathers were lying near the nest.



Circus hudsonius

June 1, 1944 Mtn. Home, Idaho, Area II, 530 p.m. The female hawk stayed on the nest quite a while in fact, until I was right over the nest. I whist led all the time as I approached the nest. There are 5 young, of varios us sizes, and one egg!

June 2, 1944

Mtn. Home. 530 p.m. .30 in of rain today. Water puddled on ground.

Female again on nest and male came this time, when she started screaming.

There were 6 young tonight. They appeared a little damp, because their pinkish skin showed thru quite a bit. Still, just down-covered; As the female jumped from the nest, the young were observed to be tumbled about. Evidently they were inside some of the female's chest and belly feathers, for warmth. The female made one very long glide-dive at me, after I had left the nest area, completely. Distance about 300 yards.

June 4, 1944

O830 a.m. Mtn. Home. Female on nest, and she gian't see me till I got right on top of her. No feathers are present on the young, as yetstill all down-covered.

June 9 1944

5 p.m. Mtn. Home Ames II SWIO n. The female real and all all down-covered.

study began April 17, 1944, just two months ago!

5 p.m. Mtn. Home Area II. SW10 r. The female was huddled over two of them, but the other four young were just around her, and one was outside the nest. The largest is now $7\frac{1}{2}$ " long and stands a good 5" high. It is still completely down-covered, but the wing quills are 2" long, and the tail quills $1\frac{1}{2}$ " long. The head and legs are well developed. It even snapped its bill at me as I came close to it! The female scratched my scalp with her claws on one dive she made today!

Area II, Mtn. Home 9 a.m. The female was about 600! from the nest, very busily engaged by pursuing Brewer Blackbires, as I approached the nest. She came in a hurry, when I reached the nest, and proceded to dive-bomb me vigorously! It was a good thing I had a helmet liner on, because she hit it 3 times while I was at the nest. There were only two young in the nest, the youngest and the third from youngest, the others being scattered under the surrounding weeds. I got a picture of the two on the nest, with the larger scolding and defying my presence. Here is a description of the largest young bird: Body was 10-11" long, and still precominantly covered with wooly down of a dirt, gray color. Bill is yellow on top, has a black tip and sides. Feet are yellow and $4\frac{\pi}{2}$ " long. Eyes are dark brown. The wing primaries are 5" long, and a dark gray-brown color. A few feathers and feather quills are present on the shoulder, and under the wing on the body. The head is 12" wide and about 2" long. By now, the nest has lost all shape and form, the area in a 3 foot radius of the nest now being welltrampled down, and open. The birds can be seen from 3 different directions, as far away as 50 feet. There was no sign of the male being in the vicinity. Mtn. Home 5 p.m. Area II. West to see how the young hawks were getting along, and not a sign of them around the nest. I dian't see any adult hawks in the area, either. I searched the surrouncing brush and grass for 100 feet around the nest. Evidently the acults have moved the young, because I know that at least 3 of them couldn't fly ! How far did they move them? How? Why? Nice questions to be answered. Unless I accidently run across them at a later date, this concludes the Marsh Hawk nesting

June 18, 1944

June 15 1944



Circus hudsonius

- Sept 19, 1946, Poiss: Can Lake, bins to. one made came in the selice, and circled over the take at 1-300 feet a few times and them processes on a . I believe this is the limit I have seen since measuring in motive service.
- 12-mile Swamp: Oras we approached the swamp, a single gray-white male was Feb. 25, 1947 seen beating slowly northward against the wind, and while we watched, he left the swamp, and cruised over a field of weeds east of the swamp, just a few feet above the dead weeds.
- Fern Ridge Dam, Lane Co: a single female flew low over a raft of baldpates Feb. 28, 1947 and mallards, but they showed no alarm, and it made no pass at them.
- 12-mile Swamp: a single male over fields near the swamp. Alit on the ground Mar. 11, 1947. several times and just sat there, so far as we could tell.
 - May 6, 1947, 12-mile Swamp. a female over the field east of thw swam, chased by Reawings, and Brewer Blackbirds.
- 12-m.le Swamp: Onone seen today, but I found last year's nest in the center May 13, 1947 of a clump of Carex oregonensis - a similar sight to the one found in 1942
- (May 5). li-rice charge Lab unitain reach an interior over the product of the production of t
- The disc is a policy of the second of the se Sept. 10, 10.7 Lot one in the order its of the Leo-tyl. The term of the special because in the discourse in the respect to the circumstance of the large grounds as it is in the contract that the circumstance is a second of the circumstance o the lime over, one has to the state of the s line en men terment a comment to the feet to the section
- 12-mile Swamp: of female or im ature of last season over the swamp area. We March 9 1948 saw them make passes at Reawing Blackbirds, M & F pheasants, and Gree-wing Teal, but with no success. Except with the Teal, it wasn't at all persistent in following prey.
- March 23, 1948 12-mile Swamp both H & F seen. Also fould fresh kill of Coot at eage
- of water in pasture area.
 Mills-Fair Oaks, Sacto Co. 1700. 4 or 5 pairs over the gravel-barrens. Mar. 25. 1949 Watched one female or immature male doing loop dives of decreasing height with complete wing-over at top of loop.
- Mar. 26. 1949 Lake, Washington, Yolo Co. 7930 a.m. For immature doing looping gives, with wing-overs at top of loop.
- Sept. 27, 1949 Sacto-Franklin, Calif. one dipped desparately when I honked the horn
- Nov. 16, 1949 8 miles north of Butte City about 2/road mile harvested rice area. Dec. 7, 1949, Salinas n., hing City-metz 7 3 p.n. Observed one making passes at the ground, in company with a hovering white-tailed kite
 - Dec. 13, 1.49 Lower Lovelock Valler, Fershing Co. Lev. 445 Observed analeone following behind a short-eare lowi, and when the owl would miss, the hawk would make a pass at the ground.
 - Feb. 20, 1950, Lincoln, Placer Co. Calif. watched a male cross hiway, and nearly run into electric wire line between poles, as it had to twist downward to avoid them.
 - June 9, 1950, Yolo Bypass, Yolo Co. 1230 p.m. several seen, and IF mushed from freshly-killed 2-gro. n pheasant. Eaten.
 - March 19, 1951 Woodward Res. Stanislaus Co. Calif. watched young male doing loops and dives of courtship.
 - Dec. 13, 1952. From Lodge Refuge, Butte Co. 2 p.m. Watched 2 making regrested dives from 5-12' Rt. at placements. 7 husbed 3 while we watched.

- Feb. 14, 1953 Chaves Co. New Lexico. Observed a few flying lengthwise along the highway at dawn (6:15 a.m. MST), 20 minutes before sunrise, which was at 6:40 a.m. Saw one alight, and drag "loot" from highway as we appreached. Apparently have caught onto easy breakfast thru road kill of previous night.
- Feb. 14, 1953, E of Plains, Yoakum Co. Texas. At 8 a.m. we saw a marsh hawk divin g at a coyote in open field. As animal ran the hawk followed, swooping.
- May 30, 1959, Carson Valley, S of Genoa, Douglas Co. Fevada 350 n.m. TDST.

 2 nests, 3 5-week old young at one and 3 1-week old young at other 150 yards apart. All 6 young banded and photographed, also setting.

2/4/67-5wf Dulce, N'14x. One in charact pure yellow pine orea.

* * * *

•

.

OSDIEN 1196 - 1275-1566-1567-1619-2043-2016- 83-167-331-337-602-607-609-754-857-869-877-893-923-924-971-1145-1175.

Panaion heliaeetus carolineasis

Lute

Location and absence tie.

Aug 18 1944

McCall, Idaho. Saw one this evening about 50 feet above the lake. It seemed larger than a Red-tailed hawk, and mostly white below. Wings bent like a gull.

Oct 10, 1944 Kings Hill to Glenns Ferry, Idaho, 10 a.m. to 3 p.r. Notes on one I observed from 1335 to 1415 this afternoon:

It came up the river, about 50 feet above the water, hovered, and dove out of sight into the water, before I realized what it was. As it came up, it seemed to have something in its talons, which later proved erronious. The bird had its right foot deformed, either club, or otherwise. Just about the time it had shaken the water off its feathers it met a female Marsh Hawk, heading downstream, as the osprey flew upstream. Both binds made some high screeching or squeeking notes at each other, and the osprey gave way, and the female marsh hawk continued on in its intended flight, passing almost directly overhead. The osprey circled directly over me several times, but I didn't seem to both him in the least! He circled around, and then hovered over a particular eddy, at the edge of the river. It hovered in such a position that its line of sight was 500 or directly crossing the rays of the sun to its right, and also so that it was "down-sun" from the objective in the water, thus not casting any shadow on the later. By this method of sighting and searching with those telescope powered eyes, it eliminated reflection of the sunlight from the water:

Cuite a strong wind was blowing. All gives were made upwind, or and quartered into it.

Dives were made at about a 60% slope from the water surface, and were usually sta ted at a height of from 40 to 60 feet. Wings were only partially folded at the start of the give, but becoming more folded as the dive progressed. They were closed, or folded tightly, just prior to the birds contact with the water. Its entire body went below the surface for about 10 seconds. Then, using its wings to pull its body to the surface, and out of the water, it would fly just above the water, up-wind, shaking its body, tail, and wing feathers vigorously. After two un-successful dives, probably caused by having only one good talon, it flew to the other side of the river and landed in 4-6 inches of water. It did nothing but sit there, in the water for about two minutes, and then it started to bathe, using exactly the same motions that I have seen a robin use, in our bird bath, at home. This bath lasted for at least 7 minutes by my watch, and it was very busy the whole time. When it had finished, it just stayed in the water and looked around. A green-wing teal flew by, circled, and landed within 5 feet of the osprey. A magpie came down and landed on the gravel beach, just 2 or 3 feet from it. Evidently both of them thought it was a cuck, or else were very curious! Oh yes, later on, I also saw it miss two other gives into that same eddy.

Apr. 22, 1947

12-mile Swamp, Corvallis part before leaving Dave Marshall spotted a "hawk" (1645 PST) flying over the swamp. We saw it was an Osprey, and it was being chased by Redwings, and did a "snap roll" to evade them and continue on its northward course. Had a fiar view of it. As it went over the north end, most of the ducks took off, and spread out in all directions.



- October 21, 1947

 In mile a conjugate of the conjugate of
- April 20, 1946 12-mile Swamp: One down in scages at south end. Thus call the next gave 3 sharp "jeeps" and Thew south a hitthe before crambing.
- June 23, 1948 Cottage Grove I reavoir? harron "hack" conditional element located an actual organization had had been collected an actual organization with another collected over a region to the concentration, and then a lamature bird, in around a restriction, and then a lamature bird, in around a restriction to the lama, a real amount, and the second or actual.
- July 28, 1940 Eavi: Luke: Thyin with move a character of the bar well.

 In the a.m. I materia one cave then the same of an action of the same of the s
 - Apr. 24, 1950, Fel R. Telta-S end of Humbolat Bay, 1 530 p.m. I pair carrying targe clumps of what appears to be seawed or eelgrass. Both birds circling in snort-radius circles to gain altitude, and then they headed east from the south end of humbolat Bay toward the higher hills to the east.
 - Apr. 25, 1950, N. Humbolat Bay, near Samoa 12 p.m. watched one dive below the bay water surface and take an 8-inch bay smelt, and then witched it escape from harassing gulls as it circled high over the bay and headed eastward toward the hills.
- March 23, 1952 New Carpar, Mendosino Co, Calif. (c. 5, H. #1). 9.45 a.m. Carrying 1 fish.
- March 23, 1952 Rockport Bay, Mendocino Co. Calif. circling over bay area and mouth of creek called with shrill whistle several times. Made two unsuccessful dives into bay.
- June 27, 1954 Pope Beach Spit, Take Tahoe Calif. Fair circling and fishing over marsh. Watched on e carry 10-11" fish down the beach to nest we had previously located in dead Jeffrey Pine snag about 25' high. Never saw bird actually go to nest, tho it landed on limb near nest. Mountain Chickadees had nest at 10' height in same snag, on side away from lakeshore. We found several catfish and hardheads or other minnows all dry, and mostly headless, lying on the ground below the nest.
- Apr.1,1961 Bodega Bay, Sonoma Co. Calif. 2 feeding over bay area, would dive from 100' or so, take fish, fly to fence post, or even on mud flat or dry grassy knowl to feed. Fish to 15-16" or so. Watched gulls chase osprey from their catch. Took pics. Gulls took over fish.
- Aug. 7, 1763 Canyon Jet. Area, Tellowst e W. K. No. Pain showers 2 at Alum Oresic and Yellowstone R. bove. Also pair with 2 months in nest or rock innacle at ookout foint below Current Jet.
 - The Cooks Key Chisfort, Long & Fla Next atop stack town 75th 2+1 gry (2 = 3) more of the Cooks Key Chisfort, Long a Fla Next atop stack town, w/ east fry the formy mothing across an backwise they.
 - 3/21/11 Execution 7 de la comme de la company de la compan



POLYBORUS CHERYWAY

PRIOUSON CARACARA

6/18/66, E of Meere Maven, Morids. I me swith like sere to temperature

| | | , | |
|---|--|---|---|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | ٠ |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| • | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

Observer Tree Evenden, Jr.

Prairie de Con 1823-1861- 2001-2089-2101-2104-231-

luco municura.

Lute

Location and observations

Feb. 12, 1949

Nigger Hill area, between forks of Amer. R., El Dorado Co. I seen about 3 p.m. cirling southward in tight circles against the wina, without using the wings too much.

Oct. s, 1950. Toulon Sink, Humboldt Fink-Lovelock Valley, Nevada: 1 icinoued among jole line ahe a of car at 30 mph and it was not using its wills all the time, either, sould perch atop a pole to rest, and then say on a see a preache a.

18 mi NE of Literfield, Lassen Co, Caly along 4.5 395.

the second of the second of

Falco pereminus anatum

Lati Location and observations

Feb. 11, 1947

12-mile Swamp. Ta single dark phase flying low (50-60) over the open swamp. which I first called a Cooper's (by the size), but it very definitely was a large falcon, with the pointed wings, presumably of this species.

Sept. 25, 1948 Grey Lodge Refuge: 1 flew over as we flushed some pintails. It made a fast pass at them, but they veered, and he didn't pursue further. His flight was very, swift.

Oct. 23, 1948 Grey Lodge Refuge: 1 in area made pass at cinnamon teal, but missed.

Nov. 25, 1948 West Moreland Pond, Portland, Oregon: watched a dark duck hawk for some time disturb all the birds around the west pond. The baldpates on the lawn took to the casting pond. Domestic pigeons and pine siskins flew about with the duck hawk flying in and about them. The hawk did not take anything and finally flew over towards the east pond. D. B. Marshall.

May 21, 1949

Coast Coi Car mal, Interes Colificing La materials and allow to a tocaling but even after long search we did not find it. The fetale was larger and lower toned in color and voice than the male. Had the time we were searching the cliff, both woulds were crying, and circuing against the thermals on a nearby west-facing fliff, sonctines coming over us.

June 23, 1949

Levil's Gate Lan Site, N. Ek. Mumbolot F., Elko Co. France: no that 25 % fect on east-facing chilf, wove river. Fort in crevie-, who I couldn't see into it, or reach it, t'e I trice. Addits no ray. Fold not sites Itung murby in sililar liches in the rica talk.

Feb. 4, 1953

E of Moreed Refuge, Moreed Co, Calif. I on pole line - rich Olive-gray back.

Feb. 27, 1953 E of Tatum, Lea Co. New Hexico. 2 p.m. 2 duck hawks seen.

Mar. 1, 1953 N of Tonopah, Nye Co. Nevada - 9 a.m. 1 on snow-covered greasewood.

My 18, 1954 Sura Oaks, Secto Calif - 6 pm.



Observed lied venden, Jr.

Peale! / Inlor

Tumo troi liela tripa tri i i . . . v. u: 566-567-

Falco peregrinus pealei

Location and object voltions

March 26, 1947 Neahkahnie Mtn one adult bird circling over the mtn. by T. McCallister.

Observer Fred Tvenden, Jr.

Black Pigeon Hawk

Number of field trian or lied our er vou: 252-254-446-472-573-721-

Falco columburius suchleyi

Late

Locations no observations

Oct. 5, 1943

South Tacoma Swamp, South Tacoma, Washington.

This is the first one I have ever seen. It was about the size of a sparrow hawk, the tail was square, and the body color was very dark. I'm not sure, but I think that there was some long-itudinal striping on the underside. It flew low overhead, and then out over the open water in the marsh.

Lec. 9, 1946

Corvallis, 1000 a... no seem in a marmolin or truly tree it both and VanBuren. It lead, linearly, lose in the tree transfer to the same tree of the same tree o

Sept. 10, 1947

Littues, and then to people und model to themse, with the time. Cand min, 55% i, him Copies also in over which have a complete and both birds have contained into the Special interpretable caimbee higher are above the laws.

•

.

Observatired vender, of.

estern linto la ma

1096-1347-2318-2523

Turco co marine beneir i

Late ' Loc tion no observations.



time. Lots of them on the telephone peles, and increase Corvailis: 02 atches one hovering over one .pot pointed into the ind. m 5, 1941 It would move he hover another spot. It was activate motivates, for a period of at about , or 4 recents. Colverlis- o. neut:: 0 neal ohr rous. Olveri; or aco ozna a o eza zione nigiliari e satisfied. chart state of the Aug 29 1944 There was a heavy concentration of these birds all through the mtn. area today. More than I have seen in one day, before. Woodburn to Salem or I noticed a great many perchea on the poles and wire Feb. 18, 1946 lines all the way. They have been pretty scarce most of the winter, that is, up to now. Dixon Creek Ridge, Corvallis, A pair are nesting in a tall snag by the Apr.14, 1946 creek. Dixon Creek Ridge, Corvallis, ONE Saw one carrying a Thamnophis about Apr.25, 1946 2 feet long: It had a hard time flying, and dropped it some distance out over the field. oison Oak Lake, Linn to \mathcal{C}_{s}^{re} several joung, or is atured three that Sept 19, 1946. there, all giving min-pitchou call note. Taw a robin may a and perch within a feet of one of them. Nothing happened. 2 miles N of Bellfountain: several seen on wire lines, etc. One spotted Jan. 21, 1947 on a fence post, and it had a dead bird in its grip. We gave chase, but it carried the body with it. After a lengthy search on the ground around this site, we found enough body and wing feathers to determine (inaccurately) that the victim was a male towhee. We found where the bird had been been first attacked, for there was just a white-spotted black wing feather, and some scattered breast frathers 20 yards away. About 40 yards away was the final catch, for here were many belly and breast feathers, and even pieces of flesh. The hawk must have hit with terrific force to rip live flesh loose. Below the perching post were more body feathers and two wing feathers. The hawk seemed to be a female. Avery Park, Corvallis Quite a few seen. Most in clean plumage, but one Mar. 11, 1947 shaggy. Also saw one carrying what looked like a mouse, up in the oaks. N Foothills of Plomosa Mtns., 9 mi SW of Bouse, Ariz: We saw one male Mar. 21. 1947 light on the side of a Sahuaro Cactus at a neat looking nest hole. As

we were too close, it did not enter the hole.

919-921-923-924-933-941-951-952-956-957-958-960-962-967-9680-974-979-980-984-990-994-995-997-499-1000-1008-101-1021-



near Beallfountain: one flusuea from top of a pole by Lewis orapecker Aug. 22, 1947 that wanted to lgich on the pole. This happened two times.

12-mile Swamp? a few singles, and I also observed a pair of them feeding March 9, 1948 together, or rather hunting, from about 30 feet and about 50' apart. In one instance, both birds headed for the same spot at the same moment, both alighting on the prey, but soon one flew from the spot, leaving the other with the prey. Sex was not aeterminea.

Poison Oak L, Linn Co. one at 30 feet over Tield, trying to buck Apr. 4, 1948 a 30 mph wina, and losing ground. It had to arop groundward frequently lest vice Klameti rake, Toper: nert in condicion line in one i July 27, 1948

clessing - note at 5 f ton erro. Reset the one poure tongt. A sie no not in grador pine at row or row. . here ago is et up on south. A joing kinds marby. The It is one. One join or inc returned to the most home, too.

Farmington Res. Site: Thying along, and did snap-roll-drop when I honked May 19, 1949 the Chevy horn. 3 times repeated on it.

June 18, 1949 Yolo By-pass, Tof Sacramento, Cal. one male being chased from tail willows by Breach Lackbirds and bullock Oricle, both of which were later seen to be carrying tood to young in their rosts.

Tomales Bay, Marin Co. 4/miles of road.

Nov. 6, 1949

Nov. 16, 1949, 8 miles N of Butte City, Calif. - about 3/mile of road

Jan. 3, 1950, Woodland-Waxwell, US99-W, Calif. observed a sparrow hawk make a pass at a recently-killed horned lark in the middle of the hiway just as we approached making the hawk veer off, and loo back over the hiway, and after we passed, it looped back around, and scooped up the horned lark and flew off with it - the hawk was first observed at least 150' away from the bird on the hiway, and there was no lost motion in moving directly for the bird, and in making the 1st pass, and second catch.

Sept. 13, 1950. Elk Grove-Logi, Calif. one starrow hawk being the sec by 8 Brever Blkbas. lov. 29, 1991 Gres Fairs, Loriel C. 1770. In the line of the first fairs of the first fair fairs. On the record of the angle of t Nov. 29, 19,1

S. I end of Take Tahoe - an irreture hawk lit on ground in pauture and was May 30,1953 mobbed by Grewer Blackbirds.

Feb. 28,1954 S. Fk, American River, near Mormon Island, Eldorad Co. Calif. Nest in cottonwood, 2 birds about.

5 Souto Co, Calif. U.S. 99 E to Water. Hoing sp. Ak. trap w/ mouse in wire case brought male ang 28, 1954 sp. Ales. in from 110 yes and 25 and 60 yes and 20'.

Carson City, Levada. Clased by robins. May 17,1255

Dec. 25,1966 Sacrand to Brease, Mole G. Califf. Chased by to as forced flighten.

2/4/56 - 3/rond mile - Tomalea Pray, aly. Aug. G 7000 and Torde iste.

nct.20,1756, T. Bacto. Colif. I Write chased by god togled a ter communication of Tellow-billed Tunins.

June 9, 1957 Marden Hiway and Elverta Road, Sacto. Co. Calif. Many seen, along with several noisy family groups, especially among cottonwoods along Sacto.R.

Novem.19/1, 1958 Sacramento-Thornton, Calif. During wind and main, several perched on wires along road or on road and held wings outspread as if exposing as much

June 17, 1960 - sope Deach State I. k, Tate Talle, Jame A. Ca, Gair, a Google Har da by a Claste C redwings,

July 28,1960, Carnelian Bay, L. Tahoe, Calif. F and young calling from hole in sine snag. June 22, 1961, Woodland Hiway, Yolo Co. Calif. 2 adults feeding ? 2/3-grown young on

willow stub, near home.
Sept.10,1961, Lassen Peak trail and summit, Shasta Co.Calif. 1 at 9300 feet on trail, and 2 over peak at 10,470 feet.

Dispute Minh. I'm some of building the Melletto on Commissed Calif.

Aug. 17, 1762, Envis Swiff Fours, Tolo Co. Calif. very Common singly and in family grows ye

Location and openiv bions

Louis i de la de la transferación de la companya della companya de

June 13, 1946 Berry Forest Camp, on M. Santiam R., below letroit. Eldon and I came upon a female Blue Grouse perched in a small t5ee, and we flushed 6 young. Perhaps there were more, but we couldn't fina them. The young were not over 6" long, but flew well, and fast, for 50 feet or so at a time.

Asiqua-Silver Creek Biage, 10 a.m. Seen frequently in the rosa, going in. Goose-Clear Lake are north of Salen. Spotted one in the branches of July 7, 1946 Dec. 24, 1946 a hazel bush, and approximate within 15-2 least or it, as lit

Apr. 12, '47 Abiqua-Milk Panch Area " several heard hooting this p.m.

still dien't fly. I die not ly roue closer.

Laclonala lorestor noise heard hooting, but on the way out, one maked ahead June 14, 1947 of the car for 10 or 15 feet and flew-jumped up on the bank peside the road where it watched us, and then non-chalantly warked our a 100 feet further, just like a tame chicken.

MacPonala Forest i ushea 6 immature biras from the eage of the roac. Mac.onala rorest adults and a young laurated from the gravel road at 8 July 12, 1947 August 7, 1947 a.m. loong were nearly it is cized, it so med.

- MacTonala Tores to flushed juite a few from the ground, and they to enter a Aug. 11, 1947 talk fir, and as I moved about would give forth a loud gutterar cackling sound, that I haven't hourd before. To berea riagetops.

Mary's Peak, Benton Co. Oregon 4 p.m. s evelas singles, also I strutting June 9, 1949 male with & female: on road. It flew with throat sacs still distended. Also a few falt-growns seen.

Aug. 20, 1950. Bumpass Hountain area, S of Lassen Peak, 8000' elev. Sierra Sooty Grouse in open area off trail above Hehen Lake - would only walk ahead of us, barely running at times - a true "fool hen".

Sct. 15, 1952 Granite Creek-Cattle Mtn. area, Madera Co. 7000' el. flush ed about 10 or 12 on ridge north of Cattle Mount ain. Red fir and Jeffrey pine with understory of Ceanothus and chinquapinand huckleberry-oak.

Camp Harvey West, Echo Lake, Eldorado Co. Calif. 1 F w/ 3 young on 9000 June 26,1954 redge N of camp.

Crass Lake, Luther Pass, Calif. played with one up bill in words by booting May 19,1955

and grunting, and retting abswers from it.

July 21, 1956 Phipps (app. Adre, added Steam Adres 198, added to the steam Adres 1 l abilt fra checlock chroat (Ejore . d l:) b. .

Sept.9-10,1961 Bluff Falls Campground, Lassen Nat. Park, Shasta Co. Calif. -3 - no fullplumaged males. Habitat of alder thickets openish underneath, with grass. As I approached they lay in open corn-lily meadow close to boulders 82" or more in height. Difficult to see them. Watched them feeding on tall grass heads. They had moved less than 100 yards in course of about 3 hrs during Such we as Eller I will be all more the self my say

telseen is got minte



number of fiel transfer and near the :

CLOTICES WIND LIGHT LIGHT

late

Location and obje v tions

June 9, 1967, Riding Mountain Nat. Park, W. Manitoba. Canachites canadens's Spruce Grouse Pair of the typical eastern form in forested know habitat between two bogs picked by GWMalaher. E. of Swanson Cr. and Whirlpool I ske. Spent 15 minutes within few feet of the pair - little understory cover but duff. Terrific display rotation by M - fanned tail, with layers of fans showing. Eye brow pockets up a inch bright red-crange. Wings drooped. Female unruffled. Fed and even dusted at roadside almost at my feet.

rich red brown



duncer of fiche unla on this pare weet

Oregon Hulfed Grouse

2-3-15-51-65-95-172-182-212-351-362-367-376-378-385-411-435-506-522-587-591-596-609-634-726-767-775-71 802-810-833-849-851-870-887-1158-1161-1178-

1179_1185_1186-Ponasa unbellus e bini

Late

Location and observ lion,

10-1-42

Pearis ferry: Or found Rufked grouse with most of Calkers from love in it and talon significant to vody, and we con become rould be a 1 med out as I have seen at reast I in the some of time of name seen to it.

Apr. 25, 1946

Dixon Creek Ridge, Corvallis, "Il of us were clustered together in one spot on the north side of the hill taking readings of temp., etc., and one of the boys spotted a grouse on a nest, and it was almost surrounded by us. One of the boys was sitting against the base of the very small fir on oprosite side to the nest, which was at the base of it. he was less than $1\frac{1}{2}$ feet from it! The bira never batted an eye, but trusted in its camoflauge. Later Dr. Gordon and I took pictures and it stayed on all the time he set up the tripod, put in the cartriage, and took the shots, less than 4 feet away! But I flushed it when I tried to get the camera less than 3 feet from it. Theore ! There were 10 eggs in the nest. The cavity was a little over an inch deep, and 6-7 inches in diameter, and lined with last fall's Big-leaf Maple leaves. Grasses and twigs made up the bulk of the nest. The nest was at the base of the small fir, on the north side, and was openly exposed from all other sides. A few short sprigs of Poison Oak were growing around the nest. One of the fellows had his Labrador retriever with him on the field trip, and it sat down within 5 feet of the nesting grouse, but never gave a sign of having smelled it. The bird was up-wind from the dog, as well. I wonder if a nesting bird like that can prevent itself from giving off scent of any kind ?????

July 17, 1946


Corvallis, Oak Creek Fidge 2 s.m. Collected a going Fulled Grause about 4 weeks old out on the riege. The rizeard and bulling with bruck-berry seeds, another, larger seed - \(\bigcup_{\text{o}}\), and some lift pieces. The crop was rilled with Cascara berries.

April 12 '47 Abiqua Camp-Milk Ranch area. heard them clearly, speed spaced similar to the "bouncing ball" notes of the Screech owl.

iec. 6, 2047 Acade Pat to Cat Office Patricia justice in the Cate of the Cate

Apr. 30, 1949 Beaver Pond, 3 mi. Tof Bruce Sta. Benton Co. arumming and along ask-willow-alder-cottonwood stream bottom along creek. Flushed 2. (IBM) 530 a.m.

6/11/67 Wy Toke Cowielen, Van Lo BC. Zale W/ Lying Deary yng,



Observer Fred venden, Jr.

rumber of field trips of ...id. obs. Voc.:

Columbian than, tailer Ground

Pediocetes phasiunellus columbianus

Late Location and observ tions

Feb. 20, 1966 - see below: 12's male with can dom't fee both and crown and tour white feethers on tail. Penale dom't barrier crown in tail.

They flush last, and are thoughd first bocom. Control on release.

This at the aberlain of the

76-

Greater Prairie Chicken - Tympanuchus cunido

Feb. 20, 1066; Chamberl din-Rid Pend Dom Amel, Tympa, Drule o Duffalo Cos. S.Dak.

12's with sharp-tails on without. Flying down from himler or and himle of fields.

They flush direct from mixed flocks and Amren Jackson one in flixed flocks the sharp-tails now are adend the chic enside as in Greek un on use of underwing American as countly a dill in flock flocks, along with know toil nethods. Since chicken body domker then observationed,

Frairie Chicken-Sharm-taile: Thouse Tybrid - about 1 of all traced by Jackson. We saw one at 1st trace site east of dissumi in S. D. Torre chicken than sharpie to "ours"

4/10/71 BeCome Konson - Fori, 1 group, Watered coming & country + Entry serge.

there is good contrast in profile in flight. Freeks Tof issuri D.



Observan Fred Ivenden, Jr.

Juge Len

1230 - 1232 - 1233 - 1395 - 2382 -

Centrocercu, die alle u

June 21, 1949 Near Twin Bridges, S. Fk. Twhbolct Fiver, Tiko Co. 2011. actit and the

2-grown young crossing gravel road in samebrush no liver ectton a citat.

June 2:, 1949 Mary's Fiver, Malvista Fanch, Fiko Cou nevaga - acult ind K myire younging gravel road in sagebrash - mayor bottom area.

Dec. 13, 1949 Pyramid Lake, Jashoe Co. Levada - 2 p. 1. 4 flushet from very sparse desert scrub along west side of laic.

in the state of th •

Merriam's Turkey

Meleagris gallopavo

- July 1, 1959, South Rim Grand Canyon, Coconino Co. Arizona. Mildred spotted two adults and 5 young in draw by abandoned dam E of Grandview Point in woodland-grass.
- July 2, 1959, North Rim, Grand Canyon N of village. Il seen in grasy meadow about 100 feet from forest of spruce and aspen,
- May 10, 1975, Garden Canyon. Tt. Wuachuca, Arizona. 705 a.m. one in road, crouched on pavement as we surprised it in the ca. It stayed right in open, permitting us to take picture are willow, with car envine running, and it hissed, growled and gurdled at us the whole time. I not out, we thereto the car to "free it", bent to pick it up and it immed up, an away, circled back ach as the road and into the woods 30-100 fe t before ding the same squatescape action;

11/11/90 Westensoland Co. Va - Frido Patomach - 3 & Stratford was have an family

4/8/8, Skylin Din 200 year Sof Uffer How Roll Va 2, for 2 and + 3 10 6 4 19 gloring

| | | · . |
|--|--|-----|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Observi irea Lvenden, Jr.

Lastern Bob - shite

3-6-18-34-67-74-75-77-80-81-90-101-112-142-747-149-150-151-156-159
161-197-212-378-384-41-610-60-411-610-637-646-666-667-613-681
682-688-6894-727-786-829-860-165-870-169-880a-887-888-890-844-895-897-401-901a-1045a-1218-1246-1437
Colinus virtinimus virtinimus (1439-2407-019).

Date

Location and object v tions

May 1, 1947

Stewart Lake, N of Corvallis 7 to 9 a.m. calling from several fields. Ore seen in an isolated Oak tree which I approached within 8 feet.

June 7, 1947

1 1 -mile Ewamp: one crossed the road ahead of the car, and fine hed and couste ahead of us at 15 mph for 15 fe t or so, before turnion (i).



Observed lieu wender, a.

Europe in Partriage


25-31-292- 293-299-328-491-609-

Pardix perdim perdim

Location and observations

tept. 1, 1949 Tule Lake, Colif. dia. 5 or 8 Chakers trans to train to the respective of the Press. They are quite take. They are trop the receptance of the respective for the respective of the

May 26, 1954 Kem Co Calif-5 mi suf Marieger along Liway



Arizona Scaled Quail Calipepla squamata pallida

Feb. 21, 1951 Caterina, Dimmit C_0 . Texas. Ran across this flock of 25-30 by accident when we stopped to give chase to a thrasher.

Chachalaca

June 20-21,1964, Santa Ala MAR, Midalgo Co. Texas. 12's and note with 3 chicker-size edgs.

Test noisy in a.m.s to 0 or so, but sometimes would a t-off calls in p.m.

,

•

× 8

1933-1935-1949-1951-1954-1956-1969-1979-1893-2000-1986-2006-2013, 2309-2326

Observer Fred Ivenden, Jr.

California Wall

und i of libert die, . .

oldburn, err Williath: Trade of the Sept. 22, 1941

ca the icac in mont to the ear. Woodburn! Saw covey of valley quail, at Woodburn Fruit Growers Sept 4, 1942 warehouse. There were two adult bards and 19 young barely halfgrown in the flock. They were foraging under a scotch broom clump.

Walla Walla, Wash. Came upon a large covey of these quail, I' + & + immatures Nov. 17, 1943 peeding on the ground under Sumae & week Sogwood. The we opproached they strated "concains" + toward of into The governor in the interior onto the low branches of impe is the se. the trunk, verliebly to eigher winter. One of them are a compared see the in Limbs of going resulting it sight wante week! I am and inch I saing teling ing.

Corvallis, Willamette, R., Gravel crusher. I saw one pair of them out in May 23, 1946 the open, on a pile of gravel, near the rock-gravel crusher. That they were doing there, I don't know. The area nearby was also quite open. They flew off to the woods when I approached.

Stewart Lake, 9 a.m. 3 very busy giving scoleing-talking rotes that July 23, 1946 sounded like a Gray ligger. They were perched as high as 30 feet up an ash tree while uoing so.

Oct. 29, 1946, Corvaluir, of aver ib. In a to be a caine maken play a pair, in a figure to the time, and in attraction the

Corval is-reprise retraction of a cover of 17 in a contraction of the first tenth of the contraction of the first tenth of the contraction of the Oct. 30, 1840 Nov. 30, 1946

Corvallis, Brook Lane, 0715 a.m.: 9 pair, all paired u were in the oaks Apr. 15, 1947 sunning and preening this a.m. beside the road, and others were just sunning. I watched for 5 minutes, and finally they discontinued their activities and left - females first.

Midale Canting river, Linn Co - 950 a.m. Il w ahead of the cal, and crossed July 313, 1947 over ahead of us at 402 m. ...h.

.ver, Park, such a pair seen, one is neet us in a maple, and giving alarm Aug. 27, 1947 notes to young manippy, and I approached.

Sept. 26, 1948: Folsom Dam Site: flushed one covey of at least 35 birds near the main ridge on the west end of the dam site. $\frac{1}{4}$ mile on I flushed a small

covey of 8 or 10 birds.

Gualala, Menaocino Co. 45& in plowed field - 3 lookouts. 8 a.m. Mar. 20, 1949 Woodbut, Oreg. - 745 a.m. Overcast low. 40 quait worked Ing our live, exocard sirect to 5 x Feb. 24, 1950 worked across Beckers garden.

Monterey, Lake in town - 45 feeding on park-like area grassland near brush Dec. 19, 1950 in close and mixed association with coots, flickers, and killdeers.

Aug. 18, 152 Bass Lake - Placerville, via Coloma, El Dorado Co. Calif. 645-745 pm. PDST. Saw 4 coveys of 2 adults each with young of 4, 5, 7 and 5 in number.

Farmington Res., Stanislaus-San Joaquin Co. Calif. saw pair adults with 12 Aug. 22, 152 young not over 1\frac{1}{2}-2 inches high. also of Hogan Res. - 10 less than 2" Li. Geysers Loop, Cloverdale-SE of Geyserville, Sonoma Co. Calif. coveys of

Apr. 19, 1954 6-20 very common. Asilomar, Pac. Grove, Monterey Co. Caif. Watched F leading male in more or less May 11, 1954

direct line for at least 300 yards thru pine woods.

Calf. Quail: 182.2

Nov. 3, 1958 - U.S. 385, near Franktown, Works I, Works Co. Nevala.

Flock of 35-40 feeding on your alogs facing & A beef him.

June 22, 1961, Bass Lake, Eldorado Co. Calif. 1 pair with 5 half-grown young

June 28, 1962 Bear 8. nouth at Westher R. St Tuba Co. Calif. few calling, and

flusted one cover of a rout 20 young only 32" bigh.

S/31/19 Regard road, Millorgan. I landed from grace & road to

near agree.

They got up to 35 each define turning of about fue.

Lophortyx gambeli gambeli

- 3-17-47, Needles Landing to Parker Dam, Calif. 14 in wash with tamarask and Palo Verde
- 3-18047, Bill Williams R. Canyon, Ariz: seemed to be small flocks at many points along the canyon. Gave one "Valley Quail call" and one of their own.
- 3-20-47 Colo. R. Indian Res. south of Parker, Ariz.: 4 at edge of arrowweed in flat country, on dirt field.
- May 14, 1952 Meadow Valley Wash, N of Caliente, Tincoln Co. Nevada pairs seen regularly along this U.S. 93 hiway.
- May 14, 1952 Pahranagat Valley, near Alamo, Lincoln Co. Nevada 1 perched at least 40 feet high near top of cottonwood. Others seen fairly high in brush, etc.
- Feb. 13, 1953 6 mi. 5W of Miami, Gila Co. Ariz. near Sleepy Hollow. 1010 a.m. Snowing, and snow on ground in this juniper-oak-yucca area, and we saw a couple of these quail sitting on rocks in the open, while it snowe.

5-6-65-211-225-401-478-512-516-587-609-634-635-641-642-649-662-672-679-686-1298-1320-1353-1385-1673-89-955-767-73-808-833-865-893-104-408-1018-1021-10402-1041-1066-1136-1149-1198-1210-Ureorty, victo and 121

Location and Jb. rv vicus

March 9, 1943 Corvallis Protes cameters by in part month your confined miling. A frankly and for your ward. Cont in we taken, Variety Guail, but this measured browner & was a to include since call notes with night and more within with the to a. Tox i very Guril.

- June 1, 1946 Yew Creek are a, along A lsea Hiway. Beautifully colored individuals seen along the way this evening.
- Oak-fir belt, 3 miles west of 12-mile Swamp: we flushed a covey of 8 or 10 Feb. 25, 1947 of them at the edge of a wheat field, and fir woods. We could not approach very close to them. One gave a short, soft, call note as they flushed. They were quite noisly in the brush.
- Dixon Creek Ridge of found a small covey in a dense briar patch on top of Mar. 1, 1947 the ridge, in logged out fir area. Call notes of alarm were sort of tremelous and exceted. Just caught a glimpse of the birds.
 - Apr. 10 '47 Dixon Creek Ridge: Flushed a pair from brush in the D. Fir clearing atop Dixon Creek ridge, and then flushed 2 more farther down the south slope in the oak area.
 - May 29, 1947 S. Spur MacLonala Forest! One was heard a re or less continuously for most of the afternoon.
 - Mills Fanch, Abiqua Creak Mushecete pair from the eage of the rold. May 31, 1947 and they flew just 10 feet.
 - June 25, 1947 Lorrang to amlauf, Louglas Co. Lt 1000 in fir lorest dieas. Came across a pair that had is cute little chicks. Wheat ly" long. I almost caught one chick. They sounded just rike a broom or chicks. The female stayed within 6 fest of me, crest erected, and gave keveral types of scolding and alarm notes all the time. Then I left, she have two plaintive notes, and had answers from the chicks, which had been miding silently till
- Maclonala Forest watched 4 g-grown young wlaking along a log. They were July 12, 1947 brownish-gray, with prominent head color bands as seen before.
- August 7, 1947 Actorate Fore to came across a covey in the road. I have to make 7 -grown young that were goite brownish in color.
- foott Vermey, Implus Cott pair with covery on 15 yours about to long. July 13, 1:48 must: very win-s onc.
- Trail Creek, Chile Dei Pail, Jack or (2. Jun 16 e.) was an Jes July 24, 1148 ar, bruss 100 fortist the ere k.
- Coolinge Managers, 6400', Freeno Co. N. Fk. Kings 1: Fout 1/1.5 acres.
 Trinity Mins for trivity Contex to Mineraville, Trivity Co. Calif. May 24, 1949 July 6, 1950
- lug 21-22, 1951 Huntington Lake, Fresno Co. Calif. 6,950' el. several small coveys were seen about the lake, but most were composed of young or immature birds.
- Sept. 8, 1951 Lassen N. P., Manzanita L, Calif. 6,000'. Covy of 8 seen, mostly immatures.
- July 11, 1959 Camp Winton, Rear River, Amador Co. Calif. adults with 10 young 4" hi. Aug. 2, 1962 Mescuits, Eldorado Co. Calif. 23-267 Tel. 3 pair, 2 with coveys of 1/6 to 1/5-grewn young.

Observal Fred Evenden, Jr.

Ituma or this taken a contract

Flumea quail

568-561-

Oreorty: pict. ict aremophila

rate

Location and place IV

March 23, 1947

Joshua Tree N. Mon., 5000' elev: heard and flushed several. They were all over the valley and rocky hillsides where we camped.

arch 24, 1947

Joshue Tree N. Mon, Calif: giving their singular notes before sup-up this a.m. I'd estimate 15 scattered around this small valley.



Late ct. 10, 1941 Corvering - 1900 Pt as to a consequence of the urch 27, 1942 white eroms. 711 Pudding R., Toodburn - crowed liter blast on nearby farm. Poticed this Mar. 14, 1948 occurrence quitefrequently this winter. Avery Purk April 19, 1948 - Escaped Lilver Placasant hen had nest on Thate of I. Fir in brush and grass. I ling-neck erg and 8 of her own work in the next. Airil 30, 1,48 Fing-Leck egg hatching, but not the liever Pheasant eggs. la-mili 'ar jer'ilu 'ng. Lir dailte and dat the lear to the se-5" with. June 18, 194 Aver, Park, Corvolli. Of tilver Shouse, to Texture or the most park law south June 20, 1948 of previous parties, or a del believe as mathias defit of a decision broken by the Co. I Witches on a ten? ?? She has but neems to be we the most, for it was ol, and I with a six how bering the clark the attempt low, gutteral of a treatment the file. There will be a star Windowst. The 11/50 nest as appropried to I to a other terms till into the Sept. 25, 1948 Grey Lodge Refuge: 100's present, in areas near flooded ponos, as well as in dry fields, of pasture, and of grain. Mostly immature birds, comparatively few adults seen. In the a.m. they were gathering grit from the gravel roads, and as the day warmed we would see them on the roads only in the shelter of willows, etc., or in the shade along fence rows, not out in the open. Jack Rabbits often seen with them. March 12, 1949 Lake Washington, Yolo Co, W of Sacramento: flushed 50 pheasants from boggy burned-over cattail area at edge of lake - alfalfa fields nearby, but none seen in them. Yankee Hill, Dutte Co. 1815. oak woodland - 2 cocks in small meadow area Mar. 30. 1949 near creek. Surprising to/see them here at elevation of about 1400 feet. Lake Washington, Yolo Co: 7530 a.m. hen on nest in "broad-leaf-mottled-thistle" Apr. 30, 1949 area, which had been trampled down by a truck, so the nest was fully exposed to the south, whence I viewed it. The first time I came within 3 fect and she dian't flush, but when I came in with a camera to 3' she flushed. I took apic of 12 eggs, very fresh-looking. Take tashington and vicinity, tole to. On it. The new late, that evently Aug. 12, 1949 acre in italization area sear chracku-citch-and sed-cite tele. The inter-Gray Louge Defuge, Public Co. Gallif. - Ethough per a late una a poste overann-Aug. 16, 1949 ning the refuse, few who seem this atte, is the seem to be likely by hole they are no ting t trie time.

Nov. 15, 1949 Williams, Colusa Co. 4 several pheasants in the rice. Nov. 16, 1949 Orland Irrig. area, 6, 3, and 3 seen, 3 were perched in cottonwood at about 15' elevation. These were in Black Butte Res. area. In the irrig. area were 7, 6 in alfalfa field, and 1 in willow, about 15' up. Sept. 20, 1950. Will of Moodland, Yolo Co.Calif. one glided with car at 20 mpl. for about 200 yards, and just after being flushed from ground near road. Sept. 7, 1951 Lake Washington, Yolo Co. Calif. at 825 a.m. - many seen (35-40) Most in ragged plumage-immatures. 2 in one willow and 2 in another willow at heights of about 10-12 feet. Aug. 22, 1952 Farmington Res area, Stanislaus Co. Calif. several 3/4-grown young. Rio Viata + Dais, Calif - 12 sel you plusests clong the roads flow in rain shower 2-3



Little From trane

329-461-468-580-2241-

Grus comadarsis caracturis

l hocation with old day differ

8-4-44

Leen at Weiser, Huntington + Durker, Italo

Nov. 19, 1946 Corvallic, "Dais in Dan" le. 04

linshed an line to our it the landers to I con it van rolling crown as and the pine on love I to.

Apr. 5, 1947

Leadbetters Penin., Willamette-Columbia Sloughs: we saw one in flight over the tops of the cottonwoods. Almost the size of a Blue Heren, and elongate neck was well observed by both Dave and I.

6/21/66

St. Lucie Club, st. Lucie dr. vie. 2 clults of countries the tentre and 10% and create and c



§ Th.

Observer Free Lychden, Jr.

Eanchil Gran=2080-2081-2241-2242-2243-2276-2286-2428-2447-2493-5-7-

Grus canadonsis tomico

Late

Location and observ sing

Oct. 4, 1942

Corvallis, Duck Trap Pond. Or Foggy in spots and cold. One flew over me at about 15 feet. Seemed to have a wing spread of 7 or 8 feet, tip to tipq Neck was not bent, but was straight. Near me it banked its course and dipped downward, very deftly for its size.

Nov. 15, 1948

Los Banos State Refuge: Wandell Miller and I heard some several miles away, and when searching thru the glass picked up a flock of 40 or so circling for a landing in the area near the San Jaaquin R., E of the refuge. Altogether there were at least 250 cranes in the field.

7eb. 3, 1949

Secremento 5872 Raymond Way - 73° q.m. 4 flying mortherly in big loop a mile across.

April 14, 1945

Philomath, Benton Co. Oregon (IBM). about 100 of them circling at a considerable distance up in the air currents. Gave coding sound like a lot of pigeons. Fig not flap their wings. Could see long necks and legs. Ends of wings carker colored. They stayed within our view for several minutes. They constantly circlea, but were headed in a northerry direction.

Apr. 5,1949

Corvallis - 1230 noon. flock of 8 rlew in at several hunarea feet elev. from the S and processed to circle over the west end of the DSC compus. This circling consisted of alternate flapping and glidings as the birds rose steauily nigher. After 10 minutes they were discernable only with a pair of \$7% binocs. They then formed a small V and proceeded straight noth. Apparently they had struck a convection current, for they rose upward with comparative east. At no time did the birds give any call. Thick

Nov. 22, 1949

Gray Louge Refuge, But to Co. Calif. 2 p.m. - a fairly compact flock of 36 individuals were observed feeding on dry ground in the center of the refuge, but just about 150 yards from the car. We drove within 75 yards of them, and then they took flight when John Sypulski tried to get even closer for a graphlex picture of them. They took flight, and gave their rolling call note as they did so. They were slow, but powerful fliers, and circled nearly back once, before heading off toward the west end of the refuge. Two

11-30-49 2-12-52 separated and headed southeast, not joining the main flock again.

W + S of Mercad Coll 100 1/4 5

Les With 185-200 in 185-200 in m. John Engle.

Atta on a constitution of the first flow of 1,000 days.

Feb. 20, 1952

Canal Ranch, S of Thornton, S. Joaquin Co. Calif. Saw about 1200 in one field this a.m. Quite a sight. Observed some portions of courtship dances similar to the Sarus Crane in India.

Mar. 27, 1952

Hogan Res., Calaveras Co. Calif. Small flocks flying north today. Bass Fake, and I's, aldorade Co. Calif. 35 to dire up comb fort of Jan. 10,1954 american liver - watched for a siles to

March 21, 1954 Jenny Lind Calenna G. Gif. 1 p.m. 21 insoling liver rea.

Feb. 15, 1959 Cond Rock Rd, 5 W. I thouten 5 bagin to Calif. 3, Cond to a see, Into Gill, 2 of the season on grand to a see

March 21, 1954

the state of the s entropy of the second of the s

Virgini (ii) 1011 659-661-67-639-639-639-655977-78-142-147-210-275-384-396-5543-583-583-612-616-67-639-639-655977-957-768-984-1011-1043-100-1156-11472-1187-1309-1375-13262-1457-15816-1581-1057-1796-2099-2101-2121-2132-21742266-2447-2495RELIUS Limicole Linicole

inte May 8,1939 May 10, 1941 Top of bedy brown in the control of the property of the standing reader for suggest, and with his or united to the standing reader for suggest, but level up the control of the control of the property of the property of the control of the control

May 18, 1941

Lemmine (valp acts of arowali, \mathcal{P}), if we have a fixed the name to tampo to tata to the second and the second to the

May 24, 1841

Lty 5, 1942

che 8 apro mante e tembo.

1 -mile state a transferie: Thompicture consult our, with spots receive a consultation of anish and the state of the state of the spots of the state of the spots of the state of the spots of the state of the stat

March 6, 1945

gradician in the colored solution of tween the bringer.

Apr. 24. 1946

McFadden 12-mile Swamp clear and hot. Saw three of them today, and found two nests. 10 eggs in one, and 11 in the other one.

May 22, 1946

12-mile Swamp. Of Saw just one today, and it flushed from just between my feett, almost.

Apr. 8, 1947

12-mile Swamp: one heard on the east side of the swamp, and I later flushed one from a clump of Carex sp. at the north end, near the fence. I had only a grimpse of this quite wary bird.

Ar. 15, 1947

12 mile Swamp': Tom McCallister flushed one this week, not many yards from where we flushed one last week.

Apr. 22, 1947

12-mile Swamp, 1530PST: first saw this species on April 8th, and located at least 2 pair today. Walking thru sedge-meadow and found ourselves "Surrounced" by a female (?). We stood still and it worked its way completely around Tom and I, never being farther out than 15 feet from us. All the time it uttered one particular call note of alarm at the rate of about 30/minute. The sound began with a barely audible sound like that of a quite taut rubber bana (med. weight) being snapped, and then the clearly audible part consisted of a sound not unlike that produced by us when we hold a blade of grass between our hands and "squawk or squeek" it. The birds note varied some. While the one bird was working its way around us thru the spike-rush and sedge, a second bird remained almost within a 3 foot radius some 20 feet away, and gave the same alarm note, and also another, deeper, "gargling" or grunting series of sounds, and these were occasionally taken up by the female, as well (assuming the sex.). Finally the F came into a sedge clump 6 feet away and stop ed scolding, and quieted down long enough for a "breather" and then we heard it "purr", or at least the sound suggested that to both of us, tho it was barely audible. Then I spotted a small black young swimming thru the spike-rush toward the parent, evidently it had remained "frozen" until the "purring" signal from the parent. The young was $2\frac{1}{3}$ " long, the tarsus was la" long, and blue-black. Iris brownish-black. Skin pinkish-white. Bouy covered with a dense, long, glossy black down, the cutest little dickens you ever saw. The bill was pinkish, white at the base, with a black band around it, not quite 2-way out. The "egg nail" was still prominent on the upper tip of the bill. One item I forget to mention was the well-developed " claw" at the thumb of each wing Both wings down-covered.



- May 6, 1947

 12-mile Swamp, "cornered" another pair of them, and Rurá a small downy black young about the size of the one caught on April 22, 200-250 feet away. Banded with size #3, 41-354647. Both parents scolded and "chipped", their mandibles together while we banded the young. Black ring on bill, but no "egg hook" present. At first it tried to swim away, and then as I came close it "froze" among the spike-rush with head low to the water.
- May 20, 1947 | 12-mile Swamp? in the south part, an adult on nest containing one egg which was a cream color, spotted with a maroon-brown. Nest od ary Carex stems 7" above water. Both adults stayed within 5 feet while I was at the nest.
- May 27, 1947 12-miles Swamp. locat c another pair in the Faming Swamp, which had young up to 3 or 4 inches in size. Both parents stayed caree of Coulon't catch the young.
- June 17, 1947 | 12-mile Swamp. heard them several times, and saw one may some 75 neet from an "open" area of Impeochamis to a Carek obsupta chump.
- Aug. 29, 1947

 Term Fiage Lam, Lame Co. Tob and I caught a brack county bard in a remy area near the lame bord or north of the day. It all to be to be an about a large The with head c/4", and bits bards and all long, and tearth at out being out being and the remy filmiest and smallest of rectains where coming out being an item, and the tips of a fet in the rest where to be seen. The interpretable already had touches of a strutt around it. It run under a hor carrie we caught it, and I benede it with 41-354648. In very material record is indicated from this pira!
 - Jan. 6, 1948 12-mile Swamp Pave M. and Tom McC. found one along the gravel road leading into the swamp area, in a flooded area about 100 feet east of the normal margin of the swamp.
- April 20, 1948

 Ik-mike Ctamp: seem to be quite a flw in the area. I'd estimate at meant 10 pairs, and I'm sume the dian't hear or see them and. In the earlin, in week still in same condition. Found another nest, near the earlin, in Carex obnupta. Clant material still wet. About 8" over eater. Cup 8" aleep and 4-5" in diameter. At I corner we found a grass platform about 5 fest up in shall ach, in Carex clurp, which we can't right out, unless it is a base for a nest of this species. Another nest platform made from Carex 2" above water in Carex clump. Tater depth at nest, was about 6".
- June 13, 1948

 The mile than post of a contract the post of the results of the post of the post of the results of the post of the results of
- June 21, 1948 | 12-mile Suca : Tourn & pairs and a single abult; also "peopers" with one pair, and to single abult.
- Sept. 8, 1948 Sacramento Refuge: calling occasionally. One found freshly dead along the road. Near adult plumage. Base of lower bill deeply notched or widespread.
- Feb. 2, 1950, 12-mile Swamp, Benton Co. Oregon. Found this one standing under a Rosa next to the horse shed. I approached within 5 feet and took pictures. Then hit the bird over the head with a stick, stunning it so Merl could pick it up. Was very, very thin, with absolutely no body fat. In gizzard were a few small fish bones and what might have been some skin from a fish. Undoubtedly this bird would not have lasted much longer. DBM
 - July 30,1952 Near Eagleville, Modoc Co. Calif. Flushed I from flooded, oiled pasture. Oct 10, 1960 Canachael, Subless Longlet 1951 forms one or more



DE IV E LEE VIL , J. .

Soru Rail

1043-1147a-1187-1903-1948-2108-2147-

142-147-152-288-396-398-595-602-621-661-695-710-868-880-887-888-968-984.

Porzena cerelina

Location fin (Danger vite)

li-malar usa j vouta de somelais fe e de e i granda. Jun Dille, the crist its in the bound of the constant of the const

lay 5, 1942

12-mile Swamp. Or Saw a couple of them this p.m. May 22, 1946

12-mile Swamp. The heard a strange series of 3 notes, that may have been Apr. 15, 1947 this species, but of course none of us are banking on it.

12-mile Swamp, checked up on last week's call and this is it. Tom Apr. 22, 1947 coaxed one up to within 10 feet of us, by repeating the "ker-wee" notes. Heard just the one pair today.

Aug. 15, 1947 "Poison Cal" Lake, birm Cofflushed one wire incl. of the grangin among the pond lilies on the Co-slae. It flow without a scalar to at 60 % lest before alighing in other pend dilies.

Milwaulere Oregon - 11 eggs in nest of dead sedge grass of in a clay of nedge . cuttons.

1. " - 10 eggs in " " seage grass surported in a Competendar prose level with

The water . The nest was covered w/ leave grass to forma doned . roof. - a "trees. Man 11, 1940 May 17, 1941


4

Coturnicops noveboracensis2

Oct. 2, 1948. Suisun State Refuge, Solano Co: Gordon Gullion and I heard a new call from the tules, and checked the call descriptions in Peterson's. This species is the one that fits it exactly! We did not get to see any of them because of the dense Scirpus growth. However, I would say there were at least 10 or so in the area.

1/20/19

And the action of the




3-16-47 Monolith to Mojave, Calif: wading in shallow water nottoo far from the hiway Sept. 8, 1948 Sacramento Refuge: purple black and gray color with redding feet and bill.

Dec. 18, 1952 - Hood-Tranklin road, facto 6, Caly 8a.m. faw 25+ in one ford of tall cattails typen water thanks.

Jan. 10, 195h C.S.H. 17 - 2.h ni. 3 of Julish Sta. 2 n. . . Frat 15-30 10 11 houles in shallow water area grown to cattail 3, in cattle carrier.

Jan. 12, 195h Same local - only 14 seem.



| 1996 | 1912 - 1918 | 1920 | 122 3 - 123 6 | 123 - 123 6 | 124 9 | 1295 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125 | 125

seemed to be scattered out over the swamp now, rather than in large flocks as before.

cct. 7, 1947

Li -mile - E. M. Lindman of the column of

Oct. 18, 1847 | li-mage stamp: Oct. denote , il not salpo att. for which we can also the first total of the library visit of the first we rubners to 1890.

Jan. 20, 1948

12-mile Swamp: watched 2 diff. ones take off - runs of 45' and 50'

12-mile Swamp: 1015 p.m.: Heard no owls so I agree the resection. Out call. Got an answer armost immediately from the role. Burning a 10-minute period I reported this home times.

The carl started out ammiousl as a ferecel Jar, but finds of a series of "keeyou" s. At first the, were as roug and first at the "warpar" nouse, but became progressively shower, and lower in liter, the last "keeyou" being barely audible at a distance of 15. or so.

linea it up and went out to it - round a floating Goot nest. Made or camp and soggy rotten seage, etc. There were 7 tannish eggs, very heavily spotted with airt, in the nest. Next cavity very wet. The eggs were compactely covered by a hancful of soggy nuterial similar to that of which the nest is constructed, so that the whole arrain loomed like just a pile of rotting debris. Took a jics of them.

June 21, 1948 17-mile S amp? The coots were not some ties ties.

July 16, 1946 Fern lacge Ferenvoir: "plenty from the product colling. Very misy groups.

Sept. 4, 1948 Westlake fark, Los Angeles: several - adults and feeding one downy young about 4" long. Its bill orangish. Another - near adult size young, gray plumage and with gray bill.

Oct. 23, 1948 Grey Lodge Refuge, Calif: 10-15,000 of them!

July 7, 1949 -12-mile Luxung - no coots.

July 31, 1949 Grace Lake, Ochit. adults and semi-downy young, ith or restrict. Aug. 2, 1949 hencotd Pool, Ochit. adults and secure young (b) loop).

Sept. 1, 1949 Lower magnetic and Tule Laker, Count. some reen. I with Labelier. ...

June 24, 1950 Monterey, Calif. 10 a.m. On the lakes were your, from count to grow a Green legs. Observed some cults in antagonistic pase with head lowered and wings raised over back.

Aug.19-20,1950. Mannanita & Reflection Lokes, Lassen Park. Total of 15. Also about 7 5/4 grown young, with wark legs. Also, 5 downy black young, with bright orange-red heads. Tatched one acult coot single out one mallare, of 7 perched on the lake bank, and chase it into the water. 3 young, each of diff. size.

May 7, 1951 Southside Park, Sacto. Calif. 1 carrying an 8" long, 3/8" dia. twig across



Sept. 8-9, 1951, Manzanita and Reflection Lakes, Lassen N. P. Calif. 6,000' elev. 20-25 on Manzanita Lake and 6 or 8 on Reflection. Several immatures, and a few downy young with red faces and bills seen, too.

a few downy young with red faces and bills seen, too.

Feb. 28, 1952 - 2,500-3,000 eating-out grain field north of forestry bldgs. in Yolo Co. Apr. 18, 1952 - checked field, but damage not evident. Much of crop headed out, except for plants in swale areas where water stood - perhaps coots eating only

delayed heading of plants.

June 3, 1952 - Leetsville, Churchill Col Nevada - 3, 1/3-grown young.


June 5, 1952 - Carson Valley, Douglas Co. Nevada - Class II - 4 and 1.

Dec. 23 1952 - Late N.W. Refuge Coluc & Calf. Flucked some + they found 35 might.

Feb. 3, 1953 Morro Bay, Calif. 100's around bay, and swimming onto bay waters with brant,

and feeding with curlews and redwits

Spay 199 Nogette Level Cand. One cost of Loover wings.



Bluck Oysvui-cutcher 25/1-2697-3564&&

UK: VI 12 THE THE TALL.

Haemitopus bleam ni

Late Location are observ time ch 30, 1941 worth end of Aguth Leach: I have one pair of and, patched with a carma Gull. They were 1/7 to be the circums with the circums only and basek. United the suggest in the soft and the circumstances. seemed to be a or a inches were write the line to a larger to color. In Thight they compared to have a singular or a sample et. Boiler Bay State Park . We spotted two of them feeding on a wave-washed March 10, 1946 rocky shelf at the base of the cliff. Took pics with 12" lens, movie. Coast F of Carmel, Tonterey Co: up to 6 in a floor on offs one rocks. May 21, 1949 Apr. 18, 1954 Ft. Bragg-Westport, Mendocino Co. Calif. 5 seen on one rock - adults. July 5, 1970, Depoe Bay, Oregon. 2 pr with nests at Surf Point. Those feeding on low-tide pools on rocks had a buzzy call

·

Observed transvenor, of.

estern : nowy lover

70-441-1039-1305-1307-1686-1948.2187-2254-2422-2583-

Onaraciia, mivosum mivosus

| Luce | Location in order cipus |
|---------------|---|
| Aug. 1 , 1000 | nernely finite, Lyunger, every - 15-communication - 100 mm. |
| Aur. : , 1949 | Action inc. 1991 |



Observer bred Lvenden, Jr.

Femipalimeted Ploves

182-113-199-991-1299-1567-1578-15873-1/953-7187-2329-2421-2521-

Characrive semi a Lastur

Tute Location and observations

pt 26, 1942

Wecoma Beach: 8:50 a.m. Foggy and misty. Came upon two or three of them feeding in the sand at the edge of a tide pool. Upon investigating near where they had pecked, I found the sand schools.

June 30, 1970 SW of Laramie, Wyo. was a fast runner on the ground. Several types of notes

6 6


2507 - 3553

Mountain Plover

Euroda mentana

Feb. 16, 1959 - El Centro Road, Sacto. Co. Calif. RW-, SW 15-25 Brkn.
2 with Killdeer, Dünlins, and W. Sandpipers on very wet muddy field.

June 30, 1970 SW of Larmie, Wyo. Fast runner on ground. Several types of notes. Sept. 00, 1970, valid the second of the second o



oison dan rake Maine d. reveras rais locar d bearinging arouse Sept 19, 1946 to use use chiti. Addin seen, and 1.0v. ..b. 1546 alocks, fred grave in alint. May 6, 1947 | 12-miles Swam, or two different pair seen that seemed to have nots that we couldn't locate. Mark 22, 1948 College Farm, W. of Corvallie de Theo. Went reported a next containing Yangs on this date on a point of land between 2 creeks, just above this juntim fix. Open site few Lards, even. Government Island, tol. i, new Postiand - round adult blire recolding and mpr. 18, 1940 feigning. Then Ton AcC. iting a young one lying alat, noti have and elacamoflauged in the rod. Infter considers to tooking as I no the lare joung ones in the same manner. A ratithin 8' radius. Tarkou and alcohou 4 young ones showly make their way to bleat theathers of whilt. Yellor tip on one if bil..

12-mile Swamp, (LBM) saw one by the horse shed. Seemed very wear and the sheet of ice on it Jan. 20, 1949 reluctant to fly. lich't call. (most ground has sheet of ice on it now) Parate Res., Mokelumne R., C laveras Co. nest on open south slope next April 2, 1949 to 12" chunk of quartz that was flush with the ground. The nest hollow is 37" across and about 1" deep. Failing Estate, Benton Co. Oreg. DBW - 200 in fallow field along 12-3-49 entrance read. Lever saw so many billdeer at one time. Paw flock of 100 in rye-crass field near the seose flock, later. Bass Lake, 2000', El Dorado Co. Calif. 1309, PDST. Watched an adult, just April 29, 1951 as drops of rain came, and adult squatted in open, giving notes, and watching us. Soon a 3"-high young came and got under wing, from behing, with head facing forward. A minute later we noted one young of same size about 15-20 feet away. Finally heavy hail came, hitting the adult bird, without bothering her, tho some pellets were about 3/8" diameter. Finally adult acting quickly got up and left, leaving 2 young unsheltered from pellets. They kept trying to pick up the pellets bouncing around them, and finally the adult which had left to seek the other 2 young got all of them into a dense, 6" stand of Erodium near the lake, where more protection was available from the hail. After hail passed, and rain tapered, the 4 young and the adult were out feeding again, then heavier rain came again, and they assembled (3 of them) under adult on spot first observed usea. One young got under from front, facing back, this time. I gave chase with a dip net, catching one young, anchanding it with a 2-band, 40-250426. Sacrano to imposs: Dec. 26, 195h

the marsh near Richardson's and continued in a southeasterly direction away.

will - wrang at Karst 12- 2 who Construment

Britishows, sitto to Curit - 3 your down formed this arm.

Ups. 26, 1959 Uf-7-6-2

May 30, 1958

Willdeer, p. 2.
um. 19, 1962, Davis Sower olds Tolm Goldalif. few, but 190 flocked at Mondland Sewer Fa

2.5

ollan Ploves. Feb. 15, 1959. Justo G. Laly .. ice reco f 2/10 09 holow, 2504. Pluvialis dominica Observer Fred G, Evenden, Jr. Black - be lied

Unland Plover

Number of field trips on which observed: Jenna Plover 250-257-309-567 491-1104-1147-1265-1291-1317-1330-1334-Jenatarola squatarola 1351-1362-1381-1386-1567-15526-1567a-1630-1683-1686-1699-1756-1757-1786-1854-300 tremia longi estuda 2427-2571-2583-2209-2239-2242-2243-2254-2265-2295-2143-2396-9-2403-284 Bertramia longicauda 2717 -257/-2573-

Date

Location and observations:

Oct. 2, 1943

McChord Field, Washington. Ford Dicks and I observed tow different shore birds, one description follows. It is the first Upland Plover that I had ever seen. It was large and rather cark brown above, lighter below, had short bill and legs. They appeared to be partly webbed. Legs were black. Neck rather long, slim but stout, and bill-to-tail length might be 12". In flight it dipped and veered from side to side. Body looked short and wings long and tapered when in flight. It emitted a scall note series or squawks.

May 17, 1944

Mtn. Home, Idaho. Reservoir. Saw one pair of them. About 10-12" long Black face, throat, neck, chest, and belly. White bordering it above on head and neck, and rump. Back of the neck and back and wings is a speckled or mottled gray and white. Wings mostly white in flight. Black bill and dark legs. It has the trait of the killdeer in running a short ways and stopping.

Dumbarton Briage Lower San Fran Ba; 1100 PST - saw just one, about 12" Mar. 27, 1947 booy length. In winter plumage yet.

Oct//13//1950///FAYALIAh/141/hds.

Moss Landing, Mont. Co. Calif. Several, but no full blacks.

Apr. 2, 1955

Noss landing - some full blacks
Rey Farm Island, Marode Co. Calif. Pars. Apr. 4, 1,55 Apr. 4, 1955

P.T. Pleasant Rd, Ef Franklin Rd. Souts to. Calif. Ocal 150 in writer Feb. 15, 1859 not over 1214" dight. I golden Thomas of Willow is



Coler r ir a stenon, &.

164-182-776-1061-1410-1535-1536-1901-1904 1956-2254-2268-2274-2344-2422-2642

2687- Suri-bire

aphriza ving.

nug. 1, 11d:

1.c. 0, 1847

Dec. 18, 1949

Rewport-waldport saw one with block turnstones and aleutian sandpier kokkyken - about C. birds in the flock.

8 C Ku 9

`

Ougervor and liverent, Ja.

Lucky Turn of ..

70-204-244-628-1057-1854-2274-2306-2982-

report include almost

Jeb. 21, 1943 Waldfort to Heate Had of werry ming ange fracion of these noise, 4 fold mine . They way of me to min, man any close to the edge of the whether whether it is not one one.

- 101. 7077, 2001. - 711, 1010. - 711, 1010. - 711.

10-1/8-164-441-568-776-178-40-1039-1068-1330-1332-1384-1386-1410-1535-1761-1761-1903-1904-1903-1904-1268-2267-2311-2344-2387-2422-422-2511-2571-268-2687-

Arenaria malanoce hara

| rute ' | Location six such to tions |
|---------------|---|
| lec. 6. 1047 | |
| | |
| Nov. 6, 1949 | Bodega Bay, Sonoma County, Calif. 2 p.m. several observed with other |
| Dec. 7, 1949 | shorebirds in a marsh at the east end of the bay. Morro Bay, San Luis Obispo of about 15 of these birds were observed squatted down on piles of seawerd above the high |
| Dec. 18, 19 | wave-wash of this clain, sands beach - no rech perched for them. |
| | |
| | Seal Rocks - saw flock of about 10, and Tep allows hear approach - Db., |
| Oct. 15, 1950 | Farallon Isalands: several of ervec on the wave-parage rocky store, and |
| | also saw 4 or 5 about 100 up island cultis and slends on very dry areas. |
| 0-1 70 7050 | |

Oct. 10,1959 Bay Farm Island, at Bridge to Yadeda, Calif. 1145 DAT. Clear and warm. with 10,000's of shorebir's with a dozen or so black turnstones on mudflats.



1/10/14 ac god, Colonday pid - son force + 1 miles of the force of colon of the grant of the gra



Observer bree lyoneen, Jr.

The first and the order of the second of the

Corvanuis: De Land the particle at the particle and the contractions of the them Jun. 5, 1541 twisted its flight to are not by the pain is also seen a let by and the ground no just a not let by the ten over the joils, in the notelection to have entry the colors

12-mile Swamp south of Corvallis ? This p.m. I flushed them by the dozens. April 24, 1946 They were everywhere I walked.

Poison Dak Lake, inn Co. two of them in the outgoing at the lake edge. Sept 19. 1946

12-mile Swamp: perhaps 25 or 30 were flushed from flooded pasture at the Apr. 15, 1947 north end of the swamp. In beautiful plumage now.

12-mile Swamp: quite a few in the flooded pasture area. Heard several April 22, 1947 "winnowing".

12-mile Swamp: heard one or two, sur oscaly winnowing slong the east June 7, 1947 side center area.

Fern Linge , and a pair of them taked from taked on aleas need to lowe, July 24, 1947 and they cite ou are no.

12-mile Swamp: a few in mershy areas around the cam, and 50-60 Plushed March 30, 1948 in north fiera.

1: -mile Fram :0730-045 a.m. at rea t 5 in order. Tooled by trinking we heard May 15, 1948 a foreed, own for a smile, but rocated i, a or a piras sciling as the high above us, and sorecline: juite low over the ream, . Throwing to highway high only, however. Ton a muce with the vinga, which are nouttered quate rapidly as the swoop with a slight drop in altitude. If whiching takes place while there is a desper sho p, the tones become richer, hore mellow, and lower.

June 15, 1848 | 18-mile Carpet dr I active to real I of the distribution the The parent and high in a fine to the base of the fine in the contract of the fine to the fine the fine that the fine the fine to the fine the fine that the fine the fine the fine the fine the fine that the fine it rank a noter; "the beauty of the beat wis noth; but, the work morning, let with most order ones or, in the angles to se, as first as earlies, and Lamilian Vincepaul, one taken bithings aliminating, its constant the site. The ficker on two on the Time ten occurs him to be an mest, or yours bines.

Jan. 17,1949 Fair grounds, Sacramento, Calif. 5 p.m. - 3 of them.

Apr. 30, 1949 12-mile Swamp (LBM): Thushed a few here and there - 1 perched on a fence post. One winnowing at & p.m.

Oct. 22, 1952 Mendota Pool, Fresno Co. Calif. flushed of them about noon. They were in such a rush in circling around me that I heard and saw that several of them clashed wings in banking and turning.



16_1797-1811-1853-1888-1953-1966-2024-2033-2049-2068-2022-2189-2209-2242-2243-2266-2363-2301-2312-2343-2371-23944-2396-2420-2420-2420-2420-2263-2556-2571-2591-2633-2659-62-63-

Numenius americanus

- Nov. 13, 1948, Palm Beach SW of Watsonville, Santa Cruz Co: 4 in marsh area among the sand dunes.
- Nov. 15, 1948, West-side Grasslands from Los Banos Refuge to Gustine 75-150 in short-grass marsh areas along the Santa Fe Grade.
- March 12, 1949 N. of Franklin, Sacramento Co. Yover 75 feeding in wet pasture with steers, first thought it was a big flock of pheasants, they looked so big at a distance.

Dec. 6, 1949 5 miles W of Tracy, Jan Joaquin G, Colf. 915a.m. 35 in alfala field.

Jan. 11, 1952 11 of Tracking with field for the following of area + 3-400 in grasslands E of MWMA.

Mar. 31, 1953 - 2pm. at let. C.S.H. 33 + 152 wol for Banos Calif. 5-600 in flock in reling grasses Wille.

Teb. 1, 1958 - Mining in a - 2 - Committee of the state o

| | • | |
|----|---|---|
| | | |
| | | |
| | | |
| ъ. | | • |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | , | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| - | | |
| | | |
| | | |
| | | |
| | | |

Huaconian Curlew Phacopus huasonicus

- August o, 1947 Fern Fiuge is am: one flew over ab ut lob years from Fig. 1 that and I, and it give its rather melodius "cm-lew" call an it is econouth part us. The bill was not outstandingly long, not oring over 4" at all, and the officerall size of the bird was not marke moust to surrout our calling it a home-billed Current.
 - Mar. 28, 1950. Rocky cliffs, Pacific Ocean, San Diego Calif. 430 p.m. saw 4 curlews first taken to be this species, "bedded down" on rocks just above the wave washes. The head striping was plain as in the Hudsonians, but the 3-4" long bill had a more pronounced bend than I have ever noticed on the Hudsonian.
 - Jan. 13, 1953 U.S. 101, Ventura-S. Barbara, Calif. many perched at hiway edge along coast.

Feb. 15, 1959 - Fraklin Rd, 5 & mi 3/ / mallin, Cly. 3 w/ 35 40 long- Coll curleurs.

Limosa fedoa

October 30, 1948. Bayshore Drive, Richmond-Oakland, 930 a.m. along main hiway with the willets. Brown-speckeed bodies. Bill 3&" long, and appeared slightly up-turned. Ti -half is black, other part is light-colored. 15 or 20 seen.

Aug. 6,1955 Fore Beach, Mouth Upon Truckee 1, Eldorade Co. G. if. 815 a.m.
3 on bound done of erect. The property of the control of the contr I : C: 1 _ (1)

eriotics and a second s

And the second of the second o

··

Spotts: Jungi, a

Location in obtain

May 17, 1941

Typh Valley, hite kivis; is saludi and the saludi.

Barlow, Oregon - 4 eggs in a depression on sandy least of river cutteres.

May 26, 1940

January 12, 1943 Willamette River, Corvallis, Org. Clear & Ged. 5-32 p.m.

Cal would solitary first that we a gray since, this below and had longitudinal white lines on wings, when in flight. When it lit it booked upe + down. Had black bill & feet.

May 13, 1947

Stewart Lake - one perched on the pond rily leaves near the pond center. It flew giving its cal notes.

May 24, 1947

Cantiam River, Foster, Linn Co. a pair on a par above the bringe. Had a nest that we could not locate.

June 25, 1947

Fern Riage Iam area - reveral pairs were seen. One pair had nort on · exposed gravel bar below the dam.

uct. 20, 1947

. i liametto I., orv. Line inlametto I., orvaniosit, ... joir raji nobel a o rad min, o den and com, chair, and raji nobel in countries.

[Goast I am, Tillar tt. 1, or Organia! the riving a communication of the riving and the countries.

June 23, 1948

modey, let con rever be he settle encore and I force a gitte light colored, and ish inciving a symmetries as the river to unather a v L

July 21, 1848 [id. chotto r. com reconsity "U-Mac p. .) on prover a first region of the release to the complete of the release to the complete of the complete to the complete of the comple at least in

July 80, 1948 T. Ampara liver to lyrth Paint: a ne aquit et yea or or a a te rocks on both sides of the liver. The Mac-years in it, and it meet, which ham been to grant factorial to be early when a late

June 22, 1945

mary's liver, Malvista Fanch, siko Co. Tevada - on gravel bars. The observed to fee on Cicacio, and by time it resinct the mean of the car via part, to kill it, I guess. Cicario were very con on on villow in the real

July 50, 1950 Aug 14-5, 1951

Silver Lake, Amador Co. Calif. 7,200 el. about le pairs of adults seen, and on one stretch of lakeshore we found a brook of 4 downy young running around. 1 was caught Sat. (the 4th) p.m. and banded, but the others escaped. We returned to the same stretch of lakeshore, and cornering the family on a rocky peninsula, Ard and I caught all 4 of the young, releasing the one banded the day before. Size 1A bands used were: 41-152117, 122, 123, and 125. Young were downy, gray color, with darker stripe down middle of back. I young escaped us by taking to the water, and as I reached for it, it definitely "duck-dived" and swam a few inches under water, using its wings for propulsion. It repeated this feat three times in a row, but only for short distances. Then tt rode in the water like a phalarope, and swam a full 40' across a small bay off the lake.

May 14, 1952

Upper Pahranagat Lake, along US 93, south of Alamo, Lincoln Co. Nevada. Imler and I observed one perched on a barbed wire fence about 10 o r 12 inches over the water. Saw 12 others at the lakes in this valley.



Page 2. June 19-20, 1954, Silver Lake, Amador Co. Calif. 7200' el. 2 pair. 1 where 4 young were banded in 1952. 1 at south end of lake in neadow area. Located nest on dry meadow near dirt road - tookpic of it with 3 eggs at 3 p.m. on the 19th, and at noon on the 20th, with 4 eggs, meading that if an egg were laid each day, the 1st was laid on the 17th. We had a very difficult time finding the nest, succeeding only after one had returned to nest, and, knowing general area, we got the other mate to call while flying nearby, and the one on nest answered with single note. Located on slight hummock in rush-grass meadow 15' from road. Cup $2\frac{1}{2}$ " to 3" dia. and law to 2" deep. Some grass, chips and leaves lining it. Slight shelter by rushes. July 12, 1957 american R., Sacts Calef, new arden Park. I pair sean in a.m. Oct 14,1960 america R. If arden Poste Socto 6, Culf - 2 adaption desco.

6/10/67 Vancours B.C. Ruth. aisjust. w/met.

1973 - at Tour floory:

9/6/13 Stagestin 201 20 - 16 action y history.

1977 - a 9 Cours, years 182 - latte Store. 7/25/76 - CECLESTICLE PICET, Sour Contile Backer Michael Man week with a hot on Cody you from Con 19 mg 1.



Ubmerver fred Lvenden, Jr.

estern colitary Auna, 1, 02

167-169-171-173-174-250-621-859-1309-2133-2261-

Tringa politarla chambar

Late

be di re but ry tim.

Oct. 2, 1943

McChord Field, Wash.

Saw this one and an Upland Plover together on the cement runway. Brownish above, but more light grayish below. Had slim neck and short bill (Legs were a little longer for size than the larger bird. They flew away together and the smaller had shorter and less-tapered wings.

May 13, 1947

Lamile Swamp? Watched one or a muddy flat among the spike-rush at the south end of thwe swamp. It was by itself. It testered and nodaca, and stom pretty low to the ground. The call notes, when flushed, were a little higher than the Spotted Sanapiper, and its flight was of long sweeps, but very irregular and zigzagging in direction. It was a dark gray-brown above, with a wash on the chest, and white below. Head had a gray-white eye ring, and there was no line over the eye. Sup osealy it had a barred rump, which I could not see, it being all write, but inasmuch as a lesser Yellowlegs is supposed to be even larger than this bid, which was about 7% inches long, it is eliminated, and I guess I must admit it as it is, even if I saw what I saw - Ha!!

April 23, 1948

If -mire framp: "To depote this one between a yellowless and a dilideer in a new growth of spine-rush. The character is a for some time and got fairly close to it. It testered is a us once, and when it finally fine of it showed as plainly is bength is stripe down its tall with black and white cross-bass. It also gove its call-sote, which sounded like a spotted sundices only higher.

المنافعة الم



Oct. 30, 1948: Bayshore Drive, Richmond-Oakland: 250& - common. Black and white patch on folded wing. Black-edged white wing in flight. Strong call note.

Bay Farm Island, Alameda: about 500 this p.m.

June 21, 1949
S. Fk. Humbolat F. Elko Co. Mevada - at least 2 pairs in vet perture area.

Nalvista Fanch. Fyruently treading, but I couldn't focate mest, especially in one spot where I was when the birds would make anyened dives at a missing roonly by inches. The must certainly be nextlement, for these actions of the adults indicate it.

June 5, 1952 Carson Valley, Douglas Co. Nevada - About noon Fred Wright and I observed at least 30 of them in flooded pastures andpools at north end of valley near river crossing along hiway.

June 27, 1954 Pope Beach Bar, Tahoe, Eldorado Co. Calif. Celeste irscher and Argantes found a willet sitting on a nest on May 30. The bird did not leave the nest while they stroked it. We found a pair of willets on the lakeshore all p.m. and found the abandoned nest. Located in slight depression between 3 closely—spaced cow-pies. There were 2 eggs, and apparently that was all laid, for each had distinct depression under them, probably from pressure of adult bird - this also explains why nest may have failed from wet ground below it. Took pic of two eggs. They were of distinct difference in size. Larger about 2½ long and smaller 2 ½/8 long, and each being proportionately different in girth. The larger egg was paler in color as well. 2 adults seen.

41,9/6 5 - Flawing, The May The -100's wheeling in several flocker over by area.

Wandering Tattler

Heteroscelus incanus

October 3, 1949 Seal Rocks, Lincoln Co. Oregon. just one on rocks - no flight patter seen. Just grayish - 10% inches long. Very erratic flight - slim-looking. 3-4 syllable melodic note in alarm flight. On rocky islands at low tide, *//2 wave-washed.

Ruff Philomachus muanna

Apr. 18, 1964, "unting Crk, "arsh, "Sl S of Ut 197, Alexandria, Ta. 7 a...

1 "winter plumage" brown specimen with Greater and Lesser Tellowlegs, etc.

On open tidal mud flats, and feeding in shallow water, and were or less aloef from yellow-legs. Also spent much time in abort vegetation, preening, and saw it stretch both wings in u word "V" once, should dark rous "homemones.

A plane flushed it.

• • A STATE OF THE STA

LESSER-2242. 2419-2420-2428-2438-2439-2447-2493-5-7-2/15/59-Exelsión Rd, Sauto Co. - 2657-63.

Logs v v Bred Lvendan, Ja. Creater Yould -- (17 - 187 - 188 - 142 - 384 - 511 - 525 - 545 - 583 - 545 - 602 - 604 - 616 - 710 - 743 - 758 - 827 - 832 - 844 - 848 - 850 - 857 - 1142 - 1147 - 1153 - 1156 - 1159 - 1164 - 1177 - 1187 - 1281 - 1286 - 1289 - 1289 - CLICLE WILLIERO, 12-mile swamp south of Corvallis: Saw a few solitary ones and one flock April 24, 1946 of 30 or so of them wading in the swamp. They wheeled and banked in flight just like Teal. They were very noisy and gave their "kip kip kip" frequently. They appeared to walk on the aquatic plants rather than wading on the shallow bottom of the swamp, altho I don't know. Feb. 25, 1947 12-mile Swamp! when we first came to the swamp, there were 5 in shallow (up to body depth), in company with 15-20 W. Sandpipers. They gave the call note as they flew. Before we left the swampy we saw them again at the north end of the swamp, where we heard their call notes again. 12-maile Swamp: Wa flock of 6 lit next to the Cinnamon Teal at the north April 8k 1947 end, and later on McCallister flushed a flock of 54 from the flooded field at the north end of the swamp. Some landed in fairly deep water and gave the appearance of swimming, although this must be doubtful. Appril 22, 1947 - 12-mile Swamp: perhpas there were 100-125 today, well scattered throughout the swamp area. new thick. It called be a to be, the called which the best of which One flying + city on to to to one, it is an a. Calf. 10-9-52 Jan. 10, 195h Ford I mi. south of C.S.M. 1, on Meelsion load. The thereon i one lesser Yellowlegs secn. Nov. 2, 1957 Yolo Bypses, Colf. It land ingen seen in 2 and their. Jan. 16, 1960 Com Kine Deterdent wordering Carribael, hatte valy 2 the grant of his Call + 1 W Sandy god Oct 14, 1960 - aug R - Sof arden faste, Gasto C. Cety-1002 most.
Fel 1, 1910 grace Car 46 y to fame Cey, sound 100, flocked upe.

A. J. K.

.

.

1

Aug. 2, 1049, Just Wor Los Banos State Leruge I observed 60-65 of them just 50 feet from the road, feeding in a shallow, muccy field jone with avocate and stillts. base 7x55 masses. They seems a long-time at the gring plumage downtcher, but the short bill a linched this ob. In breading plumage yet, and out of range as recorded previously.

Oct. 2, 1953 In a iver istracy, Alleja, Salana Ca. Falif. 1964. . 2 seen.



Tec. 6, 2.47

Con local, the control of the control

Feb. 13, 1972 Under S end, Yaquina Bay Bridge, Lincoln Co. Oregon
200 on flats at lower tide. 1 with white face on left side.

Pectoral time iper

Historia La otos

Pept. 28, 1947 Ferm Hidge Land, which contact the very not to be all the land with a resolution of the land of the following state of the land of the



Observer Fred Lvenden, or.

Baira's Land, iper 1903- 2202 -

Esolia Fisopia bairdi

Tocation we because we have

Buff-breasted Sandpiper - Tryngites subruficollis -

9/23/67, Turf Farm off River Rd, Mont. Co. Md. 21cone on closely-mowed trurf. Brely moved out of war of mower and trator - 1 walked 1 flew breefly. flashing white under wings. Both later flew to bare ground with killdeer and golden plover, but walked bak immediaely onto the closelyemowed turf. Later al flew as together. Lighter breasts thm in 2field guides.

78, tem is on all company of the object of t



Late

Loc tion . . obs. v.ti .:

Feb. 28, 1947

Fern Ridge Dam, Lane Co. a flock of 20-25 "gyrated" past us, and lit 4 mile away; and we approached within 10-15 feet of these little birds while they fed.

Mar 26, 1947

Upper Tillamook Bay: flock of 150 seen by Tom McCallister.

July 24, 1947

Fern Fidge i am Prevetal mochs or them here to be rear along the ale rose. They would circle and dight right in the rose, here there as a secer, and then food into the one more mater whom there. The most I say as 17 in one lock, but more than that the present in the stee.

August 8, 1947 Fern Fiage Lam: Der berore, this species was around the land or raw, and

were very tame, allowing us to a groad within 5 is t of the in "Ting" the fora. Many Flocks of varied sines, in the vicinity.

Ec.t. 25, 1947 (Fein Bione In .: Occ thurse limb son and group even, level to the large area with somitchers, nectured not a control in anily and the main has the about.

Feb. 8. 1948

Fern Ridge Reservoir: many in the area - we watched up to 2 0 feeding on mud flats at close range, and others were mixed in with a huge flock of assorted pipers, probably mostly westerns and a few Fed-backs. In feeding, a small bunch would run down-wind 25 feet or so, and turn into the wind, working upwind to about the point where they started working upwind before. This process was repeated time after time, till they were within 35 feet of us Oct. 30, 1948 Bay Farm Island: Alameda: at least 30,000 least and westerns in one flock.

Sept. 1, 1949 Lower Alamath D. Calif. flicks nana avery out. Dec communities westerns. Picked up a botaline victims. The



Pelian alvino scaladina

Late Locati : a no observations

May 13, 1947

of this species, in company with a more or less uniformaly light gray of the same size, perhal a abira of the opposite sex, in winter plumage, yet.

11-24-47 Lec. 7, 1847 Bill Williams R., E of Parks Dam A by gale Monson - Coloris, Assertion on the interest.

LUCE - Coloris Colori

Feb. 8, 1948 Fern Ridge Reservoir: some mixed in with a huge flock of Westerns and leasts.

April 17, 1949 12-mile Swamp of Thock of 75 working over mud around flooded field at first turn in road. Brack berries very conspicuous. The once flushed by the Cooper's hawk, but came hack. Later a flock of about 10 returned. In with the red-backs were about 25 smaller, probably western sandyipers (LEM).

Aug. 2, 1949 Mendota Poor, Calif. in tinter prumage.

Feb. 24, 1950, 39-ride Stamp, Burton Coll DBM. Throke of arout 70. Standard window plungs. Fight place coparate flow rate for a surface the flow was in a flooded field with rye grass coming up. Was at first confused as to the identity of these birds as this is the 1st time I have ever seen them in winter plumage. Had killdeers with them to compare for size.

Apr. 1, 1961 Bodega Bay, Sonoma Co. Calif. 100's, some full reddish adults with the black bellies.



.U.D.) OI 116_ 01 1 16_ 01 16_ 01_ 01 16_ 01 16_ 01 16_ 01 16_ 01 16_ 01 16_ 01 16_ 01 16_ 01 16 Limnouromus sriseus stole des

Late

Location une out el veti T.

April 27, 1941

to be a run-occurrence.

March 27, 1947

May 6, 1947

:ept. ≥ , = 47

Feb. 8, 1948

Sept. 8, 1948

Sept. 1, 1.4.

October 19, 1949

newport: Of attinuing in the second a pone of the life in the land to be a

Dumbarton Bridge, Lower San Fran Bay: / 40 or 50 Dowitchers on the flats.

12-mile Swamp: a pair seen at a distance, and then closer, and white area on back and rump was plain. Prumage was richly colored, chestnut. lan ricac lum: "" o our arver a riben et a districe, et l'inseme en entair within the tree can they comen't hause the line is a more than and hered in. There are a month about the month of the contract of the contrac to set so was in our iventer re-city, on like so to the so satisfied Line of binter. Together an allocated about 17 firms, the Lites, on the couth miles year and a dense drock or o -75 direction of the cost of the section .

Fern Ridge Dam: Na flock of 5 were feeding in a slightly isolated groups a ways from the huge flock, of other species on one of the mud-flats.

Sacra. Wildlife Refuge: scattered pairs flying around. Small flocks of 5-20 birds seen, and we saw one huge flock of 300-350& birds, containing few avocets and stilts.

Lower Alexath lake, Galif. rest to Least and Western sancyapers, this is the most numerous scorebine. ave confected 7 between viction.

4 miles 50. of Socramento, Calif. 8:30 a.m. 150-200 in irrigation overflow- sond. and I mi. b of G.D.M. 16 on Excelsion Good Payer 1 . Alch List. Law many feeding in shallow water with westerns and locate and pollow-logs today. byen more present on I r. 17 th 11:30 a.m.

Excelsion Rd, Suito Cos, Caly- 25-30 w/ 2 Japan Gallow lago.

Location and observations

Light of the first of the control of lean tally point of the lower state of the control of the leant of the

or Comment in the Comment

Apr. 12, 1944

Mtn. Home Reservoir, Idaho. Saw 5 along the edge of the water. Quite small, 4-5" body, black bill and legs. Bill 1-12" long. Tawny mottled above. Gray white below. Black spots on folded wings. Plaintive call notes. Head and neck, onto chest, a light gray-brown, streaked. A definite demarcation line bet een this color and the underside gray-white. on the chest. Their warning notes seemed to be a musical trilled-rattle. Seemed quite bold. Allowed me to approach to 10 or 15 fert, upright, before they flew.

Feb. 18, 1947

Greenberry Flats, Benton Co. 2 flocks of these sandpipers were seen in a grain field beside the hiway. One flock had 64 in it by actual count, and the other was slightly larger - perhaps 70-80 birds. They were a light grayish brown above, and a clear white below, with just slight dusky suffusion across the upper chest. The bill was slender and at least as long as the head. The legs were black or dark brown. No dark patch was seen on the forehead.

Feb. 25, 1947

12-mile Swamp a flock of 15-20 in muddy field east of the swamp, which flow into the swamp and joined the Yellow-legs, later, as we left the swamp, we saw them in the field again. I approached within 30 feet on foot, and observed them feeding, bathing, and preening while standing in 2" of HOH.

Sept. 20, 10.7 Februaries

Producter of the state of the state of the value of the state of the s erec, with so itemes, rector a smaller tr, in link or a survey. Jo. i then the goat's.

Feb. 8. 1948

Fern Ridge Dam: in a mixed flock with the leasts, a huge flock of several hundred or so, and perhaps, a few Red-backs in with them.

Bay Farm Island, Alameda: at least 30,00 westerns and leasts in one flock'. Oct. 30, 1948

Tule Lake, Carif. Logt. 1:491000's on the area. Lave picked up 8 boarden victies. July 18, 1957 Pope Reach Marsh, T. Taboe, Palif. Watche one walking on bond lily

(snadderdock leaves this afternoon.

James 6, 1960 Detecting Bottoms, Clinet K. Szicto Co, Caramichael, Calif. 1 an growel bay

Observer Fred Evenden, Jr.

Sangerling 1810-1811-1899-1938-2068-2187-182-192-193-204-441-776-1003-1039-105) -/1/0-1330-1380-1386-1567-

Crocethia alba

Location and observations Late which the analyse that while a; made, should be made at rain by made,

Dina, Tal, the low rie rigidital process of the line of the outlinet name to we came of the fill to he is a sure of the outline of the outline is a sure of the outline of the box of the or 5 minutes, to the outline of the outline of the outline of the box of the box of the or 5 minutes, the most circles. It is a sure of the outline of

151 144 (miles)

Location and opserv think

1104-1180-1192-1197-1201-

Mtn. Home Reservoir 4 to 6 p.m. NV18, 60°F May 17 1944 Saw one pair and a single, wading bealy-deep at the eage of the water.

Mar. 27, 1947 Dumbarton Bridge, Eower San Fran Bay like PST - I saw 40 or 50 of them. Mar. 19, 1949: NW of Vallejo, Solano Co. feeding with side-swipe forward notion.

April 23, 1949 Grey Loc e Refuge, Butte Co. 500 reen. reca by ralking forward, slueswiping their bi, l as they go.

May 5, 1949 Gray Loage Refuge, Sone small patch of "salt grass" where alkali concentrates were on top of the soil on this flat area contained 2 avocet and 1 stilt nest. The adults were on the nests in plain view of us, just 50-0 feet from the car. The nests were always in the open, without protection of any vegetation more than 2 % high. Found 3 avocet nests - alm contained 4 cgss. The eggs. were strongly talered at one end, a gray-green color, with darker blotchings. They were larger than the stilt eggs, and a lighter base color, as well. I took one pic.

fest. 1, 1949 , Lower Klamath Bake, Calif. Till - sha e i Docus; ducy are e, i s in considerable in hers of botuling. Its of ther cane.

May 1, 1952 Gray Lodge Refuge, Calif. many seen. Several nests with 3,4,4, & 3 eggs. John Cowan says they were present about 8 weeks ago.

June 4. 1952 Carson Sink at Pelican Island, Churchill Co. Nevada - observed my first downy young. Bill short, and only slightly recurved.

Feb: 3, 1953 Forro Bay, Calif. 190 in loose flock at SE corner of marsh- few cetting brownish-red heads.

5% corner, Take Tahoe - 'n adults at mosth of I the Truchee Higgs. Calif. Bay Farm Eshand, Alamoha Co. Calif. 100's seen - none with reddish honds. Lay 30,1953 Feb. 6, 1954 ondu, industry Tolour, Tolour, Tolour, Talling 14 sect. In late and in bear food Gustino Gun Jub, joyaco ju. 5011. 121s sect. Incomi Sept.27,195 oct. 1, 195

Yolo By son. Woodland doad, yolo Co. Caly. 3014 am. april 23 1958

June 6, 1959 Sears Point Rd, W of Wallejo, Solano Co. Calif. and Oakport Road, Alameda Co. Calif. where I watched one swimming deep tide channel Oct. 10, 1959 Oakport Road, Alameda Ch. Calif. saw 40-50 avocets, but none with red heads.

A CONTRACT OF THE CONTRACT OF 4

russer of liels traje on liel on Ivea;

Biack-necked tilt, 284-1296-1299-1307-940-947-457-1180-1192-1200-1225.1239-1243-4251-1256-1256-1281-1691-1917-1948-1962-1900-2097-21/3-2436-2531-2532-558-558-15816-1589-1600-1615-1617-1625-1655-1656-1657-681-2594-

Himantopus meniconus

Location and observations Date Mtn. home, Idaho, Reservois - 4 May 12 1944 4 of them were wading in the shallow water about up to their "ankles". These are the first I have ever seen. They give a "yap !yap ! call. White below, black above. White face with black

around the eye and on the crown. Legs are a readish pick. Very long. They drag them behind on the takeoff and hold them out behind in flight.

Throw them forward like a heron when aanding. I was within 25' of them.

May 5, 1949 | Gray Lodge befuge, ST Butte Co. Jonly saw one nest, containing 4 eggs. It was with 2 of the avocet nests mentioned above. Nests were about 15 feet apart. The stilt eggs were the same shape as the avocets, but clightly smaller, and the gray-green color was much cacker than the avocet eggs, and the blotching was much the same.

June 30, 1949 | Stillwater Glarice Againt Treat Churchild (c. Tevada: 3. monto, 3) of which were concentrated in area sere water to bbout 4" dee in the nut rans. A thin concentration of a toward. That B young.

aug. 2, 1949. Los Banos Reluce, Chif: /S of stilts seen vere in ture, its lowngrayish black, instead of pure black plumers. he -1:5 com. 3 tits for every avocet in trea.

Sept. 1, 1949. Lower Alamath L, calif. Ind - only one seen, and it has not unique

May 1, 1952 Gray Lodge Refuge, Butte Co. Calif. Quite a few present. Cowan says they have been present about 3 weeks. New nests, but no eggs found. June 9, 1957 Natomas Dist. along Elverta Poad, Sacto. Co. Calif. 15-20 in one rice

check, and many others in the area.

June 2, 1960 Pice field Sweet Eleventer Substation, Sante Co, Cally

·

Observer healtvancen, Jr.

Rea Phalarope 1201-1761-2187-2387-4-9-2688

Phalaropus fulicerius

| Lute | Location and opservation, | | | |
|-------------|---|--|--|--|
| ay 21, 1949 | Monterey and S. of Pt. Lobos, conterey Co. Several seen with Norther Phalaropes on ocean in kep becs with the sea ofters. Nost in winter brumass. | | | |
| ov. 2, 1952 | In the late afternoon I saw I full adult working in sliny, shallow pond with Nrtherns (145-150), north of Wonter ey. Dillon's Beach, Marin Co. Calif. The night of the 1st, while on the beach about 10:30 p we heard this call, but couldn't check till morning. This morning we found several (12 or so) in winter plumage, and hardly recognizable as their action habits were completely changed, as they walked on the sand, after insects. So out of character: | | | |

15612-1615-1946-2099-2105-2110-2119-2121-2347-2756-8-2523-

tteganopus tricolor

Late Location and began time

May 12, 1944

Mtn. Home, Idaho, Reservoir - Saw 20 or 30 feeding on the damp edge, or wading into the reservoir. Birds seemed mostly white on belly, and with write down back of neck, bordered by black, and this fused into reddishbrown, on side and front of neck and shoulders. They kept giving deep gutteral notes. Back was dark in comparison to underparts. No wing stripe. Some were still in winter plumage. Saw one female chase another female away from a male. When they sand from flight, they extend their wings vertically over their back.

June 21, 1949 | Battle Mtn. - blko Nevada: several on stock water pond it 58: 1. No marsh vegetation at all. Just open range and sa elruch.

June 22, 1949 | Mary's Fiver, alko Co. Nevada: dozens of pairs. Pairs often cardled within 5 feet of my hear, giving singular notes, continually.

July 15, 1949 Stillwate Refuge, Fallon New - letter from Dave Marchall - 7-17-49
6,000 at a sund flat lake at Newd of project - 16 mg mi of Ruppia
also 4 shawy Planers.

Aug. 2, 1949 LDs Banos Lefure, Calif. in winter plungs.

Aug. 2, 1949 Mendota Pool, Chail. in winter plurage.

aug. 16, 1949 May Bray Looke Ferune - butte Co. Chair. two barr cotton to in the ci

6/10/67 Vancouve B.C. Ante air parts. Then I must went if wagget Degressed, Defined, grass- lived cup.

, the st

e .

Nort ein Palatione 77-152-307-309-883-997-974-1200-1201-1299-1305-1307-1309-1318-1340-1633-160-1761-2266-2267-3268-2273-1274-2317-2317-8-9-2421-2530-2575-200-1201-1299-1305-1307-1309-1318-1340-1633-

Lobipes Lob tus

- inte location und observations
- ay 18, 1941 | 18 mile em sy cours a browning. It it is two: can be a more to while they are a minution. That is not the case of a brown or y, buth reducish at each of the case.
- May 12 1944 | Mtn. Home, Idaho, Reservoir 12 of them, out in deeper water, swimming around. Real jitterbugs! Gray head with redaish-brown on the sides of the neck.
- Beach south of Yaquina any channel. A smooth as he is an a durant (sith-out nufrus across tiont) on to largest on the cradit ones among the auner. Got within 15' as it with these, as it went the .

 It was well, Taying N.M.
 - Oct. 4, 1948 near Silver State Gun Club NW of Leetville Jct, Churchill Co, Nevada:
 7 in winter plumage black bill and feet. Black "tear over eye area.
 In small pool at edge of hiway, and allowed me to approach within 10
 or 12 feet of them. Using 7X glasses, they were big! When flushed they would fly only about 4 or 50 feet.
 - May 21, 1949 S. of Pt. Lobos I saw some on the ocean kelp bear, nor the sea lions. They were still in winter plumage. W. of Monterey I saw about 15 in a slimy, shallow pond about 6 p.m. Poth cases associated fith Foc Phalarmes.
 - Sept. 1, 1949 Tule Lake, Calif. and various ill on numbering lawses ab biron seem on mover planath. Figure us a botunism victims.
 - Oct. 15, 1950 S.F. Bay to Farallon Islands The return. 2 or 3 small groups of 3-5 seen both ways. All in winter plumage. There were 1,000's in winter plumage on the waters around the island. Some stayed within 15-20 feet of our boat. All tended to remain in loose flocks or raits on the water.
 - Oct. 1, 1950 Tono Tandi a, Intere h. 112.3 service 7. o 100ms of 10/3
 - Nov.21, 1959 Bass Lake, Eldorado Co. Calif. 1000' el. 9 a.m. 1 in winter blumage walking in curious manner on mud flats out of water. Back & forth.



Stercorarius parasiticus

Parasitic Jaeger

Oct. 15, 1950 S.F. Bay to Faralion Islands: No full adults seen, but about 8 birds lacking tail elongati ns were observed. Best tild by comparison of head markings when the approached.

Pemarine Jacker

Stereoranine pomeninus

Jan. 35, 1956, Mear Thornto. . Encto. Co. Casif. I in frost of the m clouds.

| | | | | ٠ |
|---|--|--|---|---|
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| • | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | • | |

Stercorarius longicaucus

Long-tailed Jaeger

- Oct. 15, 1950. S. F. Bay to Faralien Islands: Saw only one furn about with 8-10" tail elongations going out and one coming back. "Tailies," birds could not be identified during the a.m., but on return trip there were several that approached within 40 feet of our boat and of 7 seen in 5 minutes, 4 were quite definitely long-tailed jaegers. I watched one jaeger chase 2 gulls in a.m. and in p.m. I watched 5 long-tails course in a close group behind the boat, cutting across the wake the gulls were 10 llowing.
- June 16, 1975, St. Pawrence Esland, rear Cambell Village on top of Gevenue's tro.

 This was do mon larger, the all 3 may occur in one ones at ore time, a bi tundra Cl of mtn. a pair came within E 18 for inc.

 The rices the tundra, and as suppossible they mated for 2 2 decords, any heard giving long, draw out corean call as they sailed over the ntn. ridge above the cors and flats.

| | | | <i>'</i> . |
|--|--|--|------------|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | ` |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

Larus heermanni

- October 30, 1948. Seal Rocks, Cliff House, San Francisco. Noon: Immatures are nearly black, w/red bill. Adults with white head, black tail tip, and red bill. On offshore rocks. None known inside the Golden Bridge.
- October 5, 1949

 Yachats Park, Lincoln Co. Oregon. 1230 noon. 3 dark-headed individuals 1 full dark red bill and white trailing wing edge, and adult tail; other 2 gulls with dark-tipperd pale bills. These gulls scattered through flocks of the other gulls western, California, herring and ring-billed! On the sands exposed by the retreating tides

 November 6, 1949

 Bolinas Lagoon, Marin Co. Calif. On Stimson beech there were 15 or 20
- November 6, 1949 Bolinas Lagoon, Marin Co. Calif. On Stimson beech there were 15 or 20 on the sand beach at edge of surf, and feeding out with the retreating waves. The sand beach was sharpely terraced at this point.
- Dec. 6, 1949 Conterey Bay at Santa Cruz, Calif. 150 on the coast, and lobserved in creek or slough bathing & preening with other gulls in town.
- Oct. 15, 1950.

 S. F. Bay to Farallon Islands. Saw very few goars out but about 20 of some 80-90 gulls that followed our boat for many sides back were Heermann's. Of the 20, only 4 were definitely full adults. About 50 were seen around the islands, the ration being 1 adult to 4 innatures.
- Fecember 19, 1950 monterey what 70 (oright 15-20 of which were imputures) in ricky tide pool area at Pt near foghorn. Also a half-dozen innutures and part-acults at the monterey wharf.
- Nov. 10,1962, Menterey, Calif. 6:15-7a.m. St. 6-700 on parking area at Estero 7., and these left and were flying but to 8 side of rec't islands off Pt. Pinos at 7:15 at 1000 el. while Calif. and N. Bulls were coming east.

Observer Free Lvender, Jr.

Mestern Gull

70-80-84-84-118-124-113- (162:) - 191-218-242-249-257-355-365-366cambro of field prick to lie

iocation and openv trend

August 8, 1947 fern Fiage I am Faw a impature birds of the year figling the area, and round

one agas at the cage of the rote. Plumage saits a relation.

S. F. Bay to Farallon Islands: Some all the way out, and about 6.-70 followed our boot back for man, miles. Of the 65, 35 one lst-year im ature 20 were End-year immetures or transitions thereof, and 10 were rule adults.

They were the most common gul, on the islamus

Feb. 5, 1950 Mc loan Tourist and the most common gul, on the islamus

Several till a memorial to be bounded by.

Jecland Carlot 12/14/16 Gode States of hick

12/14/16 Gode State



Observer Fred Tvenden, Jr.

Lerus ergent tus smithsoni nue

Late | Location and obscivations

Sopt. 24, 1948 | 1. Isan. Lay well with an interest to the contract to the contract.



Lumber of fiel trips of thick observed: 102-137-190-191-192-193-204-266? - 339-365-366-446-488-489-490-515?-

pril 10, 1844 (crvali, or tom chamber a man colm of a sil a t minumetr remark. The officer an interest of the policy of the order with this continuous and a mark later part of him to the factor abort L b" long

Yachats River bay! standing in scattered flocks on sand bar in the bay, Jan. 25, 1947, and shifting with the small changes in wind direction, taking advantage of what "air pockets" they could find in this near-gale storm. An example of a response to micro-environmental demands.

Yaquina Bay, south side - a flock of around 500 standing on a sand point Jan. 25, 1947, in the bay, semi-shelter from the strong SW winds.

March 28, 1947 Berkeley on Ferry to San Rafael, 9 p.m. several flew by the ferry after dark.

Sune 2, 1950 Stillwater Refuge, Churchill Co. Nevaca LEM. Found lb nests. They are a depression lines with small sticks. One nest empty, 3 nests with 1 erg; 4 nests with 2 ergs; 5 nests with 3 ergs.

March 10, 1959, Kalama-Kelso, Cowlitz Co. Washington. 10,000's scattered along miles and miles of Columbia River and sloughs.

Number of field trip on which observed:

284- 356-17987/-803-832-791-1040-1028a-1029a-1029a-1029a-1029a-1028a-1029a-1020a-102

Larus canus brachyrhynchus

Date Location and observations

Mar 15, 1944 Pudding River, Woodburn, Oregon. 150 of them followed behind a tractor and plow, feeding on earthworms turned up; near Percy Seeley's place

Jan. 25, 1946 Hughes! Bottom, Butte Creek Road O' Several 100 of them, mostly immatures, Adults had yellow-green bill and feet. Tail white, mantle light pearly grey, rest of body white. Gray brown body on immatures.

Dec. 22 +23, 1949 New Era Clackanes Co. O regon - Heavy wind & rain storms jast kur days & ofserred small locks of these gulls the jast two days, working a freely proved field adjacent to the kiway. Most on ground - few in air.

,

Dreiver Freu Evensen, Jr.

Date | Location and observations

ug. 7, 1041 | nation of an of the many and the transfer of the same of the sam

March 27, 1947 Hayward, Calif: abundant, and following newly turned furrows of the plough, in a field next to the highway.

Sept. 1, 1949 Thile Lake, (alif. the total the mull tocies are due by laturial. Thec. 6, 1950 Oso Flaco Lake, S. Luis Obispo Co. Calif. 645 p.m. dozens pouring into the lake at sundown from the cultivated farm lands to the hand is.

- Feb. 22, 1951 Laguna Madre, N of Port Isabel, Texas perched on sand spit at edge of Laguna. Gave call seve ral times.
- Feb. 20, 1953 N of Galveston, Galveston Col Texas. 1st at Sabine Fass, then several dozen followed ferry to Galveston and I tossed them bread, which they caught in stiff breeze in mid-air.



Larus pipixcan

Franklin Gull



Oct. 4, 1943

.di dei di 11/2 di 11/ 2338-2341-2387-2583-4-2637-8-9-42-3-54-58-Larus Milacel, da

Loc tion and observations

Vashon Island, Magnolia Beach, Tacoma, Washington Encounter ed several large groups of them flying over the wateror all perched on the sand bars that stick out into the water.

March 27, 1947 Dumbarton Bridge, Lower San Fran Bay, 1100 PST, quite commo, for I saw 25 or 30 riding on the water.

May 12, 1952 Carson Valley, along U.S.395, N of Minden, Nevada - 2 p.m. 2 seen, but neither in full adult breeding plumage.

May 16, 1952 Tulare Lake, Kings Co. Calif. 1 seen, not inful adult breeding plumage. Nov. 3, 1960, Napa R. Estuary, Solana Co. Calif. several in imm. or winter plumage.

Rissa tridactyla pollicaris

- March 26, 1947 Ecola State Park, Clatsop Co: & small flocks of gulls heading north offshore. Solid black wing tips noted. Most likely of this species (by Tom McCallister)
- March 28, 1947 Manzanita Beach, Tillamook Co. one dead one one the beach. Similar in appearance to a Short-billed Gull except for the black wing tips and the black feet. Adult (by Tom Mc Callister).
- October 5, 1949 Neptune State Park, Lane Co. Or gon My first obs on this species.

 I saw 3 abou # 1/4-mile offshore, paralleling swell crests with the two shearwaters sooty and pink-footed.
 - April 19, 1951 Pt. Pinos, Pacific Grove, Calif. 1220PST. some adults and immatures sunning and preening on rocks at edge of hiway drive. Saw 55-75 off Pt. Pinos, too.

Feb. 22, 1951, Laguna Madre, N of Port Isabel, Texas. some perched on sand spits, and a few flying, thus giving good view of black and white pattern on the wings.



Feb. 22, 1951 - near Port Isabel Texas - o ver open pond near hiway - salty water.

6/5-6/ De Chiecologie NWM, Vs. Serger ten 710 + cofilower's wing cont.

Observer Fred Evenden, Jr.

CASPIAN

Number of field trips on which observed:

236 249 251,923-1182-1217-1239 1243-1275-1284-1303-1313-1544-15612-2119-2137-2152-2173-2242-2499-2421-2530-2531-2627-1646-1657-1646-1686-1687-1705-1710-1743-1855-1945-1963-1961-2103

HYDROPROGNE CASPIA IMPERATOR

Date Location and ob

Sept. 22, 1948 Steilacoom, Wash.

Location and observations

First I have ever seen. They were quite small when compared to the gulls with which they were arguing, to see which would possess the chances at a school of fish below them. They had a high pitched call. Colorations were not very distinct. Have about a 12" wingspread.

Oct. 2, 143

Vashon Island, Tacoma, Washington.

They were always pointed into the wind. Foreheads white with black more to the side of the head. Got up within 15' of a small group of 8 of them perched on the railing of Shawnee Dock.

July 27, 1948

last side, Alematic and the control of the contact of the control of the contact of the contact

Aug. 2, 1949

Memota Pool, Calif. Some us illute. Thushed and circum about me, giving very raucous call notes. Acta ed one and a 7 '-diversite the san Joa win hiver, and call up from complete number end with a my-4" fish help cross-wise in its bill. The Illument shing-shaver color - probably a caprilic results at the surface.

JUne 2, 1950

Stillwater Refuge, Churchill Co. Levaca DBM. Found 107 nests present on the island. Wests a depression but not lined. 23 nests had legs; 76 nests had legs; and 8 nests had 3 eggs.

× 4.3

,

().-

•

•

986 -1001-1003-1054-1251-1264-1265.1351-1578-1579-1580-1630-1704-1758-1829-1854-1899-193 2-1853-1854-1862-2113-2457-6-Common Tern

Sterna hirundo hiruno

to the second of the second of

£

Sterna paradisaea

Arctic Tern

Oct. 15, 1950. S. F. Bay to Faralian Islands: A seen about 1 or en out 1) on Golden Gate in morning. Perched on illoating piece of wood, but Illushed when we approached. Circled us so we had good view of them, seeing red bill.

Observer Fred G. Evenden, Jr.

FORESTER'S


Puncoick >

Caupian Tern

1265-1273-1275-1271-1284-1289-1291-1289-1291-1289-1313-1319-1351-1579-1581-1603-163-1632-1263
STERNA FORSTER! 1758-1899-1919-1938-1953-1856-1803-2036-2035-2096-2239-2243-2243-2376-23343-2-2356-2381

Hydroprogne caspia imperator

| Date | Location and observations |
|---------------|--|
| July 27, 1948 | Fart side Minath Lac: " three terr, the contraction - three |
| | Ctillwater Eget Filolife Thea, Crurchill Co. Fever: 10 Found the island of nutorass. Island and ranted down except for June, the will to quite high by the dealestalks. I mesus of the island, one with a eye and the other with A. Eges and or the nutorant stalks which has been want to slightly where the eggs were high. 71 birds on the island. 100. |
| Sept. 1, 1949 | Lower Alemath ., Gali. 75 in one laock. The |




California beset Tern

cterna albilirons browni

August 15, 1949 Benicia, Polano Co. Ca ir. Est eoge of city Ri its et agenter! Fgrry Boat House. This moon I spotted 3 very small terms with neveral ower short-tailed ones that proved to be Forester's Terms is varius at les of plumage. The S smaller ones had cusky billes, more year on at the base, and the legs that distinctly in Alo ish in tous of or nother a in the Forester's. Their call was higher pitched than the lor ater. . They they had dark patches on the sines not gree nt on the Fore tere. There is little doubt in my mind that these were Least Terms - a new line-wister for me. All these terms were morking tog ther against a north wind along the east side of this arm of Carquine: Straights, hovering about 10 feet over the water, and plunging below the surface whenever they spotted a rish, or other item of food. They would work upgind 50 ares, and then drop swiftly back to the "foot" of the feeding orda, and start the course over again. They were observed to eat their catch on the wing, and then proceeding with the fishing. This section of the bay has a number strip or summy beach, with some emergents such as desquare and cott ils on the inscession of the sand areas. Rocki jamtituded from the bay in shallow water. Turky water.

5/24/70, Ocean City, north end, Md. many with 2 eggs under them on sand fill of RE development



Thalasseus maximus

Coyal Yern

- May 15, 1948 12-mile (Wamp Pr Jac Von Torrstel first spotted it. A rule plum ged data male with white under tail ilying slowly to the north. Had a jectular type of wing-beat, as if it was about to check its forma a otion. This sugas saw this species on the pring bird count most you at this count, but this is the first time I have seen it, ever - a new sind let by last list!
- July 187, 1840 Upper Alemati Lengthing of the control of the contr in Tur, or hot walls hading.
- Gray Louge Refuse, 67 Butte Co: One was seen on way u here from Sacramento May 5, 1949 this a.m.. They are con on wherever the fields are being flooded for ricegrowing, etc. about 10 days ago I was up through this same area, but not a tern was seen. John says they just arrived at the refuse during the part
- few days, evidently working their way northward. SE of Trowbriage, Sutter Co. 3-400 over one 50-acre chunk of rice paddies May 6, 1949 that were floaded.
- June 11, 1949 12-mile Swamp, Benton Co. Oregon: 320 p.m. I coursing over north and of swamp. Came with m 40 or 5 mest of us. A full, spring-phuma sou acult.

Aug. 2, 1949 Los bands Feliage, Collin: post in a ring plumase, but some inter-or impature plumases of n. 200

August 13,1949 yolo By-pass, W of Sacranento- 645 a.m. yolo County. about 100 adulta and imm. over rice fields. Fast ones seen in area, this summer. May 1,1952 Novel for Single of the Single of the Modern of the Miles o



Rynchops nigra

Feb. 21, 1953, just N of Rockport, Aransas Co. lexas - after covering miles of gulf coast in stiff breezes past 2 days we found a raft of at least 250 of them on bar at e ge of Gulf.

6/5/90 ChiceTagus MWA. Uk. Historiatorial long laws. Lalways Community consists + actions




Californi Mun 113561888

Uria als e liversica

| Date | |
|---------------|---|
| ug. 1, 1942 | to or other. |
| March 10 1946 | Nelscott and Boiler Bay A found one alive, but heavily oil souked. It was easily approached. Saw one later in the day on offshore rocks, just north of the Devil's punchbowl. |
| | T. F. Bay to Far allon Islands: these were most abundant Alcies seen on trip out - perhaps a total of 150-175 in winter plumage, sincly, or in groups of 2 or 5. Very few seen on trip back. Very chop, rough sea on way out, and calm on return trip. About 40 about islands. Monterey Penin. Monterey Co. Calif. Seen in summer plumage, Cil Araked. |
| | To Farallers: Very few seen till at least half-any to tell of a line fall durage. Few about Signous, very content from them, to leaven. |
| July 5, 1970 | Depoe Bay, Oregon. 100's coming and going up and down the coast, both close in and offshore. |
| July 6, 1970 | Three Arch Rocks and Cape Meares, Tillamook Co. Oregon 1000's . Could hear their cries from the rock islands above the surf, long before we reached the coastal view. |
| Feb. 2 1975 | Otter Rock, Devil's uncl Boml, I nooln. Co. Oreson. 1015 a.m. watched 2500-30-0 pesled off top of rock from orth to south |

and out to sea to feed, just like ulling a take off a table!



164-441-566-567-551-884-1199-147-1201-, 1634-1688-1689-1760-1761-1877. igeon Cuillemot/455-2306-23//-2422-

Cepphus columbu

2679-60-65-

Location and opportunitions

Ecola State Park, Clatsop Co: Pall in spring plumage. March 26, 1947

Yaquina Head, Arate Beach of san heveral doren of theo recline in the work-June 15, 1947 ing sea off the sough side of the head, and maying back and forth tox the steep rocky walls of the Head, up to 50 feet above the laber. The blief stayed pretty close to a scall ledge which has have been a roll site. Two were swimming close together plishore so I threw/ a rock between them, and both dived immediately, one coming up again in mediately only offect away, and took off into the air without so much as remains on the currace - she practically flow out of the water, and the other bind stayed under over a minute, and care to ab -1 01 aday.

Yaquina head, just il is eate deach of com on increase and come, and May 31, 1948 e untime untile, on the rick. The classifier, is not a fine of choor ten chin, , can be been any recin customer.

Depot Bay, Oregon - 9 flying north in steady line.

Aug. 20-21, 1961 Depoe Ray, Lincoln Co. Oregon. adults feeding young in nests in sandstone cliffs having easterly exposure, opposite folk's cabin.

Aug 22, 1961 Battle Rock State Park, Curry Co. Oregon. 1-1:30 p.m. PDST. carrying food to young on north face of Battle Rock.

The reports to any weekly and offer to South Comy, Other adverty completed weeks. \$/26/57 -July 5, 1970

Depoe Bay, Oregon. As many as 25-30 adults off Martin's. Some coming and going regularly from sandstone paiff nest cavities.

Black Guillemot - Cepphus anylle 12/27/65, Bethany Teach-Tehobota are . Susaex Tr. Delawire. 2 off Diwey Teach.



American Marbled Murrelet

Erachyramphus m. marmoratus

June 24, 1950 Monterey-Point Lobos, Monterey Co. Jobserved one around low rocks just offshore of mouth of Malpas Creek south of Pt. Lobos.

Ancient Murrelet

| | , ** | * • |
|---|------|-----|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| • | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Cassin Auxlet

- Oct. 15, 1950. S. F. Bay to Farallon Islands: Wonly 2 seen on trip out, but 20 or 30 seen on return. At the islands, all in eark winter plumage they would dive at the slightest provocation. about 20 seen usually 2 or 3 in one small area offshold. Some flew and flog ed ahead of the boat with typical cuck "vounded and lead-away" antic.
- October 11,10. To Farallons: 2 or 3 within mile on two of into be. Auntion toward Coast in ... Some about 54 error, not berook the extent world make mins to succee before living. For community to 1 groups.

| | · | |
|--|---|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Observer Fred. G. Evenden, Jr.

Rhinoceros Auklet

Number of field trips on which observed: 192-225/-2387-9-3573-79-

| Cer | orhinca monocerata |
|----------|--|
| Date | Location and observations: |
| Mar. 28, | 1953 Monterey Penin, Monterey Co. Calif. In summer plumage - oil soaked. |
| Aug.21, | Parakeet Auklet Cyclorrhynchus psittacula 2 mi off Depoe Bay, Lincoln Co. Oregon, in Pacific Ocean. 3 of them seen within 30 feet of boat for several minutes. |



Pilot-Biologist

Aerial Colonial Bird & Manual Census - Oregon Offshore Islands, June 13, 1970

The following resume indicates over 200,000 colonial birds were mesting on the islands this summer, 1,815 sea lions were observed and 80 harbor seals were also seen.

A significant loss of sea lions on Three Arch Rocks was the most noticeable change from the census made a year ago. No dead animals were observed on the island and none were observed floating at sea or on the shore.

| | *** | | | | Sea | Harbor | |
|-----------|---------|------------------------------|-------------------------|--------------------------------|------------------|-----------------|---------|
| •• | Islands | Murres | Cormorants | Gulla | Lions | Seals | Puffin |
| Goat Isla | ınd | 2,500 | 180 | 650 | - | -(ANT) righteds | * |
| Twin Rock | | 3,800 | 180 | | - | 4944 4844 | * |
| House Roc | | -140 MH MP 144 BM | 50 | SAME TRANSPORTER | APTA curso dista | pay dis. | * |
| Whale Hea | | 8,000 | day and one | 600 | | Apir data | 125 |
| Leaning F | | | 50 | state stiger-acco | tour that tour | nero tanà | 4.0 |
| Mack Reef | | 6,000 | goal Seek oute | 200 | 445 WW 1946 | 10H duth | * |
| Crooks Po | pint | com some some some some | district county compar. | 75 | - | and the same | -X- |
| Hunter Is | land | 17% Trip days done made | 50 | 275 | arth days graph | METE 1966 | * |
| Rogue Riv | er Reef | 2,000 | 150 | 50 | 380 | 45 | 77 |
| Hubbard I | | 2,600 | 100 | - States - Special - States of | del me las | dies and | arya |
| Island Ro | oek | 3,800 | 200 | 600 | - | space comm | * |
| Red Fish | Rock | 10,500 | gare meri. Univ | AND SERVICES | | 414° 109 | aprel . |
| Orford Is | land | 8,000 | 4m6: 840-760 | med dedit famy | 600 | die eri | Day . |
| Blanco Re | ef | and one post-bid dept. | 800 | 946 949 888 | 3002 466 494 | 580 Mer | dis |
| Gull Rock | 2 | 6,000 | 900 | Trial State Comp. | MOT MAD SAME | prov dner | ¥ |
| Castle Ro | ek | 2,600 | 600 | name calvin diplo | que des ses | see one | 甘 |
| Tower Roc | k | 6,000 | 60 | State Build Amer | gree detailed | 6047 SER | ** |
| Pace Rock | | 12,000 | 300 | pleasure defined | ann time agai | 440 too | * |
| Coquille | Point | 700 | 50 | stately strain streets. | atile through | And myl. | ##* |
| Table Roc | | spile state state than state | ands down union | 400 | year thin cast | 000 - 00g | * |
| Cox Islan | id | dies car due dies ses | rest only | dated 1960s drap | - | and alles | 40% |
| Sea Lion | Caves | with the day and with | no nest- | mer specific | - | Mg/K-ramit | 200 |
| | | | ing birds | | | | |
| Heceta He | ead | cup over that and their | 125 | dec dep sea | 275 | 1949 sales | page |
| Conical R | lock | cons dest page gene sign | 110 | 200 | 125 | NAME AND | × |
| Seal Rock | | destroyer and date date | Street-stage, spiller | rest only | 300 | siritir alga. | Yest |
| | | | | | | | |

. 14

In the second of the second of

<u>Production of the second of t</u>

Takan Marian Bergaran Bergara Bergaran Bergaran

| • | * • • | : | • . | | (\circ) $\cdot (\circ)$ | |
|------|-------|------------|-------|--|--|--|
| | | | * | | , * | |
| | | | • | • | | |
| | | , 4 | | | ••• | |
| | , | - | • | · . | n 2 * · · | |
| , ii | | • | * . | # 1 A | . * | 10 m 1 m 1 m 1 m 1 m 1 m 1 m 1 m 1 m 1 m |
| | | | | | 4° • 4' | |
| | | | | | | • • |
| : | | . • | | * , | 4 , 4 , | - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 |
| • | | | | | * *** * * * * | |
| | | | | | | Section 18 Section 18 |
| - | | | | · 4 | | |
| * | | •• | | • | A STATE OF THE STA | Si ner |
| | * | | | | | ************************************** |
| | | | • | * 44 | # 7 # | |
| | | • | | | 9/6 / | |
| | | | | | 1 2 2 | , • \$. T. Co. |
| : | | 5 | | | • ' 💒 | |
| | | | | e _e | 100 | • 45.54. |
| | | | | £4. | | |
| | 5+ | | | : 7 | \vec{P}_{i_0} , | |
| | a | • | ; | ** * | d | and the second |
| 4* | | | | · · · · · · · · · · · · · · · · · · · | + Edit on the # | |
| | . 4 | | | 1 | the last of 150 to | |
| | | | | | | |
| | | • | . * | | | |
| • | | ga Marie . | 47.00 | in the second se | 1416 4 418 | |
| | | • | | | | |

| | | | | Sea | Harbor | |
|---------------------------------|------------------------------------|--|--|--------------------------------|----------------------|--------|
| Islands | Murres | Cormorants | Gulls | Lions | Seals | Puffin |
| Yaquina Head | 3,800 | 185 | 350 | sink was tops | 19475 ** 500 | |
| -Otter Crest Island | 30% SAN SEE SPACES | 90 | 460 Will 1986 | space whose values | April - Great | pine. |
| -Gull Rock | 2,000 | 400 | deer som one- | non-sam rings | 35 | alia . |
| Two Arch Rocks | 8,000 | 550 | STORE SERVICE | mate of the cutor | apin dile | , Rept |
| -Three Arch Rocks | 80,000 | 250 | ajost alkife pina | 55 | makes plants | wat |
| 7 Haystack Rock Coperfice and a | | dan ten ann | 1,200 | bijn gega gale | the stee | *** |
| - Cape Mears | John Stay Sept. After Stage | gija Apat nom | that the state | first size date | drive; disse | Side . |
| _Pillar Rock | 8,000 | 180 | dare data. Ares | rige data ibab | dans dobje | 1894 |
| Pyramid Rock | 1,800 | abot 1995 raw | violet dager dager | other-date laster | ager signs | *** |
| Jocky Cap Island | since divide little days 1904 | Higher standar mining | 300 | dang, sebah pengh | Mint the | ** |
| Haystack Rock Carrier | Tight that bus size often | special regional species | 600 | TRICK NOW ARM | eath and | 4 |
| Bird Rocks | 8,000 | 1,800 | Addit films faire | 9-40 State 10% | N= =0 | 24 |
| Sea Lion Rocks | and all the state that | 700 | and finituals | 80 | Magi. (889) | Make |
| Tillamook Head Rocks | design simple colors which address | 800 | sesse state, size- | stat des upp Mar | Marie State | 5000 |
| Light House | \$500 water state space space | 150 | disk hisk-inee | 4000 seles virgis | when the contract of | spine |
| | | distribution of the state of th | etzi-idifferendinyan da janta | application of the contract of | - | |
| TOTAL | 186,100 | 9,010 | 5,500 | 1,815 | 80 | 125 |

^{*} Birds (Puffins) are present in unknown numbers.

Raymond M. Glalm

CC:
/ William L. Finley NWR
Devid B. Marshall

| ·· . | | | | |
|------|-----|-------|---------|--|
| | : . | ** ** | My Mari | |

| | • • | | | | • | |
|---|-------|---|----|----------|-----------|--|
| | | | | | | |
| | | | | | | are out |
| | | | •- | ~ | | · |
| | | | | , | | |
| | | | | | ** * * ** | Section of the sectio |
| | | | | | | |
| - | | · | | • .; | • • • | |
| | | | | | | |
| | | | | | | |
| | ** | | • | | | |
| | * + # | | | ž | | |
| • | ** | | | | | |
| | • | | | | | 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1 |
| | | | | | | |

1 -

Observer Free Lvenuen, Jr.

Hornec Puffin

number of field trips on which our rved:

Fratercula cornical ta

Location in coseivitions

May 18, 1947 Ital Rocks, Lincoln 60th Found one dead & washed up by the tides.

| | | V . | |
|--|---|-----|--|
| | | | |
| | | | |
| | • | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

Observer Fred Lvenden, Jr.

humber of field trips on lie observed:

Turted Purrin 2389-3569&&


Mana cirrhat

late

Location and observ. tions

Way 18, 1947 See Necks, Lincoln Co .- found one dead, and washed up by the tides.

July 6, 1970 | Cape Meares, Tillamook Co. Oregon. Ad and young in water. Ad. and downy young at burrow on grass-tussock slope on mainland.



1-4-7-56-64-65-70-94-118-135-145-154-161-166-179-192-215-228-242-243-Band-tailed liceon

locatic and offer tions

oodcurn, fuga ntin com. Prince especiation of leader Five evaruous. It looked almost twice is large a first current of their bill nouse the were a pull room of implicable. In the first of incise bill nouse the rch 19, 1941 rour but of to but. There ice to talk to hit in appropriation, reg. Upper body necessors to be blue-green or wildlim action.

recrim retry, we need to be subscribed in the control of the vicinity of C to, are on the recording to the responsibility of C to, are on the recording to the responsibility of C to an area of the responsibility of C to a second the responsibilit pril 20, 1941

occurry for mot and 5,t. 17, 1841 one is the time.

Sept. 19, 1942 St. Louis, Ore.: Saw large flock of Band-tailed Pigeons heading south across part of the French Prairie country between here and St. Louis.

Notember 6, 1942 Port Townsend, Mash. A flock of 30 or 40 of these birds were feeding on the berries of the Macrone (Arbutus menziesii) this atternoon. Many of them would get some fruit in their mouth and then fly to another tree to eat it.

Moodburn - I me - the socitary Cand- intel Pina it in it is now in new and March 15, 1943 and then flew into awalut tree . It did not seem to fly far away, severingly, it had as the entretoner to refere the required sen a vary rind, may racled its react seawings

Corvallis-Reoria Ferry. 630 a.m. We saw a flock of 60-80 of them in a 1 kay 25, 1946 grain field beside the Pacific Hiway, as we went out. Al. took to the air as we approached.

Stewart Lake 8 a.m. Ora pair were observed per hed quietly in the top of an May 1, 1947

Woodubrn, Pudding R. - flushed 32 from the top of some firs in the woods May 3, 1947

May 10, 1947 2 miles north of Corvaelis, 5 p.m. 6 of them came down a valley and lit in the top of a clump of oak beside the creek, in open area.

May 13, 1947 - I saw a single, and later a flight of 15 to 20 came over the swamp,

circled and dropped into the field next to the FR tracks. Compared Santian River, Foster, Linn Co. 2 pair in a l. Fir clump of 5 trees, near May 24, 1947 an old orchard, where they were probably attracted to fice on chemies.

May 27, 1947 12-mile Swamp: 0 4 flushed iron the ground near dry old creek bed.

Milk Fanch, Spiqua Creek - Thushed 4 or 5 from a fir chump on open S slope. May 31, 1947

MacDonala Forest, Saadle area figurhed 4 from top of a L. Fir.
MacLoland Forest, Soap creek side - nest a bu feet in L. III, on L7 sade
Clembroom, and vicinity, Benton Coff by .n. on one out hilledge one over June 30, 1947 July 2, 1947

Aug. 22, 1947 40 birds in the tage of the order, and they are a derose to account another wooded area. They have been hore desping, odel to all these to see

Bradley-Jolon, Hunter-Liggett Reser vation: /10 a.m. large flocks in the Jan. 29, 1949 tops of oaks, and on the ground everywhere on plowed and stubble fields. In a strip about 2-mile wide on each side of the hiway, we saw at least 2500 birds in 10 miles of road. Also noted that often the flocks were on the ground only under the scattered oaks. Warren Smith, the local warden has had to issue many permits to satisfy the ranchers who complain of their wrecking their crops.

Sacramento-Gray Loage Refuge for t least 500 seen in small flocks in rielas Feb. 6, 1949

Sacramento by flying over State Fair grounds.

3 pm. Weed + Dunsmir, Calif. July 31, 1949



. .

Oct. 3, 1949, Huckleberry Hill, Lincoln Co. Oregon. 8-10 around cabin, mostly perched in tops of live and dead spruce, and few in locgepole pine. On Oct. 4 ceiling zero , R- and S 15-25 m.p.h. when about 20 pigeons were perched in one spruce and 2 spruce snags from 8 a.m. till at least 4 p.m. No Less than 6 were seen at these perches during this time. Rain didn't seem to ëffect them.

March 22,1950 East of Smartville, I hoba Co, Calif - Saw over 100 birds about 8:15 a.m. flying in small groups of upe to 30 birds in this area of Ligger jeine - stagarral.

Jan. 5, 1951 Hogan Res. Calaveras Co. Calif: 350 seen - first time they have been

seen in the reservoir area in the 10 years Mr. Ballard has been caretaker there. Game Warden states that they are in by the 1000's near Murphy's and Railroad Flat, further into the mountains.

March 19, 1951 Hogan Res. Calaveras Co. Calif. Mr. Ballard states that the Jan. pigeons hung around 2-3 weeks.

June 2, 1951 Clear Lake " inglas o off 1101 - 3 10 v. a. . 2. 7. Many flooler Hound will in area

Vicnty of Bass L., Eldorado Co. Calif: flock of over of in digger May 5, 1951 pine towering above live oak woodland on N slope of ridge. When flushed, kept returning to this tree.

Mar. 13, 1952 Uvas Cr, W of Gilroy, S. Clara Co. Calif. 0840PST. 150 feeding on olivesothers seen perched in digger pineand oaks on hillsides of this area. E.a Hurdren, group the Bor to Minning in I were. Have.

June 25, 1954 U.S. 50, E of Fresh Pond, 2 miles, Eldorado Co. Calif. a pair fir area. May 10, 1954 Asilomar, Pac. Grove, Monterey Co. Calif. 2 seen on TV antenna.

Sept 23, 1955 Shada Cty, Shata & C., 439 pm. pr.2-4,1755 Pacific Grave, Calif. Places of & to be present.

Dec. 30, 1953

Mark 12, 1957 Cottonwood, Caly. Many seemin fields of bottom lands.

SEPT. 25, 1957 E of El Carrier (15 10 El El Corres (15 10) april 23 1958 20 miles N of Monticello Reservoir in Lake to, Cely 15-20 in Jine - ook - want area.

Sept. 25, 1958 - Bucks Creek Powerhouse, Feather 3. Canyon, Storrie, Colif. - 35-40 Sept. 25, 1958 - Jct. East Branch and M. Work, Weather R. near Belden Galif. 6 seen.

Sept. 25, 1958 - Indian Valley, UShOAlt. Plumasco. Calif. 25-30 secn. Sept. 26, 1958 - Duincy - Portola Jct.. UShOAlt and OSH 89 - 15-20 secn.

Sept. 26, 1958 - Blue Canyan area alone USho, Calif. 35-40 seen at 430 p.m. POST. June 11, 1958 - American R. Sacramento, Calif. over 140 norched in tops of dead dottonwood by river at R a.m. They were inactive.

ang 21, 1960 Nof Polled Flat, 4.5 99, Electe Co, colf.

Feb. 75, 1963 9 of Welf Creek \$1999 9w Organ - 11 over-living
Harch 10, 1962, Gateway Retroat "ouse SW of Golfax, Macer Go. Calif. 27 O'el. Perhaps 1,500 or more in pairs to 40-50 per flock from east up out of M.Fk. canyon to SF of Golfax. Same flocks were flying back toward the canyon at 4:15 p.m. while it was snowing lightly as we left for Sacto.

6/11/67 - Victorie to Port Kenpen BC Parelet on with lines along road!

| | | | · | |
|--|--|---|---|-----|
| | | | | - , |
| | | · | | |
| | | | | |
| | | | | ٠ |
| | | , | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

Observer Fred Evenden, Jr.

Hock Tove

Tumber 1 Micle trips on 1 ien obt styck:
101-105-116-139-161- 275-339-370-1309-1519-1524-1528-1588-1588-1579-1676-1691.

Columba livia livia

White winged Dave 234

Late Location and opportunions

Red-billed Figeon

June 20-21,1964 Santa Ana Mar. Hidalgo Co. Texas. A seen, 1 on nest. Several calling. 3 at Pentsen-3. Grande St. ark.

Thite-fronted Dove

June 20-21,1964 Santa Ana NVR, Hidalgo Co. Texas. 10, with one on nest.



ne 10, 1941 - | est obdition: I notice to both in a real in . 19 17 housel cor chosy white we so tail a obesit. I consider the first on a conoodburn, Pra, zant's monas: was one since say over as ept. 17, 1941 Colvandi of them, no classically the color of the velocity un. 26, 1942 Fig. 1. Source is now to keep the albinot.

Fig. 1. Source is now to keep the line of the -st . Oc 13: 01 1 Fig 18, 1942 May 11 1944 Mtn. Home, Idaho, Cromwell's orchard. 2 pairs in the orchard. Found only one nest about 12 feet high in an apple tree. It is of twigs, and has a grass stem lining. It is shallow, the cup being about 1" deep and 3" in diameter. It is poorly constructed, a person being able to see thru it quite easily. 2 eggs in the nest, a creamy white color, and 3/4" x 7/8-I" in size, both ends rounded equally, but one end seems to have something dark, inside. The surface of the egg remind me of a "china egg" used under setting hens. The doves did not return to the nest when the rain began. May 17, 1944 Mtn. Home. One bird on the nest containing 2 eggs (took lic of eggs). May 28, 1944 Mtn. Home Area II, 5 p.m. Two young ones were in the nest when I flushed the adult from the nest. I took a picture. They were of qual size. Black skin. The cranium part of the head seemed quite enlarged. Large bill for size of the bird. Almost "tube-nosed" in structure. Body was covered thinly with a line brownish thread or hair-like fuzz about 4-3/8" long. Body about 2" long. Head 3/4" long. Mtn. Home. The two young mourning doves are now og" long. Have black skin May 50, 1944 Their body is still covered with tan fuzz, quite wiry looking. They have very fine and slender wing quills about 1" long. Mtn. Home. The two mourning doves have left the nest, but I saw them a June 15 1944 little later when they flushed from the ground in the orchard. They seemed to be > to 2/5 grown, now. Apr. 11 1946 Corvallis, Oregon, Brook Lane - Bernard Smith, Jim Thrailkill and I flushed a Mourning Dove from a newly made nest in a young ash tree, growing close beside an old tree (ash). The nest was made of grasses and small twigs, and was a shallow cup affair. It was at a height of 6 feet above the edge of the pond, and over the water, as well. There were two pinkish-white eggs in it, evidently laid within the past day or two. The nest was on the SE side of the big ash, and therefore sheltered from the NW and W wings, etc. The bird flushed from the nest flew about 50 feet away and lit to watch us. After we left, it returned to the nest and stayed on it, even when we passed near it on the way back. End of Brook Lane 18 a.m. I went out to the nest forma on Apr. 11th, to May 3, 1946 band the young. As I approached the nest I saw the adult bird (just one seen) perched on a small limb at the eage of the nest. As I approached it flew down to the water level, and proceased to put on the "wounded act" and its wing tips and tail dragged in the water, and it gave a hurt call note, as it ilew a short distance away. On the nest were two young doves, approximately 2/3 grown. They were both facing away from me, and then I came within 3 feet of the nest, they both took out or there as if they had been flying for years! One flew across the pond and lit expertly in a clump of willows about 35 feet away. The other one flew over 50 feet before it lit in a tree. I would assume that this was their first flight

5⁴7.5 . .

May 3, 1946 - continueá:

from the nest, and it was surprisingly so ong. No one had to teach them to fly, from the looks of thisgs. The nest cup, was piled full of clumps of feces, and they were in two more or less distinct piles at that.

Peoria Ferry Road, 2 miles south of Owl Creek. May 9, 1946

Found another nest in an ash tree, at about 10 fet. Was over water, but is over land since fater receded. On east side of tree, so does not get p.m. sun, directly. Same as nest on brook Lane, except this is one a horizontal limb, rather than in a tree. There were two eggs inot.

August 13, 1946 - Leep Canyon 6 fisters to Leamona 5 p.m. Arthough the canyon has been burned out, and there is very little there, Loc and I found that there was a very large flock of 50-100 love; , both adult and young perched on the fire-stripped tiess.

August 21, 1946 - Poison Oak Lake, Corvallis Long young birds present everywhere.

Sept. 3, 1946 - Goodburn to Portland Many paired up on the wire kines.

March 20, 1947 - Colo. R. Indian Res, south of Parker Ariz: 7 and 200% in one field with Redwings, Brewer's, etc.

April 24, 1947 - Poison Oak Lake, Linn Co: one bird perched 70 feet up in top of old oak, cooing loudly. Later I flushed another bird in the poison-oak-pasture area, and found a nest located on to, of a very old D. Fir stump, rotten, but dry. It was 24" across and the top was 22 feet above the ground. The stump was encircled by poison oak 6' high, except on the east side. The nest was in a shallow cavity on top ofthestump. Nest of heavier twigs up to dia., and lined with very fine woody stems. No grass. About $1\frac{1}{2}$ " deep and $3\frac{1}{2}$ " wide on the inside. 2 young were in the nest, 1 a little bit largerthan the other, but the larger one was dead, and it appeared to have a broken neck. The other was about 2" long, and covered sparsely with a grizzly gray-brown hair or down. Skin gray-black-brown. Eyes did not seem to be open yet.

12-mile (wamp, prentifur in small frocks - family groups?? June 17, 1947

"Poison Oak" L., Linn (8 - 2 pair in the area. Lushed ore naurt from the Aug. 13, 1947 vicinity of a neet located in a " bison oak" parture. The sect has situated ato plan (la l. riv stump about he" righ, ha to a surely our illoot NF of the stump, and a clump of poison oak shadin it from the C and . The nest cas a shallow saucer with a fer arise thigs placed in it. ' was one young that will be ready to lay in a ready to lay in a ready use its wings weal. About 5" long, with a general gray-crown couper. Primaries ab ut all long, tail reathers 3" long, with in whiteouter reather already showing. Even unen it beat its wind and I jience it up, it wast its breast "inflatea". Bandea | 41-354648. Ferfale & ve "woulded act" when she left the nest vicinity. This nest as located not over left from the I found this must spring atop another L. His sturp. Could be the same nesting pair, I'd guess. There was also one dad ego in the nest. The nest stump was about 75 reet from the water in the game.

10-mile ample several pairs in the real later and a taken on the north side of a pline change at another at the sect of an esh or we. The next as on a borizontal I'm ash limb, that a consequence of the states, and about 5" acr . . One accept on on the mont, in them early store west was vol -shacco.

descent - located in some a cod, that a process a board of the roughania buththat provided limity of these for the sest, sie sum of grasses, and 5" across. The rist was on the Grisley a trader, but outlet to we of damy. The nest or only I "do we the date: There were two "mare-covered" cark-shi need to binds rabban will the died-opside racing to the FI. Total larger to the part of the state of the second yet, the shart, here over 18 den. Bune 641-0040 can 45. The spined " · . + 751

June 15, 1948



- June 18, 1948
- 18-mile Samp: Plault and jour have alleady left neat.

 11-mile Samp: Prevenue pairs seen next in no use i continction. June 21, 1948 last week conjunted now, but no eggs in it. -ours not W just west of cattail clamp on dead branches of Spirot at 1' over them, opening to the all as or nest containing journ that fit the deat containing journ that fit the deat containing ned net had been.
- Farmington Fes. Site, Stanislaus Co: many in area (5-200). Hest on debris May 19, 1949 in crotch of willon on west side of creek, 4 feat above the water. 2 clean whitish eggs in a shallo saucer. West made of dry grass stems. bull willow canopy over it.
 - Woodland Willows = many pairs 5:3 opm.
 Colusa grilley, Calif clocked at 38 m.p. h. for y mile at 1:15 pm P.S.T. 7-29-49 8-16-49
- Millsaps Newville, Glenn Co. 4 p.m. about 6/road mile. Nov. 15, 1949
- Mar. 20, 1950, Red Bluff-Los Molinos, Calif. 730 a.m. 1-2 pair/road mile
- Apr. 17, 1950, N of Red Bluff, Tehama Co. Calif. 545 p.m. Had not seen any in mountainous areas north of here, even at Reading, but from Cottonwood on south to Sacramento, there were quite a few of them.
 - May 19, 1950 Farmington Res. area, Stanislaus Co. Calif. 930 a.m. 8 along hiway plus 45-50 in one open grove of trees. Adults and nearly-grown young were
- June 30, 1950, Fugene-Milton, Calavet as Co. 10 a.m. pair pas ed us going downhile at 42 m.p.h. 35 seen between t end two paints.
 - Nov. 10, 1950 Feather h, r of piges / da 1 " in croplanes, to 10 no 25- in tundy south exposure among greaged rock piles in mined area near river.
- Jan. 5, 1951 Stockton-Farmington: /1 per 2 road miles.
- March 10, 1951 Gila Bend-Tucson road, Arizona 5-6 p.m. very abundant along this hiway, especially in the cultivated areas.
- Vicinity of Bass L. Eldorado Co. Calif. found one nest on horizontal May 5, 1951 southside limb of Buckeye at about 10' elev. Pin-feather young it it, dead.
- Abandoned orchard, N o f Sacto. Signal Depot + flushed adult from July 3, 1951 nest at 7' in pear tree at 6'. One egg and 1 2-3-day-old young.
- E box backers Pass (15.4.152. 4 p.m. Mouring Dave Rest up with us for July 22, 1951 at least if mile at 62 m. p. h. - attill wind
- Sept. 7, 1951 NE of Anderson, Shasta Co. Calif. In rocky-soiled, blue-oak woodland there were few doves seen along the fences.
- Jan. 28-29, 1952 Many seen on forces near grain stubble + weed pateles N + 5 of Sacto. Colif. Gray Lodge Refuge, Butte Co. Calif. Cowan has placed wire baskets in May 1, 1952 trees on the refuge, and the doves are utilizing them for nest sites. Screening is about $\frac{1}{4}$ -inch mesh, and about 10 inches across.
 - Jan. 13-14, 1953 S. Brandara G, Giga Mo, would dover our surver and Jan. 29, 1953 - 4-5 mi. I of Tudor, butter Co. Calif. acced male reduing take perch at 35' in willow away from Mourning Dove.
- June 5, 1955 Sacto. Bypass, Yolo Co. Calif. nest at 181 in Willow. 10 adults seen.

 Nov. 27, 1955, Imbert rad front 1 of Countland, Tilis. The one in Francisco Instrume.

 Sept 27, 1956, 25-30 ad typy in week feeler 5. Land Park Hills #73 on around S.T. Sacto Colfornia 615 pm. 1957.

 June 17, 1957 In conferons forest felt along U.S. 50 east of Pollock Pines, Eldordo Co. Oly. 10 am.

 Tay 21, 21, 1957, 100 and 100
- May 36, 1959 4 Fam. PST 5 of Folian Soute to, Calif. active before any light ruly good. Back toth across roads July 3, 1960 National state Excession and the state, state in the state of the state of
 - August 1960, Hany resort in Allerette Talley wille in into Defileration
 - 50 28,1961 april 1 / hand Electric Gold of 18 'Aug.13, 1960 - Yolo-Solano Co. Calif. M. Dove census route-65 miles. 72 doves (90/100 miles) . July, 1901 - M. Leveb three $\frac{th_2}{t}$ magnifest bolders for Lorentz syllength and Lymin $\frac{t}{t}$, $\frac{t}{t}$, $\frac{t}{t}$



Observer Fred G. Evenden, Jr.

White-winged Dove Zanaide

Melopelia asiatica mearnsi

Number of field trips on which observed: 234-1841-2622-2603-2603-2603-2603-10-264-15-17-18-2620-21-22-

Location and observations: Date

Feb. 18, 1951

Parker Dam, Calif. One perched in Athel three near B. R. residences below dam. While perched the white wing edge and white areas under tail showed OK. Its cooing notes are very distinctive and clearer than mourning dove.

June 20-21,1964 Santa Ana MWR, Hidalgo Co. Texas. 1000s present. June 23, 1964 Bentsen-R.Grande St. Jark, Fidalgo Co.Texas. 10001s of birds, 2eggs/nest, and almost ator each others. Anis present.

111



Inca Dove S cardafella inca inca

ground Dove - 2618

Feb. 17, 1951. Colorado River Indian Agency, Parker, A rizona: 530 p.m. 8 seen in figs and pecans on the residence group of buildlings. Chestnut wings in flight. Have distinct 2-syllable note.

, S ·

.

California Cuckoo

1657-7770-

Yellow-billed Cuckoo Coccyzus Eneric hus occidentalis

Late

Location and observations

July 17, 1950 Mendota Pool, San Joaquin R-Fresno Slough, Fresno Co. moon hour. Heard a strange came from a willow thicket along Tresno Chaugh, and stop, ea to Look back, for we had just passed it, and then I spotted a bird flying past us toward that thicket, and I did not recognize it at all. Finally several other views of it - gray body, chestrut winds, whitish under body, black on near, undertail black and white spotted, etc. identified it for us. The birds gave low, gutteral scole notes. Before reliving the area we had seen 3 pairs, some rainly close to us as we ate runc . In willowcattail slong border strip.

May 3(,1)64. June 20,19'4

Pocomoke R. S of Delline Worchester Co. Md. nest 7 10 with he hale blue eggs. Santa Ana Refuge, R. Grande R. Texas. 12's in 1s and 2s. 6 eggs in 1 nesta6'. "To Suckeshilliese Zearn, I demonique force just on your fait, w/ wings is cooping

of Carolica ... 6/8/76 - Carolina - Ca



546 - 557 - 1052 - 1058 - 1063 - 1140 - 1203 - 1423 - 1591 - 1764 - 1815 - 1824 - 1838 - 1839 - 1845 - 1846 - 1857 - 1860 - 2037 - 2113 - 2313 - 2314 - 2391

2596-2609-2618-

Road-Runner

Geococcyx californianus

March 20, 1947 Colo. R. Indian Reserv., south of Parker Ariz: 2 crossing the road, slowly, pausing at edge. 1st one seen was on a fence post and did not fly for a moment. Glided to the ground, ran, and flew to telephone pole crossarm at 20 feet height. Opened its bill and gave its "rattle" notes (by grating its bill). Crest flexed now and then. White and red patched on head hehing eye plainly seen. Whole body streaked brown. Tail dark brown or black. It was very obliging, and did not fly when we squeaked, whistled, waved, or pounded on the car, nor until Gale got within 8 or 10 feet of the pole!!!

Reprinted from The Wilson Bulletin, Vol. 58, No. 3, Sept., 1946, p. 185

Road-runner preys on Poor-will.-On March 9, 1943, around 1:30 p.m. as I was driving along the trail about a mile northwest of Tule Tank, on the Cabeza Prieta Game Range, Yuma County, Arizona, I noticed some feathers of a Poor-will (Phalaenoptilus nuttalli) lying in the road. A short distance down the trail, I saw a Road-runner (Geococcyx californianus) running along, carrying something in its beak. When I gave chase, it dropped its burden, which proved to be the still warm carcass of a Poor-will, intact save that most of the viscera had been removed (and probably eaten), and a number of the larger wing feathers had been torn out. I find no previous reference in the literature to Road-runners preying on birds of this family.—Gale Monson, Fish and Wildlife Service, Parker, Arizona.

Nov. 5, 1948 - Dredge area, ½ mi, 5 of Fair Cakes fet. with U.S. 50, Lacto. Co. 19.

Jan. 3, 1950. W of Stony Gorge Res., Glenn Co. Calif. 1145 a.m. One on the hiway just west of the reservoir, and as we approached it flew off the road and acress the fence and then leisurely oved away from us, across sparse grassland in rolling hill country of scattered blue oaks and buckbrush - there were no color natches on its face.

May 16, 1950 Stony Gorge Dam, Glenn Co. Calif. 5 p.m. PLST. two of them at the south butt of the dam, getting drink from the reservoir, which is about full.

• • • • •

•

Eurn Jul 2580-

Jan. 15, 1963

Tyto alba pratincola

Location and on a votilize Colvarior Ly 8, 1942 12-mile Swamp, south of Corvallis of I found a light, but richly colored Apr. 24, 1946 individual in some scrub ash in a thicket out in the swamp area. It flew and Reawings chased it all the way across the swamp. Decramento, alif - 730 p.m. ne /2jing et 30' von J. 2" war" 1:0. 10-7-48 Sacramoto - 8p.m. 2 flying about reafof state Office Blog #1. 10-16-48 - 9/5 pm. - one flying at 30' & w.on I St., sast 10.0. again. 10-29-48 Near Verona, Sutter is, laid - 7 p.m. One flighty along hivay. 11-25-48 Sacramento 10 p. m. Dais flying around vicinity of 16 + 14 at ht. of 150'- 1751. Each time the beam of an arc. light apostlight againsed the seley over them, one 1-32-49 would give out with a " light - squark by. Mar. 15, 1949 Leaubetter Penin. Mouth of Tillamette Fiver, Portland, Ore. one flew out of an old barn down river from the rookery towards the nouth. Te were not able to find the bird again, and it was not present when we returned to the barn about I hour later. Could not find any signs of a nest in the barn and only occasional arospinss, which I don't think were of an ora-1030 pm. Lacto - 2 228-3 of ave - One calling from oak intackyord. april 23, 1949 10 pm. Klamatt Falle, Oregon. July 30, 1949 N of Williams, Calif, 930 pm. on fence poot in rice are. Cotaber 8, 1949 Nov. 22, 1949 garden Hiway, near Elkhorn Ferry, Sacto G, Calif - 5 5 jun Jan. 21, 1950 Nedgo of Stockton, Celif., US 99. Mar. 14, 1950 Dec. 28,1950



Observer Irea Lvenden, Jr.

Number of Field trips on Fich observed:

Racrarline's or on al

ocus asio macíarlonei

| iate | Location | Jr_C. | observations |
|------|----------|-------|--------------|
| | | | |

•

Otus asio premeter. Location and observations Woodburn, The past few nights I have heard the melogious and hollow Jan. 27, 1946 "bouncing-ball" call notes of this bird, in the trees by Giels and Woodburn, 1930 PST., I rode the bike around Gill's block to triangulate Feb. 15, 1946 the location of the owl, giving its call periodically. I deduced it must be in an apple tree behind Gill's house so took the flashlight and went to see. I made lots of noise, breaking twigs, etc., but it still called. I saw its form against the sky in the upper limbs of the tree. The full moon was out, making things very bright and easily seen. I put the light on it, but it wasn't bothered. It went on calling. It's eyes shown a deep readish-pink in the light. It turned its head occasionally and while it looked the other way I "sqeeked like a mouse" and it was comical to see it jerk around all of a sudden!. It almost lost its balance on the limb. The bird was quite light gray and tan below. I could not see the back. The head seemed light also. I went elsewhere on my bike for 15 minutes and when I hat returned it had flown to a pine tree a black away. I heard it so rode over and spotted it again with the flashlight. It continued to give its call. All evidence points to it being a male, for books say females are a reddish-brown, much darker. The males being grayish colored. Feb. 18, 1946 Woodburn, or Tongisht our Screech owl really went to town outside. It hardly paused imnits tremelos. This persisted for several hours. Woodburn, The Brewster's Screech Owl is still "tootling" rapidly and Feb. 25, 1946 often every night. Feb. 24, 1948 Corvallis, OSCCampus - 930 p.m. One giving about g of the full call, at 10-15 second intervals from tall old fir near Bandstand, about 15' up. Feb. 26, 1948 OSC Campus calling every 20-40 seconds from same fir tree as Feb. 24, and in about the same place on the tree at 20-25'. woodburn, 11 p. m., 6.5.T - One calling almost continuously from apple tree across way July 11 , 1948 August 1, 1948 onchum, Silve, et e 🗡 an. In the veni of in alm to the in Woodburn of p. m. One giving regular t low tremelo call from neighbors walnut stress = 2 moon august 11, 1948 Woodburn, home. one heard within the past 3 nites about 930 p.m. Aug. 18, 1948 1:30 a.m. Woodbur, home . One calling from serves the street . Heavy rain . Feb. 20, 1949 Woodburn Oregon. Mom tell me two have been calling every wite, lately. May 12-13, 1949 July 3, 1949 Modaburn, Oregon. Home. Mother reports hearing the owl off and on the past couple of weeks. I dien't hear it tonight till about 1030 PLCT just following a large fireworks explosion. Its call was weak. Corvallis, Oregon - nest in N& Flicker hale 12' up in head alder stat in Reavy Timber " Eggs on things " 12" . May 7, 1921 Corvallis, Oregon. 3 spgs 12' up in flike hole in the Good Mast - it offer Corvolles, Oregon - 3' up in Alekarhole in dead alles that Tiffee Corvolles, Oregon 3720' up in del flikes Role, in juice. that had 5 sparrow hank 2994 taken from I'm May 10, 1921 May 20, 1921 Corvallis, Orgon. 3 eggs 12'ngs in alder stul- Virifiel.
Indurantes, ore. - 3 eggs in hole indeed top of a maple, 20'ng. - satural cavity - Cuttress May 20, 1921 May 7, 1922 april 10, 1941 Milwania, C. 2. . 3 aga in bole 7'up in large read for stury. Hole day year Lyon by woodgester - (" May 5, 1941 Milwanker, Ote - 3 aggs on rotten word - no mest . 1 15' in dead string. I o april 17, 1943

Observer brea vender, Jr.

California Forescal

Otus acio bendirci

inte | rocution no control vian.

Number of field trips on which abrely a: 341-1542-1674-1790-2254-2326-



Observer Frea Evenden, Jr.

Flammul tea coreech Owl

Otus ilammeolu.

Location and observ tions Late

Tumber of field trips on which ous-twock: 426-7

N.C.+

Observer Free Evendon, Jr. Hornes Cal 2325-1326-24576- 2457 Eubo virginianus dei Montana - 1230-1232-Location and observation: Lute Coast S. or Lonterey: Towns many perlets in Lain, sense Lontere; sine May 21, 1949 thicket near Testern Flycate er nest. N. of Chiloguin, Klamath Co. Oregon - 1 Pacific. H. Owl Let along Riway 97 Lept 2, 1949 Lucia, Monterey 6, on C.S.H. 1. \$30 p.m. . 1 along . -. Dec. 6, 1949 garden Hiway, near Elkhom Ferry Lacto Co, Calif - 5 pm. Jan. 4, 1950 Way 13, 1950 Yosemite Valley, 6 p.m. one perched in yellow pine, with many Stellar Jays around so laing it - t was carrying a snake. From, with jays after it. . Cartal, Celf. 1 was at 645 go, n. new the Minnien. Just E of Morro Bay, S. Luis (Itages Co, Colf 5: 30 gov. One on place near rod. Still daylight. Jan. 14, 1963

May 31,1953 NE of al Tahoe, Alderado Co. Calif. Lest cavity at 15-501 on west side, huge Jeffrey pine snag. 2 downy yours, about 10" hith. Iso saw I adult. Fellets contained redents and cost, sparrow hawk, and flicter feathers.

Sacramento By-pass, Y to Co. Calif. 7 a.m. 1 abult in coltonwood at fot, Aug. 22, 1954 and 8' out from trunk.

7 mi NE of Weed, Siskyou a Calf. - One frosh killed along 4597 hartnite December 20, 1957

Observer Fred Evenden, Jr.

Forthwestern horned Owl

Location and observations

Number of Liele traps on which ees avec:

A STATE OF THE STA

Observer Fred Evenden, Jr.

lusky Horned Owl

sumber of field trips in dich our ived: 31-44-45-64-74-116-137-141-152-156-168-183-203-205-341-79?-3 425-426-508-512-589-593-613-775-847-851-861-876-931-10550-1186-4435-

bubo virginianus saturatus

Late

Location and operry ion.

June 12, 1941

Layrent'. To as at occurrent of pair on organistic aire par or worth

pril 16, 1948

Corval is take an y is a colonia lande of I le et. of our distinctions at a contrary empty recoped to a contrarty of Ill to be acce

pril 23, 1943

orvanicon (see contess of nation, en a to true in. The risk of the third the representation of the state of the second at i found on the and an ending out on the court of the sound of the loss in the tenton situation and the first being some some situations of the state tarresson is a characteristic of the conditions ${f v}$ 61, and ${f u}$ 61, and ${f v}$ 61, and ${f v}$ 61, and ${f v}$ 70, a there we to be a fill a side of the contribution of the contributi one or the rate of him link; (take and rain or table and rike.)

Oct. 1,1942

was a first to be a second Oboria from: Found a par watere sended water in the war die hary den the round the largestime of "x1" in his legen of the in it is the was absolutely no other tissue menont. The consider which is a the rest of the ..

Apr. 25, 1946. Dixon Creek Riage, north of Corvallis. Charley Cutress showed me the nest he located the week before. The adult was not around, but the young had gray-white down, with a few pin-feathers sticking through here and there. One I had a good look at seemed to be about 7" long.

Apr. 12, 1947

Woodburn, Pudding R: perched in dense cedar clump at 930 a.m.

mpr. 5, 1948

p-mile of cair Village, benton Co, & p.m. Charley Cutress took me to a nest he had spotted about Feb. 29 which had a eggs at that time. The "nest" is the open top of a snaped-off L. Fir for dia., open on top, and to the eart, but "vallca-in" to the north, west, and south. Beight about 25'. The snag was in a quite open site in old burn and to red-over area, with only a few scattered tall 1. Firs in the area, the coarest some low feet away. The snag was on a FPT ...ope, slightly sheltered by the rioge to the courth or it. The young hatched March 7, 1948. e could spot the nest over --mire away, and the Temale was in the nuche with the young, but left pefore we were 1/8 mile away, at least. It stayed in the 4. Fir woods meanty and made no sounds until we got to the snag. Then a few "who's" were enitted. As Charley went up the reader to the nest, both perents tarted giving various scolding and call notes, one of which but like a "bark"; another not inlike one note the Brick-bir ed Lagpie emits. This they chattered their birs, and the young bird gave a deeper cracking noise with its bill as we brought the bird to the ground. Its eye sight seemed guite poor, and it observed only bright things in front of its eyes, and fast movements not over a foot or so away. Tarsus was almost ful ; developed, and down-covered. Tris, bright yellow. Contour reather developing, and principles to ke" of reather vaene already exposed. Much "powder" precent. Length about 10-12". Danded it with size 8, #44-812061. oth parents were in firs less than loc feet away, the big cark male with drean white clin patch usually quiet, but occasionally "who-ing" a little. The female was not quite as large, and was almost a

| | | . • | ٠ | |
|--|---|-----|---|--|
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | · | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | • | | | |

gray-tan, rather than the cather gray brown of the mane. The gave the barking couls previously mentioned. Tuch fecult was on the snag about the top of the most, and a few recently =acquired pheasant feathers were caught on the sides of the snag, and on the grained around the snag. It was a new bird. He ut 50/ feet away to the cost was a most in the crown which an acult bird in at roost, for numerous palaets were on the crown around the mest. The more recent ones had not it into an the round of a pheasant in them. It ere the crosses and not return directly to the most was a part of a pelvis. The worlder and broken ferur, and in another was a part of a pelvis. The worlder aid not return directly to the most one described same cay the first of the firs

April 5, 1948

mile N of Maair vil and two is one by it too muc. That one to hill to the south of Moffin atte we spot tea an old Labair Pava nest 40% feet up in a Dig-leaf Tople (bucs just broaking), on an open on the Lage, not over smile from the risk way. The tree stood by itself. Charley chapsed his hand, and a bire stretched its sings and tail sand the horned out sit they off in sently. The into the out, or top of the hill to the wist. I climbed to the mest and lound, unfortunity, two esgs about 10-th cays incubation time. The bad, I enter to alma them, eleft them, and went on ever the hill, having not attributed toth of the outs, only one carling occasionally while I has up to thee.

About 15-L minutes rater we returned, and the out of team of the nest, the we had taken the eggs.

May 4, 1948 12-mile damp. In the earn south of the luce, Or. bridge. Indeviously being chance and attacked by a crow. There has been a join of their draw withis area the past month of so.

Apr. 30, 1949 3 mi. W of Bruce Sta., Benton Co (FBM). hooting at 520 a.m. 170m oak hill-side.

august 21,1949 5 mi. S of askland, Jackson G. Orgon on U.S.99 - hit at edge of road.
8/15/61 - 71. Klamoth area, Klamoth Co, Orgon - 102 penetree ottas in saled

Western Horned Owl

great-1855-

Bubo virginianus pallescens

March 19, 1947 Havasu Lake, Ariz: 2 flushed from a bluff below Blankenship bend.

May have nesting site on this bluff wal over the lake edge. We flushed them by approaching in our cabin cruiser, about 6 p.m.

Feb. 27, 1953 N of Encino, Torrance Co. New Nexico - at dusk.

apl 1/19 The Die Nessey, Rockenitabild. I figure + model in Come forth.

| • | | |
|---|--|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |



rate

Location and observations

| arch 20, 1946 | Rock Creek, near Philomath, Ore. 4-530 p.m. This one was perched on a dead snag about 60 feet above the ground, near the creek. The sky was fairly clear and the sun was bright.

May 21, 1947 S. Spur, MacDonala Forest: as we started out at 6 p.m., Aller spotted this one perched on a phone line pole by the road. This was on an east slope, and had been in the shade for a long while. It didn't fly, and I got within 20 feet of it, and finally after waving my arms it fler in a long downward gip, and an up-swoop into the top of a Pig-leaf Maule 60 feet away.

aip, and an up-swoop into the top of a Pig-leaf Maple 60 feet away.

Lec. 6, 1947

Also in a finite of the top of a Pig-leaf Maple 60 feet away.

Also in a finite of the control of the c

Jan. 4, 1948 in the state of th

Feb. 23, 1948 Sunset Hill, Corvaliis, 10 p.m. heard one calling from old orchard rea - several "hoos", enoing with 3 sharper, and more distinct notes. Loud, for its size - it made a neighbor dog bark.

Dec. 29,1953 WolfGreek word at Humbur Greek, west. (ne pendich in aller over biss. Pics.

Elf Twl -Tiorathene whitneyi

June 20,1964 Bentsen-K.Grande St. Park, Widalgo Co.Texas. 7:30-0 2 young in Pulsache tree - "soft chuc'le" colls. Imm? one dork, lighter.

Sed of the second of the secon Since the solution with the solution of the so

Late | Location and opening tions

cb. 22, 1941 | Near Peorit. This material with a in motor, high then in a rich .

About A Ball to all Evaluitions. March 6, 1943

us take i living i to the same spot as a tollow as a comment we we can be a finite, as well, finite over a significant incorporate constant xused a range time. I now my with to atmin a last on a recover RO TROUBLE LICENSE SECTION OF THE RESERVED AT THE COURT OF THE Low to the ground show or he ash a capto court, or a just his telestrated at the constant of t

Corning-Orlando - one perched on 15: crossarm of telephone pole. Pale color Aug. 31, 1948

Sept. 26, 1948: River Park, Sacramento: 130 p.m. flushed one at euge of stubble field and sand lot. The bird was in a small area containing several burrows, presumably of ground squirrels. Tame.

5 mi. I. of Coming, tehans co 12 2 m. - a pair on open know in open country april 6,1949 about 75 feet west of fonce along Riway.

5 mi. 8. of Corning takama 6 1.30 p.m. - Pair seen on mound at same place apr. 8, 1949

Gray Loage Beruge, Eutte Co. pair at nest hole near gate to reluge - 130p.m. pril 23, 1949

Gray Loage Keruge: deir still occupying burrows along tence near gate to retuge May 5, 1949 Near Orland, Calif - 2 adults on fence posts + 3 yng at mouth of hurrows. about 'z adult size. June 12, 1949

July 11, 1949, N of lun isan in Coluse Co. Calif. West burrows in canal cike. remains of water that and pri-colored recylines bout of enings.

5 mi S. of Corning, Tehama Co. 16'5p.m. 2 adulte at same spot as Gov. 6 +8.
5 yng perched on fence posts of wires 100 yds. from adults burrows. They were July 31, 1949 flying occasionally.

745 pm. I on telephone sole between Valley Home + Callegeville, San Josquin Co. ang 3, 1949

Aug. 13, 1949 6-mile unimproved rold, har it of terverton, colono Colon para just west of Lixon-lio Vista bold. Son swithin wiki, the operation top rails or proliferenced that assembly fence. They for the Late, see within 8 is toi them, but on hourt care nock and lit by sait less cheking its body and it move a seclicing note. These bires we have quite a tensy colores, compared to Economic individuals I've seen and the re-

to-mide unimported into, a sover little and a configuration of lixon, a sale Co. So the rest and the live. The live is a little in them Lept. 24,1340 I not were grown dignish, and there were find the control in I had to be rough of the row clib righten, nearby. The vertex begins to been, the they nervise as perfectly, circles and in the vertex of the many calles and the vertex of the vertex

Fruitridge ave, N. of airport, Sacto, Calif. One on low venetiment bank at 7:30 a. m. October 11,1949 3 mi N. of Rio Vista dolono of self. Inc I shoot may were administration. Biggs - Britta City Road, E of Gutte Cook, Outle to. Nov. 20, 1949

Nov. 22, 1949 17 mi. Nof Jan di on, & Lis Biops 6, Af- 6 gom - 1 do g soul.

Dec. 6, 1949 3 miles due west of Sacranento Wildlife Rejuge, Elenn Co. - 2 on actch leve al noon Mar. 20, 1950

Jan. 24, 1952 Pair at next hole on Florin Rd severt of Exclaim Rd. Sacto Co. Lough Mar. 14, 1953

Silver group fut, USSO + at Duyting you to Nivada. Fory Lun. July 4, 1959 Wif Electer Gulsta, Fute 6, City. june 2, 1960

Juny ? 1960 - Country Guis Centre - 8 am Sact to Cais!

Northern Spotted Owl

Jan. 27, 1947, near Rockie 4-corners, 5 p.m. attracted to aclump of fir by a "hoot-hoot (joined)". I found it perched high in one of the firs. It was barred and spotted below, and the undertail was light and barred some. .

٠.

Strix varia

Barred Owl

Feb. 16, 1953, on road into Singer Tract, SW of Tallulah, Madison Parish, Louisiana.

perched on hardwood at 10' by road. Observed at 40' distance. Was flushed,
but back at perch 3 hours later.

5/19/19 Sewer Cricto Conal - Mid. aside alit in open on 40 de consequent and of 19/19/19 Sewer Cricto Conal - Mid. aside alit in open on 40 de consequences.

9/9/19 Morris. many allegous young w/ short-tail.



Observer fred Evenden, Jr.

Great Gray Oal

White roll lies trips on which observed:

cotientex neurloss ne uloss

Location and observations

Fall - 1936 Hunting trip with along Pavelets NE of West Woodburn Ord



Observer Fred Evenden, Jr.

Long-eared UNL

number of riese trips of made our ryee: 512-522-584-891-895-1259-1850-2131-2326-2558-

Asio Wilsonianus

Late Location and operry along

At Paygants at Woodburn, Congenciaving winter 1941-42.

Oak-fir belt, 3 miles west of 12-mile Swamp? ???? sp???? flushed one from Feb. 25, 1947

a fir woods on the south slope of the riage. This identification is made only on elimination, for we didn't see enough to be sure. We are judging

it only by size, as the only Possible owl that it could have been.

Mar. 11, 1947 Oak-fir belt, 3 miles west of 12-mile Swamp: we decided to track down the owl flushed 2 weeks ago. We located an old oak snag about 3 feet in dia. and found an open "niche" or recess about 15' up that had fresh pieces

of grass stems, fern, etc., worked into a semblance of a nest. No bird,

or eggs were found. They should lay later, if this is the nest.

Apr. 8, 1947 Oak fir belt, again: flushed from a tail fir, within 30 or 4. feet of the snag we "/ck picked" for the nesting site for this bird. Again we caught only a glimpse of it, but again its smaller size put it in this species. The nesting sites checked again showed no use, so perhals the birds have-

n't started to nest yet., if there are two of them.

18-mile Evamp. 115. Then, and at integular intervals, therealter, a thistled "pueces e-1970. TTT" c ame from the dead as his errors the case. June 23, 1:48 on the test side. I chala whistle a good initiation with, the it we had answer inclodiately. I finally sot it silohuettse a aim t the say and saw it was take [90]... It is torvals it would be not be so here; eached in the work, and when it has be a female or important hor co-

la-mile S. camp: Thus have from a sh-ock area in the sales on a colored or July 8, 1948

swamp at noonting.

moreomer a on the art at a.

Dec. 26, 1953 O'Neill's, 5. " waterland, alif. I abult appear in a phonomer where

one roosted last your.

C'heill's, Courtland, CliC. I adult with 2 smaller birds (last mern's Jan. 11, 1954

young? in camber tree. C'Meill's, Courland, Jalif. Now are 4 long-cars in comphor tree. Jan.13,1.54 Cedar Breaks Nat. Mon. IronCo. Utah. One after chiomunk in spruce July 2, 1959 forest at dusk, by our camp lantern.light.



Observer Freq Evencen, Jr.

itaber of affect or in

Short-eared Owl

137-205-210-272-293-198-301-308-309-310-317-331-375-352-425-592-785-803-823-835-869-885-960-1396-1398-1492-1492-1498-1615-2276-2392-2525-2581-

Asio filmmeus in me tue

ocution and popular clima Late corveil is Of- . tom er amul april 10, 194. on the substitution of the control o o to a installed a lieuwood to a com-Lewisburg or an one individual ditting of fonce good new all Feb. 25. 1:45 irol the nighway. increaces in it -miles areas 0.0 March 6, 1046 the line is not in the instance in our missay to an ap. Isopping it a at. Mtn. Home, Idaho, Area II, south of town. Located a nest in an alfalfa May 14 1944 field, after flushing an adult bird. It is on the "hump" between small irrigation charmels. The "hump" is 15" wide. It is a saucer hollowed out in dirt, about 1" deep and 6" in diameter. Bottom is covered with ary short pieces of grass, stems, etc. Mtn. Home, Area II, We investigated the short-eared owl nest that we found May 16, 1944 on the 14th. Flushed one bird off the nest. The nest contained one dirty-white egg, bluntly oval, ends are blunt, but partially pointed. Egg is l_a^{\perp} to l_a^{\perp} " by l" in size. Mtn. Home, Area II, The nest found on the 14th had one egg in it on the May 17, 1944 16th and two eggs in it today. The advit aion't leave the nest till I was almost on top of it! Mtn. Home, Idaho - nest still had only two eggs. Owl stayed on till I was May 18 1944 within two feet of the nest. Mtn. Home, Idaho, Area II (ob by Joan). Owls nest now has three eggs. May 19, 1944 Owl stayed close by. May 14 May 16 May 17 May 18 May 19
Nest complete/l egg 2 eggs 2 eggs 3 eggs It looks as if the owl follows the habits of the Marsh Hawk and lays an egg every other day until the clutch is completed, with the first egg being laid on May 15, in this case. Mtn. Home, Area II, Tried and tried to locate the owls nest, but since May 28, 1944 I left the field had been irrigated and the alfalfa and weeds we e over 3 feet high. I did locate the nest site, finally, but there were no eggs, either they were lost in the flood of irrigation water, or else the owl moved the eggs to a new site, which I haven't yet located. Mt. Angel one flying across a field, landing, and then taking off April 12, 1947 to chase a Brewer Blackbird Fern Ridge Reservoir: one observed on open mud flats at south end, not Feb. 8, 1948 far from a raft of geese, and a huge flock of sandpipers. Peoria, Linn Co. O one flying over stubble riela. Secuca to reap Mar. 14, 1948 either & or 4 times, and then glice a very short distance. Meadowiew, Lane Co, Oregon. - 1 seen.

april 8, 1949

Dec. 13, 1949

Dec. 13,1949

Pernley Sink, Lyon Co. Levada - a total of 5 seem perched on low bushes about 5 p.m., or flying over marshy areas along the hiway. A winter or migration influx? Some appeared rather small. Since there are few records for this owl in Mevada, special note should be made of these obs (also Toulon) Lower Loverock Valley, rershing Co. Levada - 445 p.m. a total of 5 seen flying over pasture and alfalfa fields in the area. One made passes at something on the ground, and when it missed a male marsh ahwk, following behind made a pass, too.

Sof Frankling secto 6, Caly. Flushed 3 from I weekly field in a me Jan. 28, 1952

1.12 27, 1958 5 of A Talus Com Love on El Cotto Dd, Saite G. Colf. 8 of of in rice of the laid Roway, over the many officerent builts on 1.04 of 3, mm.

Saw-whet Owl

(Ee/- 26/1)
Number of field trip on which seen:
336-338-

Cryptoglaux acadica acadica

Oct 13 1944

Mtn. Home Reservoir, Idaho. In one of the quince trees in the deserted ranch yard of the canyon. The quince still had their foliage, while the other trees were losing theirs. On Sept 19 the owl was in the Honey Locust, as tose leaves were still quite dense. At any rate, the little fellow didn't so much as both to move, when we found him. He just locked at us, with those big eyes with the bright yellow iris. We stood within two feet of him, and Joan could have caught it very easily, if she had so desired. He showed no fear of us. (He?) was perched on a horizontal limb of the quince, clutching the hind half of a rodent that he had been feeding on. It was completely eaten to within one inch of the hind legs. The tail was about 22 inches long, light below and ark above. Body fur was grajish, with a reddish tinge on the back (Microtus??) But back to the owl. We spent fully a half hour watching him, and he had no fear, just curiosity. Occasionally he would slike that nicitating membrane across his eye, or blink them shut, revealing the heavily grayish furred eyelids. It was 7-8" long and the tail seemed quite short, altho the large head seemed slightly out of proportion. The wings, head, and body were a soft chocolate brown, ilecked with white and gray. A few larger light patches on the wings. The underparts were grayish-white, with broad, reddish-brown strives on the chest and belly. The feet were fuzzed right down to the black toenails. (fuzzed white and yellowish-brown). The bill was black. A prominent white "V" is over and between the eyes, sor of like eyebrows; quite a bit of white on the forehead. The face was brownish-gray.



No. of field trips on which seen:

Huliotta Poor-Will 2617-

Phalaenoptilus nuttalli

Snake R. Canyon near AAB, Mtn. Home, Idaho. noon.
Several of these sitting on horizontal limbs of trees around a deserted farmhouse near the river. Had white corners to tail instead of crossbarring.

Oct. 15, 1948 Nigger Hill, El Dorado Co, Folsom Dam Site: found a nearly fresh Dusky Poor-will at the edge of road, evidently hit by an auto during the twilight of morning or the previous evening.

June 21-24, 1949 Elko County, Nevada - com on. Observed some perched on fence posts. April 17, 1950, Red Bluff-Corning, Tehama Co. Calif. 1 seen at 6 p.m.

teber 24, 1954, Just word into 33 miles it intremux utation, it. 1. The moved area a telephological end of the liver. If the first of the hard pas endows, and use instructor in the thirty.

the true = Since In to be post to lide of for 5 days I'd better write you a card I be the week gets for along. The big and but in he was the big south-Frake (Intensity 2.0) that centered along the north side of the diseit. Rocks rolled like marble Ith were the weeks and the state of the in a grant of the state of the state of the Commission of the the order of a star included the top the same was the mount of the second the contraction the the outle of the way is I had been with a will serious. The was the the wine - 5%. The process of mine attended Yours there know a come in the con the the Miller of the server the year while to be ; in. Hours of the down? were with Jank

Texas Tilling Nighthawk

Chordeiles acutipennis texensis

March 28, 1950, Balboa Park, San Diego - 630 p.m. twilight - one flew about the car while we were parked on the ridge top east of the zoo entrance.

Cuban Mighthawk
Ch. vicinus
Vay 20, 1965, Plantation Mey, mor Sirunts, Meys, Floridi. over house - has ?-part call.

. Siopecrwill

May 11, 1.67, Con Canal at Prickward Road. I on limb and bot ered by scolding titmice and collow withher.

Observer Free Evenden, Jr. . uro : or right trage on which our ivea: racific ighthams

2-5-10-17-21-79-90-158-159-161-166-168-178-238-240-242-720-321-321
923-924-928-929-732-1228-1238-1238-1253-1267-1302-1308-7209-1673-1959-2119-2130-226-2368-2464-2470
Chorceiles minor has an in 7-22-1253-1267-1302-1308-7209-1673-1959-2119-2130-226-2368-2464-2470-Pacific lighthamn Chorceiles minor becamin 7632-2637-Date Location are or envasions at home, clossen Ordin a covers of a color of the other a line in the iune 6, 1940 Corvellie: 01 Il an the s'is to his transity of the sing of sales of a half and 17 1/30 p.m. flying Victory is some June 1, 1.41 June 20, 1942 Silverreck. Or Second. n. 1. 21, 1942 West Woodburn: Foggy morning at 7:30. Flushed it from the grass at the Sept. 2, 1942 base of a clump of willow. It had light plummage, and it surprised me with the silentness with which it flew. Kennethallalker reported nearing a Nighthawk just south of Corvellia along the relief therey He heard it several times the convinced. Corvallis. Feb 13, 1943 me Chard 7 and, which. 13258 WIT 7-5-43 Parifice 11 exhibitions aling frequently, and windly which a the ground (times 5), 4° K I the ground devil) ne Chard Field Wash. 7-15-43 Calling at 0100 P.W.T. June 10 1944 Mtn. Home, Idaho. During the past few evenings I have heard several of them overhead. I heard one at 1230 a.m. during a heavy rain storm at the base, too. June 15 1944 Mtn. Home, Area II 9 a.m. There were at least 8 or 9 of them in one small area in Mnt. Home this a.m. Also saw one of these that seemed to be short and quite fat. It, too, had white throat band, and white wing bars. It gave no call notes. Mtn. Home AAB, Idaho. A pair have a nest inthe rocks and gravel just off June 17, 1944 the EW runway, along the south side. 1245 noon, Woodburn, Ore. First Nighthawk of the year. My folks and I June 7, 1946 were in the back yard when I heard this old familiar call, and I loked up and saw itw "floppy" flight as it headed northward. Breitenbush Eiver Forest c amp - Saw and heard one between rain showers. June 13, 1946 Heavy rain all p.m. Corvallis, and Salem, Ore. They are common now, flying in broad daylight June 19, 1946 Ump Pioneer to Mario Lake trail, Minh Col Saw one come in low over August 7. 1946 the riage about li a.m. Corvallis, $80^{
m o}$ F. a fright of 9 of them showly feeding is they writed Sept 19, 1946 southward. To call n tes were miven. Tom accallister heard one this a.m., and this evening they were alying June 7, 1947 around the lig building on the Compus after as k.o. MacKenvie F, RE of (pringfield a pair in curting fight over gravel bar June 26, 1947 at big bena in river at 7 a.m. Corvallis - called by 7 m.m. Aug. 20, 1047 " 645 pion. nug. 21, 1947 " " " " o ..h. (There wie 10-15 togeth r. 12-16 Stamp: estimate 11-15 recent; come a little 1 sger the others. Aug. 23, 1947 June 23, 1948 Seldom calling, and most feeding over open water areas, occasionally cipping low to scoop up water with their lower bill. Graceful and beautiful faiers. Muc Lake, I of the better the case on classical ever the south end July 28, 1948 of the lake.

Woodburn, home: none heard since quite cool weather came Aug. 14, p.m.

Aug. 18, 1948

the state of the s

.

.

June 4, 1949 Portland, Oregon, 815 PST - several calling and Tlying near amanuel los. None had been seen or heard yet in Calif. so their ar rival here first seems peculiar.

June o, 1949 Corvallis, Oregon - several seen and heard.

July 30, 1949 - 11 p.m. Klamath Falls, Oregon - Calling over town.

Sept. 5, 1950 - Corvales O. ogn

Aug. 4-5, 1951 Silver Láke, Amador Co. Calif. 7,200' elev. 1 was seen coursing over the lake the afternoon of the 4th. By 9 a.m., the 5th, between 18 and 24 of them were feeding over this same arm of the lake where one had been the day before.

May 14, 1952 - Les Vegas Wevada - several at E and of town along US 466 at 9 pm. 1.0. 5.7.

July 24-30, 1952, Patterson G.S. Modoc N.F. Hodoc Co. Calif. 7,2001 el. every evening, some mornings, early, and in afternoon when thunderstorms browed.

aug 21, 1961 - Logor bray, Sincelin G. Ovegon - in evening over bry week.

any 27, 1961 - Ewelle, Maniell F. G. G. - at 745 pm (1851).

2/1-4/96 La Cross Wis Diff say than sell - They came to picking, w/ gettered chatter notes.

,

.

.

· Dut "

·

Observer Fred Evenden, Jr.

Late Location and observations

Topoburn, Playment's works: ov : . this aljing in sand it come over-.t. 17, 1941 head. One set his is on then by the outside of a harp-shanned by a.

Corvallis, Lower Campus, 4 p.m. Saw 3 of them flying low over the lawn. May 16, 1946

Salem, Oregon, "illamette U. campus, 650 p.r. 10's and 100's of them Sept 9 '46. circling in large groups over the campus this evening. The air was filled with their "squeeks".

avery Park, 04-5- p.m. a flock of from & to 40 were circuing rapidly Lept 18, 1946 through the wir at treaty level, near the ease of the forest. I watched some actually land, momentuming in the top of a 1. Fir to reed on insects there, in company with 5 Tobins, 7 teast which s, and a flictor - there are 4 V to write there all at one.

Set 19, 1846, Poison Cak Lane, inn Co. 15-10 of them circling and feeding near the eage of the weag at three-ton Level

May 5, 1947 Corvallis Par. Gordon told me about some he'd seen at a residence where they have congregated in years past, and so I checked their arrival to agree with those I heard on Saturday at "oodburn. I saw 7 altogether.

Marion, Miller's Butte - 5 chasing a Sharp-shinned Hawk which was May 17, 1947 carrying one Vaux Swift

Aug. 25, 1947 Corvailie of 15-10 at these consider, are degine mich at by

Corvalli: Gravel lite - 15 - . . yind no to man over the river, and mult-Sept. 7, 1947 by time, charles, and receips in the rain.

Coast Fork, Willamette R. Bridge, on Willamette Pass Hiway, E of gooken, 1130 P.S.T. Sept. 2, 1949

July 24, 1952 Burney Falls State Park, Shasta Co. Calif. very common in amphitheatre at the falls.

Sept. 6, 1952 12-mile Swamp, Bruce Sta., Benton Co. Oregon. 35 over swamp, calling, and occasionally with songs, too.

Blackbird Forest Camp, S. Pr., amer. R. Aldorado Co. Calif. about 50 circling Aug. 23, 1953 over river canyon area at 430 p.m.

Oct. 16, 1955 To Farallan Islands - I came over boat I di. d of JE island group.

At Sloughhouse, Sacto. Co. Calif - 3 or 4 with swallows at 9:30 a.m. April 9, 1958 49299 New L. Brillin, + Mat work, South G, Coly - 12's accom-

ang 15, 1961 June 9-10, 1962 China Flat Opgrd, silver Work, Ildorado Co. 2 heard and seen above tree tops and slower at of Silver Forhat (p.m. (roosting in tall yellow pine or fir? - feeding in toos on insects about buds??

July 20-22,1962 Chapman Crt, Cogd, Taboe NF Sierra Co. Calif. 5000 tol. about 15 or 20 overhead in area - suspect they reested in crevices or helps in tall Red Fir (A. magnifica) at our own cumb. In air for as or yet 6:30 a.m. to late asi: how. FDST. Rusy and noisy most of dar overbead, the gone for s ort inte vals.

.

•

Black Swift

- Aug. 3,1949. Yosemite Valley, Undera Co. Calif. just a short climpae of for them sweeping across face of cout-incing clim on rose to betch hetery. Their coloration and size one unsistable, even the file of seem them before.
- Sept.10,1961 Lassen Peak Summit, Shasta Co.Calif. 6 of them skimmed swiftly over top of peak at 1:30 p.m. heading southerly, and without deviating from course.

CHICKEY SIGFE

May 21,1961, Vinous t. The Club, (tt. wa Co. thic. tried 3 with V-set wings into wind and hold still for while in one position.

5/14/16 - Pennyhite Stock -Ch Sunft war concluy of digiting to Car in a



Observe Free Evenden, in.

Aeroneutes s metali acetalis

I ate | Location and object to in

Yay 21, 1949 Coast S. of Corner, Contemp Condender over read clifts - nest cavity in Serice of 250' sea chill now the lack have nest to two coulon's rand.

June 23, 1949 S. Ek. Humbolat Fiver, Lko Co. Nevaca - on 10-ma e treel, and in car, on near mouth of S. Ek. One pair observed to be joined, and templed about 200 feet towards the earth before outling what - presumbly the

May 13, 1950 Glacier Point, Yosemite Valley, 7200' elev. pair chasing each ther overhead at the point, and giving call notes - 1st time I've heard them.

March 31, 1953 Packeco Pass, C.S.H. 152, slift first Wof Sumit 1:30 p.m. 2 sean.

april 23, 1958 Montuelle Dan Site, Calif Gen. PST. #2/Ging about dom. in the campon.

Black-chinnel Hummingbird archilochus alexandri

March 18, 1947, Bill Williams R. Delta Canyon, Ariz: several seen. 1 old nest found at 4 feet in thorny bush. May have been a Costa nest. These seem to be in heavier vegetation and more moist areas with tall mesquite in canyon while the Costa is out in the drier open areas in plateaus and dry washes. A good problem to answer for no Costas were seen in the canyon. The Black-chinned feeds on many plants — chiefly the bud of the Creosote bush at this time, for they are flowering in profusion.

March 22, 1947 N base of Chuckawalla Mtns, Calif: 8 a.m. a pair were busy courting.

May 5-8, 1975, .onson's Tucson Arizona: nest at 71 in olive in closed courtyard.

2 tiny errs 5/5. h/19 was "irst setting the. 5/6 - h goung, and hegg at 6 b. .. 5/7 2 young. """. size. 5/1 rate of growth was rap d, dubling, then trebling one size in clube of days

·

Archilocus colubris

ubv-threated Hummingbird


5/19/69 above great force, mit - ad on meet & 26 own canel on horing experiench. 3/23/76 F 16 is wet on day (1.6 1.7) of consent of



1108-1878-1897-1902-1903-1904-1912-1959-1978-2062-2187-2209-2254-2272-2274-2321-2322-2324-2325-2368-2411-2422-24 45-2446-2470-1-2-3-2569-2594-269-2594-269-2594-2272-2274-2321-2322-2324-2325-2368-2411-2422-24 45-2446-

Calypte anna

March 19, 1949, Inverness on Tonales Bay to Pt. Reyes, Maris Co. - 1st I'v esen.
Oct. 1, 1953 Putah Cr. dangen, (.5 ml. / ml. inters, 1 to Co. M. Mr. h. molling r
impatures on Corb hanthup, blue ock, and fooding on scarlet Bilia.



Number of field trip on which seen: 344-536-538-539-540-544-545-546-5\$1-552-1108-1207-1837-

Nov. 7 1944

Mockingbird Canyon, Arlinton California, 9 a.m. They were very common. The body was quite greenish, more somber on the female, and the female has a small light collar below the darker color of the head. Purple flared throat patch of the male, changed shades in different lights. Also there is a purple patch on the top of the head. The tail is blackish. They are quite tame, and I approached within a Yew feet of the perched birds. They give a very irritating squeaking call Maw. 19, 1947 - Havasu L. 2 mi & of togeth: willow from mattress being used in construction of the next

at the water gange stacon.

Feb. 17, 1951 Vidal Jct.-Earp, S. Bernardino Co. Calif. 2 seen enroute, and Gale Monson states they would have to be this species, tho he has not seen them yet this spring - First 1951 records.



Blue-throated Hummihabird

Lamoornis clemenciae

2541-2542-2546-2604-2607-2611-2613-2615-2620-2621-

Proad-billed Hummingbird Cynanthus tatirostris

1549-2555-2554-2558-2560-2604-2611-

Broud-tailed Hummingbird

Selasphorus platycercus

of did look andray about the New Miden week 10 miles 25 got and fly

2604-2613-

A told Develophing

Durened i Trons

•

Observed Tred ventile . . ours, who exists one, exist in the first content ly 3, 1040 rch 7, 1941 torvallis, arous home: " at two of there for the first arrivals as the est one land of a filter division of a land of the land of the land that cover the land of the first of the land of the land that cover the land of th June 10, 1941 the call L. It is available of it is employed for / hi: iv . net, or 5- aloge a t: " in spins how of them sixthy as a co-rection of the ent urch 15, 194% .ug. 9, 1841 Corvalles tryon tare ' lego ampere -Feb. 9 1943 L. W. Mary . Teverson where a lapture Hermanyon od . E. T. H igo saw it too. The ay as cory winey. Woodburn - Saw, my first one of the year, at though L. r. Stevenson Mar. 12, 1943 reported them on Feb. 9 (a14) - Horsestone Lake, St Paul, Oregon - 11am. 3 - 23-43 Timber doing "lovedies" He limber to wout 50 , well arest dove straight down to the small swithed on a strate at the bottom of the dive, the sound of the wings checking the fall in as great you such a witter bird the gove commend notes not heard at any other time. - Canas, Wash. 4-11-43 or doing several love diver. - 1 re male did 4 consectative love dives 6-28-43 Palmer's Hill Camas, Washington 5-7 p.m. May 20 1944 One pair have nest in an Indian Peach bush about 4' above the ground. It is within 2 feet of a woodland roadway up a hill behind Palmer's. Nest is 2" across and 2" deep. The cavity is 14 inches in diameter and about 1" deep. It is cotton linea. (Or thistle down). The outside is covered by moss and lichens. It contained 2 white eggs on May 9, and they hatched on May 17, according to Peggy. Corvallis Gravel Pits of Saw my first one for the year. March 19 1946 found nest on limb of Ash tree at 15-20 feet. In shade on NW side. May 25, 1946 at Peoria Ferry, Benton Co. hea Butte, lake and Butte of still quite common around here.

Rouge River, Illahe, Curry Co: 10 a.m. several pair doing love dives.

August 9, 1946

April 26, 1947



Rulous Humingining

i age

May 1, 1947 - Bot Morris reports at yaping out of it is in see in a String at befeet on the N side of the lives it was in a thereing of word zona. The next was on the "outside - lagsaids" of a dooping 3" Granch. The rest you is make, I considered him many

Macionala Forest of watches one brillian male recains on St. John's lort July 12, 1947 and Camada Thistie. Saw two pair at the same time in this vicinity.

Liadle Santian Liver of nest in red huckleberry bush rowing on perpendicular July 13, 1947 N-slope chill at 20 feet above the river.

heard the first one of the year, and saw it later, just like last year, in my own back-yard (410 N. 16th, Corvallis). Hovering over one on Grape. March 26, 1948

Pucking P, odburn tea tea hovering near top of 60 woot count, and I 73 May 8, 1948 watched him record crop to or, till he reached the bottom, checking each hole made by the Fou-broakted Supsucker workings - 1 (Ling for say, ?????)

Smater Lake \mathcal{Q}^{\star} fin to sales that $t=\infty$, on the structure substitute for the tJuly 25, 1940

Inverness on Tomales Bay to Pt. Reyes, Marin Co. Fany seen. Several Mar. 19, 1949 feeding on south slopes on Arctostaphylos.

Gualala R, Mendocina-Sonoma Co477 a.m. feeding on salmonberry flowers. Mar. 20, 1949 Lithia eth, o't no, ton - o'vial reen - t fintingam civing. aug. 20, 1949

Hogan Res., Calaveras Co. Calif. sun-posing on ground in flower bed at Ballards lg. 22, 1952 Morro Bay, falif. in park among cypress and rocky point shrubbery. Love dives. Feb. 3, 1953 april 3, 1954 Win Land Park Jacramonto, Calif. Full M':

June 26,1954 Camp Harvey Mest, Scho Make, Aldorado Co. Calif. 1 M on 9000' ridge N of camp.

.

.

.

•

Obsetv r Free Lwinern, Jr.

Milen's Hu ling in

Rumber of field trips on Valon observed: 85-1543-1614-241-2667-

tel aphorus alicai

late | Location and Control

•

Observer brea Evenuen, Jr.

Calliope Hum. incbird

lunter of field tripe or which observed:
88-10-4_1259-1260-14197-1632-1693-1696-1925-2134-2320-2321-2366-

Stellula carliope

Late

Location is observations

Nov. 14, 1948 Monterey, Calif. 9 a.m. 2 present, perched on "head" of Aguava or century plant in part area along the edge of the bay. Chased each other.

Dec. 30, 1949 Fardee Res., Calaveres Co. Calif. 3 ... one small hummer of this or the Costa species feeding on flowering our ant just south of the lain dam, soaring up above the bush, and perching on a nearby buckeye. De-me! quite small. Good winner record.

.

Sept. 12, 1942 Woodburn: This was a surprise, indeed. A pair of Belted Kingfishers were flying around our neighborhood. One lit on a light wire in front of our house. We had the lawn sprinkler going and one of them came down and bathed in the spray. There was also a young flicker, a chickadee, and a young robin enjoying the spray at the same time. The Kingfishers gave their rattling call at irregular intervals.

Jan. 17, 1945 Corvaint: . - mint in a graduate and all literate and at the limb.

While a telling it flay from one tree to another, it stopped its form a complete and hevered for hoor 4 seconds while beating its wings, and it lost no elevation, but aid but up slightly pefore going forward again.

Oct. :0, 1946, Forvilli: () ... increase the rearry. Is to medice in the algorithm of the such fence at the parce (each hacth), on , purchase there a minuses in the such settles.

April 12, '47 Mt. Angel: saw one perchec on wire line over road, close to flooded fields. River 1 mile away.

June 15, 1.47 Seal Rocks, and coin Co- pair have nest home on sand bluir whose base is at high-time mark. It is in the sand at 14 reet above base or the chiff

July 2, 1847 — Hamile tham, "- h pair of there execute the test show on the case, and we which continue the chartening pover too them, and a proches for the high in a small ach.

August 2, 1947

Avery Park - 3 s.m.: one make observed rhying over this point stretch of the river, with its underpoints provided by trailing in the water, and the wing tops mitting the dater everyonce and a file. Every 15-fer is that are pred into the water, and then required laying, only to studing and again, to 7 or 8 times at its T tatched. Its a thoracal tenot to cutch fiel, but note to bathe, I assume.

Aug. 14, 18-7 Inllamette I, corvallis of Secon. one sound coich high, hover, and then scroop up, therm a capa, climbing up again, nover, scrop upstream, clive dorm-ward and public out without hitting the water, and skinning above it, as it crossed the relate .

Feb. 1, 1949 Long Tom R, 3 miles N of Monroe, Benton C8 (THMCA). 2 seen - both with small spiny-reyea fish in their sills.

Biawell Park, Chico Calif. 1 p.m. - observed 2 perched on limb over north edge of creek, about 12 up, with the mule bird atop the female, holding on with his bill in her crown feathers. Later he fell off and flew away, and then she left, and she was the first to return, followed soon by the male, but both flew away again before any further activity could be observed. Previous to my first sighting these birds I neard a great deal of chattering for a period of several minutes, which resulted in my finding them.

Nov. 14,1950 - 30-40 mps. - 305 Proc. One of several minutes.



May 19, 1951, Merced R. Yosemite N.P. Calif. one flew at 35-35mph, upstream, paralleling us for about $1\frac{1}{2}$ miles.

Nov. 7, 1951, Little Grass Valley, S.Fk, Feather A. near LaPorte, Flumas Co. Calif. just after noon we found 4 hanging around a dead lodgepole pike at the edge of the river. Very noisy, chasing each other, and giving a variety of calls and scolding notes not be and before.

Morch 23, 1952 Mill Cr., N. of Ft. Bragg, Mondocino Co, Calif.

October 24, 1954 U.S. 40 near Cosco grove Calif. I along stream.

Cot. 23, 1055 provider ., niche - ord will to bear to the

Die. 27, 1957 Cow Creek, 4599, Not Wolf Corch, Josephine Go Oregon. Heavy Rain I seem in the seem of 11, 1958. Ann. R., So irden Park, date G. Colif. One adult Mon next at 7-9an.

W/ F along rive. Vast wiland was at end of 5½' leap horizontal burrow, with 2

grows in bottom where feet worked. Birds would have to story to go then it.

3 you, black skinned, of "leaded" w/ coard feether. Displayed no curiority to sunlight

June 23, 1958 - anex. R. sorts a colif - 3 pm 3 forms now almost fully feathered like abilt, w/ some chestomet mixed up three chest band fadjacent white. I triad to second "crapt of cringed" out up long slick at which point they seemed all militrity of "crapt of cringed" into bottomed burrow, in dist. One see wind head a lit of ever when chross rim out appeared immobilized - braining or least? abuilts stayed 200' away across rim while of woo of brestow.

June 11, 1958, American to Garamerto Calif. I pain, Fout of burrow at 7 a.m. Caucht

M in net as it flew from the burrow about 8 a.m. Intrance 121 above water.

Apr. 26, 1959 - NW of Bridgehouse, Sacto. Co. Calif. Watched pair in courting flight

and chase.

· ·

Colaptes cafer cafer lete Nov. 30, 1941 a large small to the interpretation of the con-Avery Park, Corva lis: adults and joing of the year common. Their 25 Aug 146 favorite perch was a short lime high on a tall fir, in the cun, where they were usually seen preening. avery Park / Corvalians. The recoing on the ground, schauching about tept 17, 1940 big-leaf Mayae Leavel. Sept 18, 1946 Aver, suck," one gave a car. similar to tail alarm notes. onitor, Ilt. nocl, the obsume a cant, character to the life of the observed in the life of the observed in the life of the life. Nov. 28, 1946 Tov. 50, 1840 -Most were in the grown, weil intermed. Poison Oak Lake, Linn Co ? - many seen, and most were feeding on the ground, Jan. 16, 1947 flying into tree when flushed, and as usual, go into the dipping act. Jan. 21, 1947, Beaver Creek School Common on the ground in all the fields. Saw upwards of 25 in one grassy field that may have been an old pasture. Avery Park, Corvallis, 7 p.m. saw one fly from a maple tree to the airt Apr. 21, 1947 pathway, pecked around in a small area for a few minutes, and then proceeded to dust himself in that spot. That's digging your own bathtub ! Sept. 12, 1947 Big Take has and 945 h. . . : On tallied 14 bines require on the more is approximate the following the figure t , t and t , Feb. 23. 1948 N 15th & Jackson St, Corvallis 745 a.m. Ø/ cool, calm. 3 of them, sex, not determined. One on crossarm of telephone pole, the other two on wire lines, 5' to 5' away, and below the first bird, who seemed indifferent to the aisplay activities of the 2 other biras. These two would utter excited callnotes, and at intervals fan the tail graciously, or nervously, and tilt the head upwards over $\frac{1}{2}$ -way, or to the side, still giving voice at same time. Twice the two biras got side by side, almost facing each other, wing the same thing. Another time facing the same direction. Dipping and bowing also took place all this time. Cotaige Grove Practical Paratrat 25 feet in broken, rott , but iving June 23, 1548 fir, in party can a core. Memiur to NI. Fire ent in section, Tesoing young : ile I vatchec. Corvallis-12-mile Swamp (IBM) Picked up one dead along the road. It is a Jan. 25. 1949 hybrio yellow-shafted x red-shafted. Has red cresent on back of neck. Fed and black mustaches, underside of wing and tail feathers mostly yellow. Sex? Had been feeding on poison oak berries. a: Corvallis, Oragon - 5-11-21 - Tayyo, 10' up in dead offgets tree - Kole 2' it 5-13-21- 8", 4"... " -"18" deep. 5-14-21-7", 20" " deed ald stib - 20" " hole. 11 11 11 5.24-21 - 6", 8" " naturas caring of " 5-26-21 - 7", 6" " dead cake state Jana Maryon, Maryon Maryon June 3, 1941 Johnson, Porten dy Corgon - Bying in a thore 3 below the write Se recht al sent in dead about



10.1772-1/15-1774-1775-1774-178-1774-178-1781-1781-1782-1724-1722-1724-1727-1729-1730-1731-1732-1744-1745-1749-1755-17624-1762-1762-1764-1765-1766-1767-1766 univer of field tripe in Chich operived:
26 - 84-85-165.168-169-172-173-174-175-263-264-265-266-269-271-272-273nea-shafted Flicker 276-277-278-279-280-328-332-333-336-337-338-339-340-540-543-Twin Falls - Hagerman - Bliss Adob Observed a pair of them on a telephore May 3, 1944 pole cross-bar. They bobbed their necks up and down and from side to side all the time uttering queer notes. Black Butte Res. area, Stony Creek, Tehama-Glenn Co. about 1/lor 2 acres. Nov. 16, 1949 Stone Lake, Lambert, Sacto. Co. Calif. 1 flicker nest on NW side of dead May 8, 1951 snag at 20', with tree swallow nest on SW side at 17' height. May 9, 1951 Armstrong Ranch, near Riego, Calif. nest on S side of cottonwood at 18'. July 27, 7,52 Nother (4.600) Flight 2035-2219-2220-2224-2205-2226-2226-2227-2232-2234-2237-2239-2242-2243-2242-2318-2319-Oct. 6, 1952 May 10, 1954 Southside Park, Sacto. Calif. 3 pair seen, one pair courting-dance on limb. Asilomar, Pac. Grove, Monterey Co. Calif. nest in Monterey Pine on west side at 23' ht. Dia. 12" there. Sacramento By-pass, Yolo C. Calif. 6:45-8 a.m. 40-50 seen, oslty igust 22,1954 loosely-paired. c. 5, 195h, Relson Tos. area, Todor Ch. Culif. The Rills; at Carrie Court Court Tonbor Hawk Dec. 26, 1951. Bryte, Tota to. chif. I dissing narrow Tawk. Oct. 31, 1957 Camichael, Calif. From AS.T. Watered one personing for award orienter protection with wind

May 31, 1959 Pope Beach marsh, Lake Tahoe, Eldorado Co. Calif. nest at 42 on


May 31, 1959 Crater Jake, 80 0' el, V slope upper Hope Valley, Albine Co. Calif. West hole at 1/4' on W side of 30" diam. red fir snas.

SW side of 9" diameter lodgepole, and only 15' from lake.



Mearn's Gilaed Flicker

Colaptes chr, soices mearnsi



S.P.W-2226-2228-

Observer Fred Evenden, Jr.

L ate

Location and observations

pt. 22, 1940

caca's priore on cutte brook? round no make for maile white he also about a lest whole with. They silently, and not very high, he was bout a resolution the arrance it the highest point blacks in the up.

ovember 5, 1942

Port Townsena, Tas h. This afternoon and the morning of the 6th I saw a large male of this species feeding on the berries or rea fruit of the Macrone (Arbutus menziesii). He would cling to the small branches like a Chickadee and feed on the fruit.

pril 6, 1946

Wren, and 2 miles W on Newport Hiway. Heard one give his ringing call from some timber nearby.

May 18, 1946

Maclonald Forest 7 a.m. Saw two and heard others.

July 29, 1946

Little Abique of p.r. I we two of them that have a nest in to of a 60 foot snag. The hore opening to the Sw. One would leave the nest when the other returned with food, so there must have been young in it.

Jan. 21, 1947, Near Independence School, Benton Co., first saw it flying high, in the open, 200% yards from any woods. It looped around, and landed on a slim fir in the woods.

April 12, 1947 Ridge Road between Abiqua Camp and Milk Ranch: 4 p.m. Male calling loudly and fless to a tall fir snag and the "leaped" across 15 feet north to another tall barkless snag in which a nest cavity is. He lit below the hole and "hikea" up to it, and went in without hesitation. Whether or not it is a new hole, I couldn't tell, for it was on the SW side at 35 feet. The tree diameter at the hole level is 15", and the hole is elliptical in shape: () being $3\frac{1}{2}$ wide and $5-5\frac{1}{2}$ high. The snag is at the edga of a gully on a rather steep south sloke, which has been burned and logged off - only scattered trees and snags remains.

The male stayed in the nest about 3½ minutes, and hesitated on coming out - finally putting out his head and holding his bill open. In about 1 minute, out he came and flew to a barked snag 20 feet NW of the nest snag. Finally, after giving more call notes, it flew north over the brow of the hill.

May 31, 1947 -Same as pril 15, 1047 - of I came to the nesting snow, the female entered the nest, and stayed in when she saw I was parked nearby. The remained in for over 15 minutes, at which time I became tired of vaiting, and moved.

Aug. 11, 1947

Machonalo Porest, a pair were attracted by imitating their call, and by pounding on a "sounding boar d" I. Fir. They came within 15 Feet or me, and here very curious. They findly flew from tree to bree together.

March 26, 1948 Oak Fiage, 2 mi. I of Corvallis: Workings come on in the alder-ash swamp

area, particularly in the very rotten snags of alder. Tork observe a few reet and 35 feet also ve ground. Fnoto taken. Tinter workings, for they were reasonably fresh.

Mar. 2, 1950

Indean Valley, N. Fk. Cache Cr. Lake Co. Calif. 1130 a.m. one on floor of valley in white oak-digger pine open woodland. This bird seemed to be way out of habitat. All hills covered by chaparral.

Feb. 16, 1953

SW of Tallulah in Singer Tract, Madison Parish, Ja. Southern pileateds heard and saw many of them, much to our dismay, for they made us think of Ivory-bills each time. U.S.50, E of Pollock Pines 12 miles. Eldorado Co. Calif. Flow across good. to S. 5'8 P.S.T.

Man 30, 1959

, Richardson Grove, Humboldt Co. Calif. Heard one almost simultaneously with a song by a wren-tit nearby.

Aug.23,1961

Sequera 14 le. 76 get & cot of a feeling degrood besties liter to faceres.

6/4/74 Senseal, Mid - Cailling low on waln sooled energy, 10/5 m.

Feb. 2Φ, 1951 Carrizo Springs, Dimmit Co. Texas - First seen by Milared as they played around on a telephone pole beside the hiway. Colorful.

Feb. 26, 1953 Breckenridge Park, San Antonio Texas. Jatched them doing courting dances.



549-544-546-547-548-5-3-1838-1839-1840-1842-1843-2546-2600-2601-2602-2609-2615-2618-2221-2622-2629-2625-

Gila Woodpecker

Centurus uropygialis uropygialis

March 18, 1947 Bill Williams R. Canyon, Ariz: quite common in the canyon on the ground as well as in willow and Populus. Very noisy with a varied assortment of notes. Quite tame.

Centurus carolinus

red-bellied innehecker

| | | | • |
|--|--|---|---|
| | | | |
| | | | |
| | | • | |
| | | | |
| | | | |
| | | | |

I'clanerpes erythrocephlinus

Red-headed Woodpecker

6 A/66. Jake W. Mitton area, Tarder Dr. Stoff it Prinns, ark. I seen. The male came from real body in stub (1) of toler one cole in 12k-like view cords at 1 laws area now reminences.

| | C. C. Marie, 194 |
|---|------------------|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | • |
| | |
| | |
| | |
| | |
| | |
| | |
| • | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

2444-44-2617-2613-2617-

Observer Fred Eventon, 32. number of field tri, s on thatch observed: 16-83-85- 233-531-565-574-887-9012-439-951-954-955-957-958-962-979-June 11, 1948 Form riese: Of june attend I have character into a concentration to a confi the old Their in a comme satuation to I spoud a conjugace r ilytha come who are not read if the contract on a file and ! went in but we, , but a postume of Michigan particle . The contract construction of the contract construction of the contract c bachen up - . Who employed to the track to be I have a employed the velocity! mich't note in it has a flor i. Very pretty, at thet. The Charlier tells on the one of single the relief of in the old of the end of Fernal in the property of the state of the first tell of the second of the old of the old of the second of the second of the old of the old of the second of the second of the old of the old of the second of the old of th July 23, 1868 Firm described cults output block of tries to be a local series as been in a larger mobile tries, to on velocity of the color of terms. Valley Springs, Calaveras Co: Watched 2 of them 100' apart, catching Nov. 6, 1948 insects on the wing about 3 p.m. Salinas Res. San Luis Obispo Co of one in nest hole in snag on east slope Jan 4. 1949 Hole at 15' on the W side, or toward the slope. Quite open area of woodland. Folsom Dam Site, Sacto. Co. Mpaired, and hanging about the tall digger Mar. 27, 1949 pines, both living and dead. Several observed on trippione poles at West Point, Clev. 2000 feet - Calcurras Co. 1233 moon april 2, 1949 Quinda, Yolo Co. Wij. april 19, 1949 Natomas, Sacto Co: hest in 5-dead, topped Eucalyptus at 35' on east side. May 1, 1949 Open area to the east. This species sensitive to Tangy's horn. E of Oroville, Butte Co 7 a.m. 5 acusts on top of one telephone pole. May 6, 1949 July 11, 1949 Chico, Carif. heaviest concentration I have ever seen. Westing at heights of 15-45 feet in lubras lobate in hidre. Park. he dime oung at 7 and 8 p.m. Medford, Jackson County, Oregon - In black oakse E of town. aug. 21, 1949 Armstrong Redwood State Park, Sonoma Co. N of Guernieville, Calif. 3 p.m. Nov. 6, 1949 one heard in the heart of the redwoods, and later seen at a height of about 175 feet in one of them. A few myrtle and small tan oaks nearby,, but all small in stature and nowhere near the elevation utilized by the woodpecker. Nov. 16, 1949 Black Butte hes. Area, Stony Creek, Tehama Co. Cormon in White oaks and Fremonts cottonwoods. Many nest holes seen in cottonwoods. Noisy till about 10 a.m., one much quieter later on. Also observed them hanging on the vertical cliffs of Black Butte, and entering the vertical crevices. This was also observed on the Urland Buttes.

Dec. 30, 1949 hogan Ros. Area, Valaveras Co. Calif. took ics of white and mainly blue scorn scorace in white oak and fence posts. The

Jan. 4, 1950 Black Butte Res. area, Tehama Co. Stony Creek. 40-5./s. mi. These biras were still very active at 830 p.m., flying, calling, perched on posts and

poles, etc., even though nearly dark. august 13, 1950

Philo Mendoreno Co Calif - 2.5 mi NW of Philo, Calif.

Shelaon, Sacto. Calif. 3 pairs with new nest holes in 1 pole.

I have over the divertically about 6'0 feel Touristics. May 8, 1951 april 23, 1952 is mentioned, but not interpreted in Best's Vol. 174, p. 219.

Armstrong Grove state Tark, alif. coveral in reducid-tan or careo. Wigh in Oct.3, 1953

Patific Grove, Walif. Several in rine woods. - some live oak present.

Acres 1984

Observer Fred Lvenden, Jr. Late Location on ob exvitions Payzant's to colors & but may of the mally into and in mag loca Ly 12, 1940 field that make no vigitable, or cover on it as any eggs. They was probably the rest insuces. oochum: or courn: On the first indicate the bound of the most before the constant of the July 17, 1941 ter to a state children in the contract of the ocueumnon where $p^{m_{ij}}$ is the result of the color of the first of the second of the secon agust 5, lu41 Like 100011 volume, as sequenced to star and a second of the alseal faw one gair of them in a dead tree along the survey. Feb. 21, 1943 Philomath to Corvallis Airport. Saw our first one of the year, perched April 26, 1946 on a fence post near the road. Corv Mi., " Cama, IMO a. . e of them lains at f - . . That tober 10, 1146 to the cost, some bring over a u-block disk. Oak Grove south of Philomath: estimated 15-20 in this grove, for they were Jan. 21, 1947, seen everywhere. Many were more or less paired off and giving their "whistle" call notes, as they gracefully chased each other from tree to tree Seemed almost to be courtship flights, being paired off. One Lewis dominated a Varied Thrush perched in the top of an oak, by taking the perch from it. Stewart Lake, Benton Co: 2 of them in the tops of the oaks around the pond. Feb. 6, 1947 they took off every few seconds in pursuit of insects, and aid mame fancy flying once in a while to catch their food. Oak-fir belt between Beaver Cr., and Bellfountain. a pair were taking over Feb. 18, 1947 a knoll scattered with oaks. They would chase the flackers, capture insects, chase each other, and utter call notes once and a while. Another pair seen in scattered oaks on the valley floor. Aug. 22, 1947 near Bellfountain: Poir flushed from side of graves road where they had been at a small mater despure area by a punche. On- They gong, and the other thew to a pole on which a charmon Falls that per thed and the litter gave buy twice.

Aug. 29, 1947 - Bellifountain-Philosomath Pover a half acren seen. One lies to a tail snag where a Gairdner occuped er was in the top, and di died a tiles before Timally raying in and basising the smalless bits about the parch, sater said at circles once note and alit on the imag. april 4, 1949 Juinda, Jolo County 1/1a. ... Quite a few in the Querus lobata of the area. Nov. 16, 1949 Black Butte Res. Area, Stony Cr, Tehama Co. very common, particularly in the blue oak areas. At least 10 seen along every mile traveled by car.

Jan. 4, 1950 Black Lutte Res. Farea - about 30/sq. ini. blue oak woo bland, and almost as dense in white oak-cottonwood open river bottom along the wont' nork. Sept. 7, 1951 NE of Anderson, Shasta Co. Calif. In rocky-soiled, blue-oak woodland. A good many were seen. Seen in footbills from Jone to Reasure, only; in ElDorale Collect. Cold + daily.
3 mi Nof Crown Landing, Calif. 1 on parts " Dec. 18, 1952 May. 31, 1952 S of inderson, hasta Co. Calif. 2;15 p. .

New anderson, " " " " " " " ook woodland

5/7/12 Btt G Colf ola y 11599F. 2N 3 sum. (Com Jan. 2, 1954 Sept. 23, 1955 Dec. 29, 1437 Apr. 9, 1958 C.S.H. 16 - From Bradshaw Road Sacto. Co. to Delaney Corner, Amador Co. Calif. Apr. 29, 1958 Capay Valley, Yolo Co. Calif. Many pairs along hiway.

July 4, 1959 Carrin Coty, o many to wearing of 47ing may 5 and from we. any 15, 1961 Hat Greek - M. Josophiway CSH 89 Sof U3299ScI - Master Co. Carry 2977, 1961 - One at anteligne Calif - 11am. Loto 6.

Ţ

Observer Brea Lvenden, Jr.

Willer of field true on lich outervie: 85-327-548-921-2220-2242-2204-2346-2357-2478-2463-2326-

Red-nuped adjourner

Sphyrapicus virius nue elis

11.t

Location and our covabions

July 13 1944

Good river, above Ketchum near Easely Hot Springs. Jako Saw a pair of them. Male had red throat patch and crown. Female had grayish throat patch and red crown. Both were mottled whitish gray on the back. They seemed slimmer and smaller than NH Red-breasted Sapsucker. Had black head, with 15 white lines on each side. I located a nest, containing young, in a quaking aspen. It was 7 feet above the ground, and had a southern exposure. The nest cavity was $1\frac{1}{2}$ " in diameter and 3-4" deep, at which point it bent downward: There was another nest cavity, in the same tree, only 6 feet high. It also had a souther y exposure. This appeared to be 2 seasons old, because in another aspen, just 4 feet away, was another nest cavity, that was newer and probably was last years work. It faced No and was 52 feet high. When the parent birds brought food, with me standing 5-6 feet away from the nest, they would fly to the hole in the tree opposite the nesting cavity, and go through actions of pretending to be feeding young, in the old hole. After that, they would casually fly or jump the distance to the nesting cavity. All this time the young were making noises of chirrs and chirps. I saw one adult feeding a stonefly and another time a burrestid beetle about 15" long.

Mar. 20, 1947 Cienega Springs north of Parker Ariz: winter resident. Feeding on dates of the date palm. Hung on line pole, motionless for at least 5 minutes.

May 31,1958 Kings buy grade, Gorgan Co Neverla. Wilcha HYBRIO we find the season of days of the contraction.


Sept 17, how, livery Park. It has come of election visits a cirierent trees within a minutes. The ci tance covered has about the yards. The election and the act each of them. If also observed 5 joing birds, just gitting their additionable age. These young birds were continually bickering and contring to each other, and any stuck aretty close together. If the bent to another tree, this others soon protoped there, also. They would tree to chase each other off the error, and thereby occupy that perch. To one individual somed to dominate the others. If ye is heave way for the other. Desides handing on the trunks, they has perches on horizontal times, or even hung upside down, till a chickage or nutrator, from a cluster of the leaf maple sumaras.

April 12, '47 Woodburn, Pudding R: male aigging nest in live ceuar on east side at 3%'.

Saw another one about 150 feet away, feeding in Big-leaf Waple.

Apr. 12 '47 Butte Creek Grove above Monitor: one drilling a hole at 35 feet on the SE side of a Big-leaf Maple

May 3, 1947 Woodubrn, Fudding R: one male lit on the side of a cedar trunk and a Pine Squirrel scolaed it loudly, and then tried to chase it away.



:

- May 31, 1947 Abiqua Creck Foad, above Abiqua Camp, and below Milk hands They had nest at 18 feet on SF side of 6" 1. Fir snug on the exposed south alope about 125 feet in m the Tileated Toda, ecker nest.
- July 12, 1947 Maclonala Forest, Peavy Ogbin area saw several individuals and then in one tree were a adults, I makes and 5 young binds, playing a ound with each other, and giving noisy calls as I have described previously.
- each other, and giving maisy calls as I have described previously.

 July 24, 1947 Fern Ficse land enother "family" of them doying at und in a dead 2. fir.

 Lept. 18, 1947 Aver, Park, (orver side of the order of the order of the order of the order.
 - June 4, 1949, Avery Pakk, Corvallis, Oregon 68. a.m. nest hole in 14" diameter 1. Fir snag at 12'. 13" dia. hole facing to 1. live maple of cas. Young squeeking, especially when the parents brought 10.0. The nale brought food but dian't enter the hole. He made two visits a nimute apart.

The femile care once, went in and gave the young food, and ther left the nest home carrying a small mass, presumably recall ries.

May 31, 1958 Kingsbury grade, Douglas Co. Nevada - Waterda Hill Street North of the Missing of this arms.

Aug. 17, 1961 Clear Lake, MtHood N.For, Wasco Co. Oregon. Family of 2 ads and 3 immatures working alder snags at edge of lake. Noisy at 7:15 a.m.



July 29 1944

MCCall Icaho, Payette river. Saw what must have been this bird, with a black head and cape, with white stri, es on the face.

August 14, 1946, August 15, 1946

June 21, 1949

Milacat Camp, Mill Cr., Crook Co. .. ilawooo Camp, Ochoco hiway summit, Ameeir Co. Feeding on Aspen trees. Twin Bridger, S. Fk. Humbolot Liver, Llko Co. Hevaus: Jair hop nest in tall aid mine me the bridges at a height of objut "8" on the north sign

or the tree. July 19, 1958 E of Velma 7. Pasin, Widorado Co. Calif. 8000 el. Mouns in nest on the south side of lite line at 12 ht. 11 diam. hole entrance. Moism. and the second of the second o

Location and op 17 si na

August 7, 1946 Apr. 5, 1947

Camp Pioneer, Linn Col - I banded one adult this a.m. 7 41-284295. Leadbetters Penin. Willamett-Columbia F bottoms. Flushed an adult (sex??) from a newly constructed nest cavity in a barkless, 15' cottonwood snag (about 20" in diameter). The nest hole was at $7\frac{1}{2}$ feet on the NNE side. 1 3/4 inches in diameter. The hole went in about 2" before grop, ing into the "gourd" part of the nest. Lepth horizonally from the entrance to the back of the cavity was a out 75". The cavity was straight walled from there to the bottom. The adult flew to a cottonwood some 40 feet away, and when we retired to watch the nest, it did not fly directly back, but flew

by the nest snag and over to another cottonwood some 100 feet away. Corvallis: MAcDonala Forest? I male working a 5" dead fir snag in the heart Apr. 24, 1947 of the forest. He was very busy "sounding" the snag and we approached pretty close to him. He pecked everywhere, and when he located a qua spot he fed there, presumably.

Tombstone Prairie, Santiam Pass 2 a pair couring on the side of a Sitka May 16, 1947 Spruce snag in the wolds.

May 22, 1947 Noti Bog, Lane Co., a pair had a nest in a dead 7" D. Fir snag, in a secondgrowth open area near where we ate our lunch. It was on the NE side at 22 feet. There were evidently young in the nest, for both birds were busy coming and going from the nest hole.

This call be been made to beet attached to grant this talks July 26, 1946 the althoroughly is the next of the training for the amelian conduction of the condu emberson in the ..., to other reconstructions. To be an important

July 23-30,1952 Patterson G.S., Modoc N.F., 7,200', Modoc Co. Calif. 2 ad. and 1 young. The young often perched on small twigs like other birds, not clinging apr.19,1440to big limb and brunk bark.

11 - 5 a \$50' in " - " ap1.24, 1943

June 19-20,1954 Silver Lake, Amador Co. Calif. 7200' el. Pair with nest in 28" dia. lodgepole snag over water at 18 (snag in water). Hole facing east. Watched hole from 3:59 to 4:25. Snag in shade during entire time. BB from kids gut hit snag and bird came out at 4:04. Worked lazily 2 trees and in at 4:08 and stayed, but came out much later about 4:45 to 6:30 or so.

July 25, 1954 Camp Harvey West, U. Echo Lake, Eldorado Co. valif. Several pair - Nest in dying aspen at 112 feet. No acults at nest at 1:30. Hole facing southwest. Hole about 10" deep. Aspen in thicket. 2 young in nest. Bunded male, but female flew before I could band it. First flight downward to ground just 40' away. Second flight about 100' to 6' on pine, and third flight across end of lake, a distance of at least 4 mile. All these flights within 3 minutes of nest departure. This at 1:30 p.m. on June 25. At 9 a.m. on June 27, we noted a pair of Tree Swallows apparently taking overthe nest hole, as they went in and out of it while we watched.

Ashen Grove, Wof Caller Teaf Take, Chist. nest in Wiside of Ordian. dying ashen at 5) feet, and only 15' from Tree Cwallow nest. June 1, 1958

Michigan Bas, Calif. One on lever plat along C. 5. H. 16 in relatively open country. Forwards, much,



527-1201-1904-

| | • . | · | |
|--|-----|---|---|
| | | | • |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

Observer Free vencen, or.

Latchelder's occupance

(84)-85-262-264-277-287-2020For--2226-2226-2226-

injobates outs and extreme

Late Locavion am persit vile.

944-951-1006-1070-1072-1099-1108-1112-1129-1132-1182-1190-1210-1372-1419-1478-1632-1649-1666-1667-1673-1690-1703-1793973-1889-1904-1918-1935-1955-1971-1988-2061-2084-2295-2295-2299-2434-24576-2573-2581-2671-

April 24, 1949 : ranklin Fanch Sloughs, F of Courtland, Sa to Colorest in 14' cotton-wood snag at 11' on fiside. Unshadeo.

Natomas, Sacto Co mest hole at 15' on "side of dead alder, just straggling pieces of burn on grayish wood. Home under stight projection, so that when

adult inside pokes head out of hole, it isn't noticeable.
Nigger Hill, Folson Res. area, Eldorabolo, Calf- on digger pine in woodland area - 32 g.m.

June 22, 1961, Woodland Hiway, Yolo Co. Calif. 2 adults and 1/2 young.

in the constant of the state of the



Location and objections

ember 7, 1940

Iver, Prin Corversio: Processiale Caraction of plum recurs committee.

April 12, 1947

Woodburn, Pudding R. Pond: "saw a female working on a nest hole in a willow in the water on the south side at 28 feet (I heard another working noislly off to the NV of this tree). Soon she gave a call note and flew away to the NW and the male came from the NW and started working where the female left off. Very little noise was made here, but the wood pieces flew anyway. The female then worked noisily on the tree the male had been on. The first cavity was already about 25" deep and they were starting to "turn it downward". The second tree was a willow 170 feet NM, also still in the water. The hole was at 25 feet on the N side. It was about 1" deep. This second willow was aead.

Apr. 12 '47

Abiqua Camp Milk Ranch Ridge: one nest building, but I didn't locate it.

May 3, 1947

Woodburn, Dudding R Wusing hole described above on Apr. 12th.

flee: Coming to on 5-14-21 6 eggs et 15' in head agale tree, lake 14" Creg. 5-15-21 5 " " 8' namall head aller this, " 12" ". " 30' " large " " " 5-15-21 5 1. 8 10' n 11 n n J. 11-21 5 1 5-28-21 4 11 11 7 11 another 11 11 6-4-21 4 11 11 31 11 11 11 11 5-16-22 4 11 1 151 11 11 " Coul 5 11 11 15 11

. . .

•

Dendrocopos borealis

Red-cockaded Joodpecker

| | | · | • |
|--|--|---|---|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

Dryobates scalaris ssp.

- March 18, 1947, Bill Williams R. Canyon, Ariz: 4 (2M & 2F). Quite bright crown on the male. This bird has typical Dryobates habits, flight and notes.
- March 20, 1947 Cienega Springs north of Parker, Ariz: 1 male. Nest in Palo Verde stub in a live thicket of the same. SE exposure at 11 feet.
- Feb. 17, 1951 Colo. R. Indian Agency, Parker, Arizona 530 p.m. one feeding inside base of dense Atriplex bush in poor light, at base of SW slope hill. Calling all the time. Back and wing barring distinct.



Nuttall Woodpecker

- February 6, 1949 Gray Loage Refuge about noon. One female in small willow along aitch in marsh area.
- Warch 26, 1949 Sacto F., near Knight's remy 24 to.m. saw at least 6 at one time, and several other. They were working primarily on cottonwood and whilew trunks, and several were observed feeding in the topmost branches on the buds of the cottonwoods, which have not yet opened. Communicated with each other, too. Common.
- Mar. 21, 1950 Bidwell Park, Chico, Calif. 130 p.m. observed a couple working the upper twigs of large white oaks and sycammores, about 60-75' above the ground. They called now and then. Since I have always seen them working trees and bushes close to the ground, this ob is of interest.
- May 9, 1951 Garden Hiway, No of Sacramento Calif. at 930- found nest hole underside of a 5"-diameter cottonwood limb, on W side, 33 feet above Sacto. R. 2 adults were carrying food to the nest; once one returned to nest hole before the other adult had left, so a short note was given, and the late departer, left.
- Dec. 2., 1954 Parts, old c. Thatford bediever bracket flurright W. in cottonwood. June 22, 1961, Woodland Wiway, Yolo Co.Calif. 4 adults and 3 young.



| Liuriber | Oi | Little true | 10 | District | 10 | District

iryobates ulbolarvaur albelarv tur

Late location am objety tion

Aug. 12, 1946

Indian Ford, leschutes Collonce or tice every day, a pair of adults and their young would come down to Indian ford Creek to "Mater". They'd hang on a quaking aspen at the creek's edge and then worktheir way down to the mater, dipping in their bills, or set on the ground at the edge of the creek, to drink.

June 18, 1950

Lake Tahoe, wert side, bloorado and Placer Con. Calir. 130 p.: 128 3 today. One was a female lovering over the hivey, catching insects as a "billers" in mid-dir, and so is stopled to watch her. He seem observed that she was carrying rood to readily-heard joung. The nest hole was a new observed that she was carrying rood to readily-heard joung. The nest hole was a new observed to be about 1," in diameter, a little higher from top to bottom. The hole was on the south side of a lo-Toot red fir stump, at a height of about 11 feet. The stump was full-larked, with no bare wood showing. The stump was only about 15 feet from the west, lakeside havey (C.C.H. 82) and was used to support a neon sign advertising Polit 's Taboe Park Cottages. The make bird was not seen during a period of 15 minutes we ware there, but the remain visited the nest at least 15 times during the 15 minutes, even the globy presence held her away for at least 8 minutes of trut time.

Sept. 9, 1951 Manzanita I, Lassen N.P. Calif. About 9 a.m. I watched one fly out of a tall Jeffrey Pine and chase a sharp-shinned hawk, even the the hawk had not come near the tree top where the woodpecker was working.

May 30,1955 7-Sp.m. irrentes, Al Palpe, Calif. one drum od quite steadilm on heavy snag in forest, at care of clearing, for one hour or sc.

July 9, 1955

Silver Fork, and just left 2 nexts along the stream, one mile apart.

Nay 16-21.,1% is northly the letter refer to a set of the contribution of the latter of the state of the contribution of the state of the set of the se

June 16, 1960 / mi E of Pollock Pines USUR, Elderado & Coff - 111 100m



167-168-4 723-1694-

Jewic -t

The state of the s

ru. 7, Ira signation to the time to the same and the same



ricoia la at . . ci t :

| :.te | · a.t. · · · · · · · · · · · · · · · · · · |
|--------------|--|
| .uc. 7, 10-1 | |
| | Lama of the term o |
| | In the last of the control of the co |
| Apr 25 1944 | Sum it of Bennet Mtn., Mtn. Home, Idaho. A very noisy pair trying to sing, or something - flew from a clump of firs as we approached. |
| Aug. 7, 1940 | Camp Pioneer, Linn Co. Plight in camp early in the mornings. They were working P. contorta trunks this a.m. |















